

TINKER

Software Tools for Molecular Design
Version 3.9
June 2001

Copyright © 1990-2001 by Jay William Ponder
All Rights Reserved

Copyright © 1990-2001 by Jay William Ponder
All Rights Reserved

User's Guide Cover Illustration by Jay Nelson
Courtesy of Prof. R. T. Paine, Univ. of New Mexico

TINKER IS PROVIDED "AS IS" AND WITHOUT ANY WARRANTY
EXPRESS OR IMPLIED. THE USER ASSUMES ALL RISKS OF
USING THIS SOFTWARE. THERE IS NO CLAIM OF THE
MERCHANTABILITY OR FITNESS FOR A PARTICULAR
PURPOSE.

YOU MAY MAKE COPIES OF TINKER FOR YOUR OWN USE,
AND MODIFY THOSE COPIES. YOU MAY NOT DISTRIBUTE ANY
MODIFIED SOURCE CODE OR DOCUMENTATION TO USERS AT
ANY SITE OTHER THAN YOUR OWN.

v3.9 6/01

TINKER

Software Tools for Molecular Design
Version 3.9 June 2001

| <u>Table of Contents</u> | <u>Page</u> |
|--------------------------------------|--------------------|
| Introduction to the TINKER Package | 5 |
| Installing TINKER on your Computer | 7 |
| Types of Input & Output Files | 9 |
| Potential Energy Programs | 12 |
| Structure Manipulation Programs | 18 |
| Force Field Parameter Sets | 22 |
| Use of the Keyword Control File | 29 |
| Notes on Special Features & Methods | 59 |
| Descriptions of TINKER Routines | 65 |
| Contents of Common Block Variables | 124 |
| Index of Function & Subroutine Calls | 150 |
| Examples using the TINKER Package | 174 |
| Benchmark Results | 176 |
| Collaborators & Acknowledgments | 179 |
| References & Suggested Reading | 181 |

1. Introduction to the TINKER Package

Welcome to the TINKER molecular modeling package! TINKER is designed to be an easily used and flexible system of programs and routines for molecular mechanics and dynamics as well as other energy-based and structural manipulation calculations. It is intended to be modular enough to enable development of new computational methods and efficient enough to meet most production calculation needs. Rather than incorporating all the functionality in one monolithic program, TINKER provides a set of relatively small programs that interoperate to perform complex computations. New programs can be easily added by modelers with only limited programming experience. The series of major programs included in the distribution system perform the following core tasks:

- (1) build protein and nucleic acid models from sequence
- (2) energy minimization and structural optimization
- (3) analysis of energy distribution within a structure
- (4) molecular dynamics and stochastic dynamics
- (5) simulated annealing with a choice of cooling schedules
- (6) normal modes and vibrational frequencies
- (7) conformational search and global optimization
- (8) transition state location and conformational pathways
- (9) fitting of energy parameters to crystal data
- (10) distance geometry with pairwise metrization
- (11) molecular volumes and surface areas
- (12) free energy changes for structural mutations
- (13) advanced algorithms based on potential smoothing

Many of the various energy minimization and molecular dynamics computations can be performed on full or partial structures, over Cartesian, internal or rigid body coordinates, and including a variety of boundary conditions and crystal cell types. Other programs are available to generate timing data and allow checking of potential function derivatives for coding errors. Special features are available to facilitate input and output of protein and nucleic acid structures. However, the basic core routines have no knowledge of biopolymer structure and can be used for general molecular systems.

Due to its emphasis on ease of modification, TINKER differs from many other currently available molecular modeling packages in that the user is expected to be willing to write simple "front-end" programs and make some alterations at the source code level. The main programs provided should be considered as templates for the users to change according to their wishes. All subroutines are internally documented and structured programming practices are adhered to throughout. The result, it is hoped, will be a calculational system which can be tailored to local needs and desires.

The core TINKER system consists of nearly 120,000 lines of source written entirely in a portable Fortran77 superset. Use is made of only some very common extensions that aid in writing highly structured code. The current version of the package has been ported to a wide range of computers with no or extremely minimal changes. Tested machines include: Compaq Alphas under Tru64 Unix or OpenVMS; Hewlett-Packard, IBM RS/6000, Silicon Graphics and Sun workstations under the vendor's Unix; Apple Macintosh; and Intel PCs under Windows9X/NT/2000 and Linux. At present, our new code is written on Athlon Linux and Compaq Alpha platforms, and occasionally tested for compatibility on various of the other machine and OS combinations listed above. At present, we are in the process of

converting our primary development efforts from Fortran77 to a more modern Fortran dialect. A machine-translated C version of TINKER is currently available, and a hand-translated optimized C version of a previous TINKER version is available for inspection. Conversion to C or C++ is under consideration, but not being actively pursued at this time.

The basic design of the energy function engine used by the TINKER system allows usage of several different parameter sets. At present we are distributing parameters that implement AMBER-95, CHARMM27, MM2, MM3, OPLS-AA, OPLS-UA and our own TINKER parameters. In most cases, the source code separates the geometric manipulations needed for energy derivatives from the actual form of the energy function itself. Several other literature parameter sets are being considered for future development (ENCAD, MMFF-94, MM4, UFF, *etc.*), and many of the alternative potential function forms reported in the literature can be implemented directly or after minor code changes.

Much of the software in the TINKER package has been heavily used and well tested, but some modules are still in a fairly early stage of development. Further work on the TINKER system is planned in three main areas: (1) extension and improvement of the potential energy parameters including further development of our polarizable multipole TINKER force field, (2) coding of new computational algorithms including additional methods for free energy determination, torsional Monte Carlo and molecular dynamics sampling, advanced methods for long range interactions, better transition state location, and further application of the potential smoothing paradigm, (3) a friendlier user interface for protein/nucleic acid/polysaccharide computations including direct input/output of Protein Data Bank files, and (4) a Java-based GUI front-end to tie together the TINKER programs and provide for basic visualization.

Questions and comments regarding the TINKER package, including suggestions for improvements and changes should be made to the author:

Professor Jay William Ponder
Biochemistry & Molecular Biophysics
North Building, Room 2811, Box 8231
Washington University School of Medicine
660 South Euclid Avenue
Saint Louis, MO 63110 U.S.A.

phone: (314) 362-4195
fax: (314) 362-7183
email: ponder@dasher.wustl.edu

In addition, an Internet web site containing an online version of this User's Guide, the most recent distribution version of the full TINKER package and other useful information can be found at <http://dasher.wustl.edu/tinker>, the Home Page for the TINKER Molecular Modeling Package.

2. Installing TINKER on your Computer

The TINKER package is distributed on the Internet via either the web site or the anonymous ftp account on **dasher.wustl.edu** with an IP number of 128.252.68.48. This node is an AlphaServer 4100 file server running Compaq Tru64 Unix located in the Ponder lab at Washington University School of Medicine. The package is available via the web and standard browsers from the TINKER home page at <http://dasher.wustl.edu/tinker>. Alternatively TINKER can be downloaded by logging into **dasher.wustl.edu** under anonymous ftp (Username: **anonymous**, Password: "your email address") and downloading the software from the /pub/tinker subdirectory. The complete TINKER distribution as well as individual files can be downloaded from this site.

On **dasher.wustl.edu**, the TINKER package is present as both a compressed Unix tar archive and as a complete set of uncompressed source and data files. Binaries are provided for Intel PCs running Windows 9X/ME/NT/2000, PCs running Linux, and for Apple Power Macintosh running Mac OS 9 or earlier. All of these executables are available in standard compressed formats as individual programs or as complete sets of executables. It is expected that other Unix users and PC users who need specially customized versions, will build binaries for their specific system. Sites with access to the Unix tar, compress and uncompress commands should simply obtain the archive file **tinker.tar.Z**. Alternatively, a file **tinker.tar.gz** containing a tar archive compressed with GNU gzip is also provided. If you choose to download individual files, you will need at a minimum the contents of the /doc, /source and /params subdirectories. Also required are the compile/build scripts from the subdirectory named for your machine type. Other areas contain test cases and examples, benchmark results, and machine-translated C code. The entire TINKER package, after building the executables, will require from about 40 to over 100 megabytes of disk space depending on the components installed and the use of shared libraries in the executables.

The documentation for the TINKER programs, including the guide you are currently reading, is located in the /pub/tinker/doc subdirectory. The documentation was prepared using the Applixware Words and Graphics programs. Portable versions of the documentation are provided as ascii text in **.txt** files and in **.ps** Postscript and **.pdf** Adobe Acrobat file formats. Please read and return by mail the TINKER license. While the intent is to distribute the TINKER code to anyone who wants it, the authors would like to keep track of the sites using the package. The returned license forms also help us justify further development of TINKER. When new modules and capabilities become available, and when the almost inevitable bugs are uncovered, we will attempt to notify those who have returned a license form. Finally, we remind you that this software is copyrighted, and ask that it not be redistributed in any form.

The compilation and building of the TINKER executables should be easy for most of the common workstation and PC class computers. We provide in the /make area a Unix-style Makefile that with some modification can be used to build TINKER on most Unix machines. As a simpler alternative to Makefiles for the Unix versions, we also provide machine-specific directories with three separate shell scripts to compile the source, build an object library, and link binary executables. Three similar command files are provided for Open VMS, and for PC and Macintosh systems. Compilation on Unix workstations should use the vendor supplied Fortran compiler, if available. The public domain GNU g77 Fortran compiler available from <http://gcc.gnu.org> is also capable of building TINKER on Unix machines and under Linux. The Linux executables we provide are built with either the Portland Group (PGI) compiler or the Absoft ProFortran compiler, both of which generate somewhat

faster executables than g77. For the Macintosh we have favorable experience with the Absoft ProFortran compiler running under the Macintosh Programmers' Workbench (MPW). On PC's running Windows 9X/NT/2000, the distributed TINKER executables are built on an Intel Pentium III CPU under the Compaq Visual Fortran 6.5 compiler. While the CVF 6.5 compiler has Athlon-specific optimizations, we have not yet investigated these options. The Microsoft Fortran Power Station 4.0 and Watcom F77 compilers are also sufficient for building TINKER, and we provide scripts for each of these PC compilers. Alternative Windows compilers such as those from Lahey/Fujitsu and The Portland Group should work as well, but we have not evaluated them yet. Please see the README files in each of the machine-specific areas for further information.

The first step in building TINKER using the script files is to run the appropriate "compile" script. Next you must use the "library" script to create an archive of object code modules. Finally, run the "link" script to produce the complete set of TINKER executables. The executables can be renamed and moved to wherever you like by editing and running the "rename" script.

Regardless of your target machine, only three small pieces of code can possibly require attention prior to building. The first two are the system dependent time and date routines found in `clock.f` and `calendar.f` respectively. Please uncomment the sections of these routines needed for your computer type. The final set of source alterations are the master array dimensions found in the include file `sizes.i`. The most basic limit is on the number of atoms allowed, "maxatm". This parameter can be set to 10000 or more on most workstations. Personal computers with minimal memory may need a lower limit, perhaps 1000 atoms, depending on available memory, swap space and other resources. A description of the other parameter values is contained in the header of the file. Note that in order to keep the code completely transparent, TINKER does not implement any sort of virtual memory or heap data structure. This requires that `sizes.i` dimensioning values be set at least as large as the biggest problem you intend to run. Obviously, you should not set the array sizes to unnecessarily large values, since this can tax your compute resources and may result in performance degradation. The worst case we know of at present is for some of the Compaq Alpha-based machines, where running a "small" problem with TINKER executables dimensioned to "large" sizes can result in a 25-50% CPU time penalty, especially if only the default compiler options are used.

Specific questions about the building or use of the TINKER package should be directed to ponder@dasher.wustl.edu. TINKER related questions or comments of more general interest can be sent to the Computational Chemistry List (<http://www.ccl.net>) run by Jan Labanowski of The Ohio Supercomputer Center. The TINKER developers monitor this list and will respond to the list or the individual poster as appropriate.

3. Types of Input & Output Files

This section describes the basic file types used by the TINKER package. Let's say you wish to perform a calculation on a particular small organic molecule. Assume that the file name chosen for our input and output files is `sample`. Then all of the TINKER files will reside on the computer under the name `sample.xxx` where `.xxx` is any of the several extension types to be described below.

SAMPLE.XYZ

The `.xyz` file is the basic TINKER Cartesian coordinates file type. It contains a title line followed by one line for each atom in the structure. Each line contains: the sequential number within the structure, an atomic symbol or name, X-, Y-, and Z-coordinates, the force field atom type number of the atom, and a list of the atoms connected to the current atom. Except for programs whose basic operation is in torsional space, all TINKER calculations are done from some version of the `.xyz` format.

SAMPLE.INT

The `.int` file contains an internal coordinates representation of the molecular structure. It consists of a title line followed by one line for each atom in the structure. Each line contains: the sequential number within the structure, an atomic symbol or name, the force field atom type number of the atom, and internal coordinates in the usual Z-matrix format. For each atom the internal coordinates consist of a distance to some previously defined atom, and either two bond angles or a bond angle and a dihedral angle to previous atoms. The length, angle and dihedral definitions do not have to represent real bonded interactions. Following the last atom definition are two optional blank line separated sets of atom number pairs. The first list contains pairs of atoms that are covalently bonded, but whose bond length was not used as part of the atom definitions. These pairs are typically used to close ring structures. The second list contains "bonds" that are to be broken, *i.e.*, pairs of atoms that are not covalently bonded, but which were used to define a distance in the atom definitions.

SAMPLE.KEY

The keyword parameter file always has the extension `.key` and is optionally present during TINKER calculations. It contains values for any of a wide variety of switches and parameters that are used to change the course of the computation from the default. The detailed contents of this file is explained in a latter section of this User's Guide. If a molecular system specific keyfile, in this case `sample.key`, is not present, the the TINKER program will look in the same directory for a generic file named `tinker.key`.

SAMPLE.DYN

The `.dyn` file contains values needed to restart a molecular or stochastic dynamics computation. It stores the current position, current velocity and current and previous accelerations for each atom, as well as the size and shape of any periodic box or crystal unit cell. This information can be used to start a new dynamics run from the final state of a previous run. Upon startup, the dynamics programs always check for the presence of a `.dyn` file and make use of it whenever possible. The `.dyn` file is updated concurrent with the saving of a new dynamics trajectory snapshot.

SAMPLE.END

The `.end` file type provides a mechanism to gracefully stop a running TINKER calculation. At appropriate checkpoints during a calculation, TINKER will test for the presence of a `sample.end` file, and if found will terminate the calculation after updating the output. The `.end` file can be created at any time during a computation, and will be detected when the next checkpoint is reached. The file may be of zero size, and its contents are unimportant. In the current version of TINKER, the `.end` mechanism is only available within dynamics-based programs.

SAMPLE.001, SAMPLE.002,

Several types of computations produce files containing a three or more digit extension (`.001` as shown; or `.002`, `.137`, `.5678`, *etc.*). These are referred to as cycle files, and are used to store various types of output structures. The cycle files from a given computation are identical in internal structure to either the `.xyz` or `.int` files described above. For example, the vibrational analysis program can save the tenth normal mode in `sample.010`. A molecular dynamics-based program might save its tenth 0.1 picosecond frame (or an energy minimizer its tenth partially minimized intermediate) in a file of the same name.

SAMPLE.ARC

A TINKER archive file is simply a series of `.xyz` Cartesian coordinate files appended together one after another. This file can be used to condense the results from intermediate stages of an optimization, frames from a molecular dynamics trajectory, or set of normal mode vibrations into a single file for storage.

SAMPLE.PDB

This file type contains coordinate information in the PDB format developed by the Brookhaven Protein Data Bank for deposition of model structures based on macromolecular X-ray diffraction and NMR data. Although TINKER itself does not use `.pdb` files directly for input/output, auxiliary programs are provided with the system for interconverting `.pdb` files with the `.xyz` format described above.

SAMPLE.SEQ

This file type contains the primary sequence of a biopolymer in the standard one-letter code with 50 residues per line. The `.seq` file for a biopolymer is generated automatically when a PDB file is converted to TINKER `.xyz` format or when using the PROTEIN or NUCLEIC programs to build a structure from sequence. It is required for the reverse conversion of a TINKER file back to PDB format.

SAMPLE.FRAC

The fractional coordinates corresponding to the asymmetric unit of a crystal unit cell are stored in the `.frac` file. The internal format of this file is identical to the `.xyz` file; except that the coordinates are fractional instead of in Angstrom units.

SAMPLE.XMOL

The ARCHIVE program has the option of converting a series of **.xyz** cycle files into an XMakeMol XYZ file. These files can be displayed as a movie using the XMakeMol display program. Note that the **.xmol** file format does not contain TINKER atom type information, so it is not possible to convert an **.xmol** file back into a TINKER **.xyz** file.

SAMPLE.MSI

The ARCHIVE program has the option of converting a series of **.xyz** cycle files into a MSI InsightII coordinate archive file. These files can be displayed as a movie using the InsightII display program. Note that the **.msi** file format does not contain TINKER atom type information, so it is not possible to convert a **.msi** file back into a TINKER **.xyz** file.

PARAMETER FILES

The potential energy parameter files distributed with the TINKER package all end in the extension **.prm**, although this is not required by the programs themselves. Each of these files contains a definition of the potential energy functional forms for that force field as well as values for individual energy parameters. For example, the **mm3pro.prm** file contains the energy parameters and definitions needed for a protein-specific version of the MM3 force field.

4. Potential Energy Programs

This section of the manual contains a brief description of each of the TINKER potential energy programs. A detailed example showing how to run each program is included in a later section. The programs listed below are all part of the main, supported distribution. Additional source code for various unsupported programs can be found in the /other directory of the TINKER distribution.

ALCHEMY

A simple program to perform very basic free energy perturbation calculations. This program is provided mostly for demonstration purposes. For example, we use ALCHEMY in a molecular modeling course laboratory exercise to perform such classic mutations as chloride to bromide and ethane to methanol in water. The present version uses the perturbation formula and windowing with an explicit mapping of atoms involved in the mutation ("AMBER"-style), instead of thermodynamic integration and independent freely propagating groups of mutated atoms ("CHARMM"-style). Some of the code specific to this program is limited to the AMBER and OPLS potential functional forms, but could be easily generalized to handle other potentials. A more general and sophisticated version is currently under development.

ANALYZE

Provides information about a specific molecular structure. The program will ask for the name of a structure file, which must be in the TINKER `.xyz` file format, and the type of analysis desired. Options allow output of: (1) total potential energy of the system, (2) breakdown of the energy by potential function type or over individual atoms, (3) computation of the total dipole moment and its components, moments of inertia and radius of gyration, (4) listing of the parameters used to compute selected interaction energies, (5) energies associated with specified individual interactions.

ANNEAL

Performs a molecular dynamics simulated annealing computation. The program starts from a specified input molecular structure in TINKER `.xyz` format. The trajectory is updated using either a modified Beeman or a velocity Verlet integration method. The annealing protocol is implemented by allowing smooth changes between starting and final values of the system temperature via the Groningen method of coupling to an external bath. The scaling can be linear or sigmoidal in nature. In addition, parameters such as cutoff distance can be transformed along with the temperature. The user must input the desired number of dynamics steps for both the equilibration and cooling phases, a time interval for the dynamics steps, and an interval between coordinate/trajectory saves. All saved coordinate sets along the trajectory are placed in sequentially numbered cycle files.

DYNAMIC

Performs a molecular dynamics (MD) or stochastic dynamics (SD) computation. Starts either from a specified input molecular structure (an `.xyz` file) or from a structure-velocity-acceleration set saved from a previous dynamics trajectory (a restart from a `.dyn` file). MD trajectories are propagated using either a modified Beeman or a velocity Verlet integration method. SD is implemented via our own derivation of a velocity Verlet-based algorithm. In

addition the program can perform full crystal calculations, and can operate in constant energy mode or with maintenance of a desired temperature and/or pressure using the Groningen method of coupling to external baths. The user must input the desired number of dynamics steps, a time interval for the dynamics steps, and an interval between coordinate/trajectory saves. Coordinate sets along the trajectory can be saved as sequentially numbered cycle files or directly to a TINKER archive `.arc` file. At the same time that a point along the trajectory is saved, the complete information needed to restart the trajectory from that point is updated and stored in the `.dyn` file.

GDA

A program to implement Straub's Gaussian Density Annealing algorithm over an effective series of analytically smoothed potential energy surfaces. This method can be viewed as an extended stochastic version of the diffusion equation method of Scheraga, *et al.*, and also has many similar features to the TINKER Potential Smoothing and Search (PSS) series of programs. The current version of GDA is similar to but does not exactly reproduce Straub's published method and is limited to argon clusters and other simple systems involving only van der Waals interactions; further modification and development of this code is currently underway in the Ponder research group. As with other programs involving potential smoothing, GDA currently requires use of the `smooth.prm` force field parameters.

MINIMIZE

The MINIMIZE program performs a limited memory L-BFGS minimization of an input structure over Cartesian coordinates using a modified version of the algorithm of Jorge Nocedal. The method requires only the potential energy and gradient at each step along the minimization pathway. It requires storage space proportional to the number of atoms in the structure. The MINIMIZE procedure is recommended for preliminary minimization of trial structures to an *rms* gradient of 1.0 to 0.1 kcal/mole/Å. It has a relatively fast cycle time and is tolerant of poor initial structures, but converges in a slow, linear fashion near the minimum. The user supplies the name of the TINKER `.xyz` coordinates file and a target *rms* gradient value at which the minimization will terminate. Output consists of minimization statistics written to the screen or redirected to an output file, and the new coordinates written to updated `.xyz` files or to cycle files.

MINIROT

The MINIROT program uses the same limited memory L-BFGS method as MINIMIZE, but performs the computation in terms of dihedral angles instead of Cartesian coordinates. Output is saved in an updated `.int` file or in cycle files.

MINRIGID

The MINRIGID program is similar to MINIMIZE except that it operates on rigid bodies starting from a TINKER `.xyz` coordinate file and the rigid body group definitions found in the corresponding `.key` file. Output is saved in an updated `.xyz` file or in cycle files.

NEWTON

A truncated Newton minimization method which requires potential energy, gradient and Hessian information. This procedure has significant advantages over standard Newton methods, and is able to minimize very large structures completely. Several options are

provided with respect to minimization method and preconditioning of the Newton equations. The default options are recommended unless the user is familiar with the math involved. This program operates in Cartesian coordinate space and is fairly tolerant of poor input structures. Typical algorithm iteration times are longer than with nonlinear conjugate gradient or variable metric methods, but many fewer iterations are required for complete minimization. NEWTON is usually the best choice for minimizations to the 0.01 to 0.000001 kcal/mole/Å level of *rms* gradient convergence. Tests for directions of negative curvature can be removed, allowing NEWTON to be used for optimization to conformational transition state structures (this only works if the starting point is very close to the transition state). Input consists of a TINKER *.xyz* coordinates file; output is an updated set of minimized coordinates and minimization statistics.

NEWTROT

The NEWTROT program is similar to NEWTON except that it requires a *.int* file as input and then operates in terms of dihedral angles as the minimization variables. Since the dihedral space Hessian matrix of an arbitrary structure is often indefinite, this method will often not perform as well as the other, simpler dihedral angle based minimizers.

OPTIMIZE

The OPTIMIZE program performs a optimally conditioned variable metric minimization of an input structure over Cartesian coordinates using an algorithm due to William Davidon. The method does not perform line searches, but requires computation of energies and gradients as well as storage for an estimate of the inverse Hessian matrix. The program operates on Cartesian coordinates from a TINKER *.xyz* file. OPTIMIZE will typically converge somewhat faster and more completely than MINIMIZE. However, the need to store and manipulate a full inverse Hessian estimate limits its use to structures containing less than a few hundred atoms on workstation class machines. As with the other minimizers, OPTIMIZE needs input coordinates and an *rms* gradient cutoff criterion. The output coordinates are saved in updated *.xyz* files or as cycle files.

OPTIROT

The OPTIROT program is similar to OPTIMIZE except that it operates on dihedral angles starting from a TINKER *.int* internal coordinate file. This program is usually the preferred method for most dihedral angle optimization problems since Truncated Newton methods appear, in our hands, to lose some of their efficacy in moving from Cartesian to torsional coordinates.

OPTRIGID

The OPTRIGID program is similar to OPTIMIZE except that it operates on rigid bodies starting from a TINKER *.xyz* coordinate file and the rigid body atom group definitions found in the corresponding *.key* file. Output is saved in an updated *.xyz* file or in cycle files.

PATH

A program that implements a variant of Elber's Lagrangian multiplier-based reaction path following algorithm. The program takes as input a pair of structural minima as TINKER *.xyz* files, and then generates a user specified number of points along a path through

conformational space connecting the input structures. The intermediate structures are output as TINKER cycle files, and the higher energy intermediates can be used as input to a Newton-based optimization to locate conformational transition states.

PSS

Implements our version of a potential smoothing and search algorithm for the global optimization of molecular conformation. An initial structure in **.xyz** format is first minimized in Cartesian coordinates on a series of increasingly smoothed potential energy surfaces. Then the smoothing procedure is reversed with minimization on each successive surface starting from the coordinates of the minimum on the previous surface. A local search procedure is used during the backtracking to explore for alternative minima better than the one found during the current minimization. The final result is usually a very low energy conformation or, in favorable cases, the global energy minimum conformation. The minimum energy coordinate sets found on each surface during both the forward smoothing and backtracking procedures are placed in sequentially numbered cycle files.

PSSRIGID

This program implements the potential smoothing and search method as described above for the PSS program, but performs the computation in terms of keyfile-defined rigid body atom groups instead of Cartesian coordinates. Output is saved in numbered cycle files with the **.xyz** file format.

PSSROT

This program implements the potential smoothing and search method as described above for the PSS program, but performs the computation in terms of a set of user-specified dihedral angles instead of Cartesian coordinates. Output is saved in numbered cycle files with the **.int** file format.

SADDLE

A program for the location of a conformational transition state between two potential energy minima. SADDLE uses a conglomeration of ideas from the Bell-Crighton quadratic path and the Halgren-Lipscomb synchronous transit methods. The basic idea is to perform a nonlinear conjugate gradient optimization in a subspace orthogonal to a suitably defined reaction coordinate. The program requires as input the coordinates (TINKER **.xyz** files) of the two minima and an *rms* gradient convergence criterion for the optimization. The current estimate of the transition state structure is written to the file TSTATE.XYZ. Crude transition state structures generated by SADDLE can sometimes be refined using the NEWTON program. Optionally, a scan of the interconversion pathway can be made at each major iteration.

SCAN

A program for general conformational search of an entire potential energy surface via a basin hopping method. The program takes as input a TINKER **.xyz** coordinates file which is then minimized to find the first local minimum for a search list. A series of activations along various normal modes from this initial minimum are used as seed points for additional minimizations. Whenever a previously unknown local minimum is located it is added to the search list. When all minima on the search list have been subjected to the normal mode

activation without locating additional new minima, the program terminates. The individual local minima are written to cycle files as they are discovered. While the SCAN program can be used on standard undeformed potential energy surfaces, we have found it to be most useful for quickly “scanning” a smoothed energy surface to enumerate the major basins of attraction spanning the entire surface.

SNIFFER

A program that implements the Sniffer global optimization algorithm of Butler and Slaminka, a discrete version of Griewank’s global search trajectory method. The program takes an input TINKER `.xyz` coordinates file and shakes it vigorously via a modified dynamics trajectory before, hopefully, settling into a low lying minimum. Some trial and error is often required as the current implementation is sensitive to various parameters and tolerances that govern the computation. At present, these parameters are not user accessible, and must be altered in the source code. However, this method can do a good job of quickly optimizing conformation within a limited range of convergence.

TESTGRAD

The TESTGRAD program computes and compares the analytical and numerical first derivatives (*i.e.*, the gradient vector) of the potential energy for a Cartesian coordinate input structure. The output can be used to test or debug the current potential or any added user defined energy terms.

TESTHESS

The TESTHESS program computes and compares the analytical and numerical second derivatives (*i.e.*, the Hessian matrix) of the potential energy for a Cartesian coordinate input structure. The output can be used to test or debug the current potential or any added user defined energy terms.

TESTLIGHT

A program to compare the efficiency of different nonbonded neighbor methods for the current molecular system. The program times the computation of energy and gradient for the van der Waals and charge-charge electrostatic potential terms using a simple double loop over all interactions and using the Method of Lights algorithm to select neighbors. The results can be used to decide whether the Method of Lights has any CPU time advantage for the current structure. Both methods should give exactly the same answer in all cases, since the identical individual interactions are computed by both methods. The default double loop method is faster when cutoffs are not used, or when the cutoff sphere contains about half or more of the total system of unit cell. In cases where the cutoff sphere is much smaller than the system size, the Method of Lights can be much faster since it avoids unnecessary calculation of distances beyond the cutoff range.

TESTROT

The TESTROT program computes and compares the analytical and numerical first derivatives (*i.e.*, the gradient vector) of the potential energy with respect to dihedral angles. Input is a TINKER `.int` internal coordinate file. The output can be used to test or debug the current potential functions or any added user defined energy terms.

TIMER

A simple program to provide timing statistics for energy function calls within the TINKER package. TIMER requires an input `.xyz` file and outputs the CPU time (wall clock time on some machine types) needed to perform a specified number of energy, gradient and Hessian evaluations.

TIMEROT

This program is similar to TIMER, only it operates over dihedral angles via input of a TINKER `.int` internal coordinate file. In the current version, the torsional Hessian is computed numerically from the analytical torsional gradient.

VIBRATE

A program to perform vibrational analysis by computing and diagonalizing the full Hessian matrix (*i.e.*, the second partial derivatives) for an input structure (a TINKER `.xyz` file). Eigenvalues and eigenvectors of the mass weighted Hessian (*i.e.*, the vibrational frequencies and normal modes) are also calculated. Structures corresponding to individual normal mode motions can be saved in cycle files.

VIBROT

The program VIBROT forms the torsional Hessian matrix via numerical differentiation of the analytical torsional gradient. The Hessian is then diagonalized and the eigenvalues are output. The present version does not compute the kinetic energy matrix elements needed to convert the Hessian into the torsional normal modes; this will be added in a later version. The required input is a TINKER `.int` internal coordinate file.

XTALFIT

The XTALFIT program is of use in the automated fitting of potential parameters to crystal structure and thermodynamic data. XTALFIT takes as input several crystal structures (TINKER `.xyz` files with unit cell parameters in corresponding keyfiles) as well as information on lattice energies and dipole moments of monomers. The current version uses a nonlinear least squares optimization to fit van der Waals and electrostatic parameters to the input data. Bounds can be placed on the values of the optimization parameters.

XTALMIN

A program to perform full crystal minimizations. The program takes as input the structure coordinates and unit cell lattice parameters. It then alternates cycles of Newton-style optimization of the structure and conjugate gradient optimization of the crystal lattice parameters. This alternating minimization is slower than more direct optimization of all parameters at once, but is somewhat more robust in our hands. The symmetry of the original crystal is not enforced, so interconversion of crystal forms may be observed in some cases.

5. Structure Manipulation Programs

This section of the manual contains a brief description of each of the TINKER structure manipulation, geometric calculation and auxiliary programs. A detailed example showing how to run each program is included in a later section. The programs listed below are all part of the main, supported distribution. Additional source code for various unsupported programs can be found in the /other directory of the TINKER distribution.

ARCHIVE

A program for concatenating TINKER cycle files into a single archive file; useful for storing the intermediate results of minimizations, dynamics trajectories, and so on. The archive file can be written in TINKER format, or in formats usable with MSI's InsightII (their CAR file with **.msi** extension) or with XMakemol (their file format with **.xmo1** extension). Only active atoms are written into the InsightII and XMakemol output files, allowing display of partial structures. The program can also extract individual cycle files from a TINKER archive.

CORRELATE

A program to compute time correlation functions from collections of TINKER cycle files. Its use requires a user supplied function **property** that computes the value of the property for which a time correlation is desired for two input structures. A sample routine is supplied that computes either a velocity autocorrelation function or an *rms* structural superposition as a function of time. The main body of the program organizes the overall computation in an efficient manner and outputs the final time correlation function.

CRYSTAL

A program for the manipulation of crystal structures including interconversion of fractional and Cartesian coordinates, generation of the unit cell from an asymmetric unit, and building of a crystalline block of specified size via replication of a single unit cell. The present version can handle about 25 of the most common space groups, others can easily be added as needed by modification of the routine **symmetry**.

DIFFUSE

A program to compute the self-diffusion constant for a homogeneous liquid via the Einstein equation. A previously saved dynamics trajectory is read in and "unfolded" to reverse translation of molecules due to use of periodic boundary conditions. The average motion over all molecules is then used to compute the self-diffusion constant. While the current program assumes a homogeneous system, it should be easy to modify the code to handle diffusion of individual molecules or other desired effects.

DISTGEOM

A program to perform distance geometry calculations using variations on the classic metric matrix method. A user specified number of structures consistent with keyfile input distance and dihedral restraints is generated. Bond length and angle restraints are derived from the input structure. Trial distances between the triangle smoothed lower and upper bounds can be chosen via any of several metrization methods, including a very effective partial random pairwise scheme. The correct radius of gyration of the structure is automatically maintained

by choosing trial distances from Gaussian distributions of appropriate mean and width. The initial embedded structures can be further refined against a geometric restraint-only potential using either a sequential minimization protocol or simulated annealing.

DOCUMENT

The DOCUMENT program is provided as a minimal listing and documentation tool. It operates on the TINKER source code, either individual files or the complete source listing produced by the command script `listing.make`, to generate lists of routines, common blocks or valid keywords. In addition, the program has the ability to output a formatted parameter listing from the standard TINKER parameter files.

INTEDIT

A program to allow interactive inspection and alteration of the internal coordinate definitions and values of a TINKER structure. If the structure is altered, the user has the option to write out a new internal coordinates file upon exit.

INTXYZ

A program to convert a TINKER `.int` internal coordinates formatted file into a TINKER `.xyz` Cartesian coordinates formatted file.

NUCLEIC

A program for automated building of nucleic acid structures. Upon interactive input of a nucleotide sequence with optional phosphate backbone angles, the program builds internal and Cartesian coordinates. Standard bond lengths and angles are used. Both DNA and RNA sequences are supported as are A-, B- and Z-form structures. Double helices of complementary sequence can be automatically constructed via a rigid docking of individual strands.

PDBXYZ

A program for converting a Brookhaven Protein Data Bank file (a PDB file) into a TINKER `.xyz` Cartesian coordinate file. If the PDB file contains only protein/peptide amino acid residues, then standard protein connectivity is assumed, and transferred to the `.xyz` file. For non-protein portions of the PDB file, atom connectivity is determined by the program based on interatomic distances. The program also has the ability to add or remove hydrogen atoms from a protein as required by the force field specified during the computation.

POLARIZE

A program for computing molecular polarizability from an atom-based distributed model of polarizability. A damped interaction model due to Thole is optionally via keyfile settings. A TINKER `.xyz` file is required as input. The output consists of the overall polarizability tensor in the global coordinates and its eigenvalues.

PROTEIN

A program for automated building of peptide and protein structures. Upon interactive input of an amino acid sequence with optional phi/psi/omega/chi angles, D/L chirality, etc., the

program builds internal and Cartesian coordinates. Standard bond lengths and angles are assumed for the peptide. The program will optionally convert the structure to a cyclic peptide, or add either or both N- and C-terminal capping groups. Atom type numbers are automatically assigned for the specified force field. The final coordinates and a sequence file are produced as the output.

RADIAL

A program to compute the pair radial distribution function between two atom types. The user supplies the two atom names for which the distribution function is to be computed, and the width of the distance bins for data analysis. A previously saved dynamics trajectory is read as input. The raw radial distribution and a spline smoothed version are then output from zero to a distance equal to half the minimum periodic box dimension. The atom names are matched to the atom name column of the TINKER **.xyz** file, independent of atom type.

SPACEFILL

A program to compute the volume and surface areas of molecules. Using a modified version of Connolly's original analytical description of the molecular surface, the program determines either the van der Waals, accessible or molecular (contact/reentrant) volume and surface area. Both surface area and volume are broken down into their geometric components, and surface area is decomposed into the convex contribution for each individual atom. The probe radius is input as a user option, and atomic radii can be set via the keyword file. If TINKER archive files are used as input, the program will compute the volume and surface area of each structure in the input file.

SPECTRUM

A program to compute a power spectrum from velocity autocorrelation data. As input, this program requires a velocity autocorrelation function as produced by the CORRELATE program. This data, along with a user input time step, are Fourier transformed to generate the spectral intensities over a wavelength range. The result is a power spectrum, and the positions of the bands are those predicted for an infrared or Raman spectrum. However, the data is not weighted by molecular dipole moment derivatives as would be required to produce correct IR intensities.

SUPERPOSE

A program to superimpose two molecular structures in 3-dimensions. A variety of options for input of the atom sets to be used during the superposition are presented interactively to the user. The superposition can be mass-weighted if desired, and the coordinates of the second structure superimposed on the first structure are optionally output. If TINKER archive files are used as input, the program will compute all pairwise superpositions between structures in the input files.

SYBYLXYZ

A program for converting a TRIPOS Sybyl MOL2 file into a TINKER **.xyz** Cartesian coordinate file. The current version of the program does not attempt to convert the Sybyl atoms types into the active TINKER force field types, *i.e.*, all atoms types are simply set to zero.

TVIEW

This is a molecule viewing program derived from the well-known RasMol program of Roger Sayle. TVIEW is modified to remove most of the protein-specific options and to directly read the TINKER `.xyz` file format. The original RasMol program has been altered to allow selection and specification by atoms instead of residues. We hope to provide additional functionality in future versions of TVIEW, especially the ability to animate the viewing of sequences of coordinate snapshots from a minimization or dynamic trajectory.

XYZEDIT

A program that performs a variety of manipulations on an input TINKER `.xyz` Cartesian coordinates formatted file. The present version of the program has the following interactively selectable options: (1) Offset the Numbers of the Current Atoms, (2) Deletion of Individual Specified Atoms, (3) Deletion of Specified Types of Atoms, (4) Deletion of Atoms outside Cutoff Range, (5) Insertion of Individual Specified Atoms, (6) Replace Old Atom Type with a New Type, (7) Assign Connectivities based on Distance, (8) Convert Units from Bohrs to Angstroms, (9) Invert thru Origin to give Mirror Image, (10) Translate Center of Mass to the Origin, (11) Translate a Specified Atom to the Origin, (12) Translate and Rotate to Inertial Frame, (13) Move to Specified Rigid Body Coordinates, (14) Create and Fill a Periodic Boundary Box, (15) Soak Current Molecule in Box of Solvent, (16) Append another XYZ file to Current One. In most cases, multiply options can be applied sequentially to an input file. At the end of the editing process, a new version of the original `.xyz` file is written as output.

XYZINT

A program for converting a TINKER `.xyz` Cartesian coordinate formatted file into a TINKER `.int` internal coordinates formatted file.

XYZPDB

A program for converting a TINKER `.xyz` Cartesian coordinate file into a Brookhaven Protein Data Bank file (a PDB file).

XYZSYBYL

A program to convert a TINKER `.xyz` Cartesian coordinates file into a TRIPOS Sybyl MOL2 file. The conversion generates only the MOLECULE, ATOM, BOND and SUBSTRUCTURE record type in the MOL2 file. Generic Sybyl atom types are used in most cases; while these atom types may need to be altered in some cases, Sybyl is usually able to correctly display the resulting MOL2 file.

6. Force Field Parameter Sets

The TINKER package is distributed with several force field parameter sets, implementing a selection of widely used literature force fields as well as the TINKER force field currently under construction in the Ponder lab. We try to exactly reproduce the intent of the original authors of our distributed, third-party force fields. In all cases the parameter sets have been validated against literature reports, results provided by the original developers, or calculations made with the authentic programs. With the few exceptions noted below, TINKER calculations can be treated as authentic results from the genuine force fields. A brief description of each parameter set, including some still in preparation and not distributed with the current version, is provided below with lead literature references for the force field:

AMBER.PRM

AMBER-94/96 parameters for proteins and nucleic acids. Note that with their "Cornell" force field, the Kollman group has devised separate, fully independent partial charge values for each of the N- and C-terminal amino acid residues. At present, the terminal residue charges for TINKER's version maintain the correct formal charge, but redistributed somewhat at the alpha carbon atoms from the Kollman group values. The total magnitude of the redistribution is less than 0.01 electrons in most cases. The file provided with TINKER reproduces the original parm94 set; torsional parameter changes for parm96 are noted in that section of the file. The newer, polarizable parm99 parameter set is not distributed with TINKER at the present time.

W. D. Cornell, P. Cieplak, C. I. Bayly, I. R. Gould, K. M. Merz, Jr., D. M. Ferguson, D. C. Spellmeyer, T. Fox, J. W. Caldwell and P. A. Kollman, A Second Generation Force Field for the Simulation of Proteins, Nucleic Acids, and Organic Molecules, *J. Am. Chem. Soc.*, **117**, 5179-5197 (1995) [PARM94]

P. Kollman, R. Dixon, W. Cornell, T. Fox, C. Chipot and A. Pohorille, The Development/Application of a 'Minimalist' Organic/Biochemical Molecular Mechanic Force Field using a Combination of *ab Initio* Calculations and Experimental Data, in **Computer Simulation of Biomolecular Systems**, W. F. van Gunsteren, P. K. Weiner, A. J. Wilkinson, eds., Volume 3, 83-96 (1997) [PARM96]

G. Moyna, H. J. Williams, R. J. Nachman and A. I. Scott, Conformation in Solution and Dynamics of a Structurally Constrained Linear Insect Kinin Pentapeptide Analogue, *Biopolymers*, **49**, 403-413 (1999) [AIB charges]

W. S. Ross and C. C. Hardin, Ion-Induced Stabilization of the G-DNA Quadruplex: Free Energy Perturbation Studies, *J. Am. Chem. Soc.*, **116**, 4363-4366 (1994) [alkali metal ions]

J. Aqvist, Ion-Water Interaction Potentials Derived from Free Energy Perturbation Simulations, *J. Phys. Chem.*, **94**, 8021-8024, 1990 [alkaline earth Ions, radii adapted for AMBER combining rule]

Current force field parameter values and suggested procedures for development of parameters for additional molecules are available from the AMBER web site at UCSF, <http://www.amber.ucsf.edu/amber/amber.html/>

CHARMM.PRM

CHARMM27 parameters for proteins. Most of the nucleic acid, lipid and small model compound parameters are not yet implemented. We plan to provide the CHARMM27 nucleic acid parameters in due course as a separate parameter file.

N. Foloppe and A. D. MacKerell, Jr., All-Atom Empirical Force Field for Nucleic Acids: 1) Parameter Optimization Based on Small Molecule and Condensed Phase Macromolecular Target Data, *J. Comput. Chem.*, **21**, 86-104 (2000) [CHARMM27]

N. Banavali and A. D. MacKerell, Jr., All-Atom Empirical Force Field for Nucleic Acids: 2) Application to Molecular Dynamics Simulations of DNA and RNA in Solution, *J. Comput. Chem.*, **21**, 105-120 (2000)

A. D. MacKerell, Jr., *et al.*, All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins, *J. Phys. Chem. B*, **102**, 3586-3616 (1998) [CHARMM22]

A. D. MacKerell, Jr., J. Wiorkeiwicz-Kuczera and M. Karplus, An All-Atom Empirical Energy Function for the Simulation of Nucleic Acids, *J. Am. Chem. Soc.*, **117**, 11946-11975 (1995)

S. E. Feller, D. Yin, R. W. Pastor and A. D. MacKerell, Jr., Molecular Dynamics Simulation of Unsaturated Lipids at Low Hydration: Parametrization and Comparison with Diffraction Studies, *Biophysical Journal*, **73**, 2269-2279 (1997) [alkenes]

R. H. Stote and M. Karplus, Zinc Binding in Proteins and Solution - A Simple but Accurate Nonbonded Representation, *Proteins*, **23**, 12-31 (1995) [zinc ion]

Current and legacy parameter values are available from the CHARMM force field web site on Alex MacKerell's Research Interests page at the University of Maryland School of Pharmacy, <https://rxsecure.umaryland.edu/research/amackere/research.html/>

DUDEK.PRM

Protein-only parameters for the early 1990's TINKER force field with multipole values of Dudek and Ponder. The current file contains only the multipole values from the 1995 paper by Dudek and Ponder. This set is now superceded by the more recent TINKER force field developed by Pengyu Ren (see WATER.PRM, below).

M. J. Dudek and J. W. Ponder, Accurate Electrostatic Modelling of the Intramolecular Energy of Proteins, *J. Comput. Chem.*, **16**, 791-816 (1995)

EMR.PRM

Reduced EMR model adapted for flexible sidechains. Only a few amino acid residue types have been implemented.

R. V. Pappu, W. J. Schneller and D. L. Weaver, Electrostatic Multipole Representation of a Polypeptide Chain: An Algorithm for Simulation of Polypeptide Properties, *J. Comput. Chem.*, **17**, 1033-1055 (1996)

ENCAD.PRM

ENCAD parameters for proteins and nucleic acids. (*in preparation*)

M. Levitt, M. Hirshberg, R. Sharon and V. Daggett, Potential Energy Function and Parameters for Simulations of the Molecular Dynamics of Protein and Nucleic Acids in Solution, **Comp. Phys. Commun.**, *91*, 215-231 (1995)

M. Levitt, M. Hirshberg, R. Sharon, K. E. Laidig and V. Daggett, Calibration and Testing of a Water Model for Simulation of the Molecular Dynamics of Protein and Nucleic Acids in Solution, **J. Phys. Chem. B**, *101*, 5051-5061 (1997) [F3C water]

HOCH.PRM

Simple NMR-NOE force field of Hoch and Stern.

J. C. Hoch and A. S. Stern, A Method for Determining Overall Protein Fold from NMR Distance Restraints, **J. Biomol. NMR**, *2*, 535-543 (1992)

MERCK.PRM

Preliminary MMFF vdw parameters using buffered 14/7 function.

T. A. Halgren, Representation of van der Waals (vdW) Interactions in Molecular Mechanics Force Fields: Potential Form, Combination Rules, and vdW Parameters, **J. Am. Chem. Soc.**, *114*, 7827-7843 (1992)

MM2.PRM

Full MM2(1991) parameters including π -systems. The anomeric and electronegativity correction terms included in some later versions of MM2 are not implemented.

N. L. Allinger, Conformational Analysis. 130. MM2. A Hydrocarbon Force Field Utilizing V1 and V2 Torsional Terms, **J. Am. Chem. Soc.**, *99*, 8127-8134 (1977)

J. T. Sprague, J. C. Tai, Y. Yuh and N. L. Allinger, The MMP2 Computational Method, **J. Comput. Chem.**, *8*, 581-603 (1987)

J. C. Tai and N. L. Allinger, Molecular Mechanics Calculations on Conjugated Nitrogen-Containing Heterocycles, **J. Am. Chem. Soc.**, *110*, 2050-2055 (1988)

J. C. Tai, J.-H. Lii and N. L. Allinger, A Molecular Mechanics (MM2) Study of Furan, Thiophene, and Related Compounds, **J. Comput. Chem.**, *10*, 635-647 (1989)

N. L. Allinger, R. A. Kok and M. R. Imam, Hydrogen Bonding in MM2, **J. Comput. Chem.**, *9*, 591-595 (1988)

L. Nørskov-Lauritsen and N. L. Allinger, A Molecular Mechanics Treatment of the Anomeric Effect, **J. Comput. Chem.**, *5*, 326-335 (1984)

All parameters distributed with TINKER are from the "MM2 (1991) Parameter Set", as provided by N. L. Allinger, University of Georgia

MM3.PRM

Full MM3(2000) parameters including pi-systems. The directional hydrogen bonding term and electronegativity bond length corrections are implemented, but the anomeric and Bohlmann correction terms are not implemented.

N. L. Allinger, Y. H. Yuh and J.-H. Lii, Molecular Mechanics. The MM3 Force Field for Hydrocarbons. 1, *J. Am. Chem. Soc.*, *111*, 8551-8566 (1989)

J.-H. Lii and N. L. Allinger, Molecular Mechanics. The MM3 Force Field for Hydrocarbons. 2. Vibrational Frequencies and Thermodynamics, *J. Am. Chem. Soc.*, *111*, 8566-8575 (1989)

J.-H. Lii and N. L. Allinger, Molecular Mechanics. The MM3 Force Field for Hydrocarbons. 3. The van der Waals' Potentials and Crystal Data for Aliphatic and Aromatic Hydrocarbons, *J. Am. Chem. Soc.*, *111*, 8576-8582 (1989)

N. L. Allinger, H. J. Geise, W. Pyckhout, L. A. Paquette and J. C. Gallucci, Structures of Norbornane and Dodecahedrane by Molecular Mechanics Calculations (MM3), X-ray Crystallography, and Electron Diffraction, *J. Am. Chem. Soc.*, *111*, 1106-1114 (1989) [stretch-torsion cross term]

N. L. Allinger, F. Li and L. Yan, Molecular Mechanics. The MM3 Force Field for Alkenes, *J. Comput. Chem.*, *11*, 848-867 (1990)

N. L. Allinger, F. Li, L. Yan and J. C. Tai, Molecular Mechanics (MM3) Calculations on Conjugated Hydrocarbons, *J. Comput. Chem.*, *11*, 868-895 (1990)

J.-H. Lii and N. L. Allinger, Directional Hydrogen Bonding in the MM3 Force Field. I, *J. Phys. Org. Chem.*, *7*, 591-609 (1994)

J.-H. Lii and N. L. Allinger, Directional Hydrogen Bonding in the MM3 Force Field. II, *J. Comput. Chem.*, *19*, 1001-1016 (1998)

All parameters distributed with TINKER are from the "MM3 (2000) Parameter Set", as provided by N. L. Allinger, University of Georgia, August 2000

MM3PRO.PRM

Protein-only version of the MM3 parameters.

J.-H. Lii and N. L. Allinger, The MM3 Force Field for Amides, Polypeptides and Proteins, *J. Comput. Chem.*, *12*, 186-199 (1991)

MMFFPRO.PRM

Protein-only version of the MMFF94 parameters. (*in preparation*)

T. A. Halgren, Merck Molecular Force Field. I. Basis, Form, Scope, Parameterization, and Performance of MMFF94, *J. Comput. Chem.*, *17*, 490-519, 1996

OPLS.PRM

Complete OPLS-UA with united-atom parameters for proteins and many classes of organic molecules. Explicit hydrogens on polar atoms and aromatic carbons.

W. L. Jorgensen and J. Tirado-Rives, The OPLS Potential Functions for Proteins. Energy Minimizations for Crystals of Cyclic Peptides and Crambin, *J. Am. Chem. Soc.*, **110**, 1657-1666 (1988) [peptide and proteins]

W. L. Jorgensen and D. L. Severance, Aromatic-Aromatic Interactions: Free Energy Profiles for the Benzene Dimer in Water, Chloroform, and Liquid Benzene, *J. Am. Chem. Soc.*, **112**, 4768-4774 (1990) [aromatic hydrogens]

S. J. Weiner, P. A. Kollman, D. A. Case, U. C. Singh, C. Ghio, G. Alagona, S. Profeta, Jr. and P. Weiner, A New Force Field for Molecular Mechanical Simulation of Nucleic Acids and Proteins, *J. Am. Chem. Soc.*, **106**, 765-784 (1984) [united-atom "AMBER/OPLS" local geometry]

S. J. Weiner, P. A. Kollman, D. T. Nguyen and D. A. Case, An All Atom Force Field for Simulations of Proteins and Nucleic Acids, *J. Comput. Chem.*, **7**, 230-252 (1986) [all-atom "AMBER/OPLS" local geometry]

L. X. Dang and B. M. Pettitt, Simple Intramolecular Model Potentials for Water, *J. Phys. Chem.*, **91**, 3349-3354 (1987) [flexible TIP3P and SPC water]

W. L. Jorgensen, J. D. Madura and C. J. Swenson, Optimized Intermolecular Potential Functions for Liquid Hydrocarbons, *J. Am. Chem. Soc.*, **106**, 6638-6646 (1984) [hydrocarbons]

W. L. Jorgensen, E. R. Laird, T. B. Nguyen and J. Tirado-Rives, Monte Carlo Simulations of Pure Liquid Substituted Benzenes with OPLS Potential Functions, *J. Comput. Chem.*, **14**, 206-215 (1993) [substituted benzenes]

E. M. Duffy, P. J. Kowalczyk and W. L. Jorgensen, Do Denaturants Interact with Aromatic Hydrocarbons in Water?, *J. Am. Chem. Soc.*, **115**, 9271-9275 (1993) [benzene, naphthalene, urea, guanidinium, tetramethyl ammonium]

W. L. Jorgensen and C. J. Swenson, Optimized Intermolecular Potential Functions for Amides and Peptides. Structure and Properties of Liquid Amides, *J. Am. Chem. Soc.*, **106**, 765-784 (1984) [amides]

W. L. Jorgensen, J. M. Briggs and M. L. Contreras, Relative Partition Coefficients for Organic Solutes from Fluid Simulations, *J. Phys. Chem.*, **94**, 1683-1686 (1990) [chloroform, pyridine, pyrazine, pyrimidine]

J. M. Briggs, T. B. Nguyen and W. L. Jorgensen, Monte Carlo Simulations of Liquid Acetic Acid and Methyl Acetate with the OPLS Potential Functions, *J. Phys. Chem.*, **95**, 3315-3322 (1991) [acetic acid, methyl acetate]

H. Liu, F. Muller-Plathe and W. F. van Gunsteren, A Force Field for Liquid Dimethyl Sulfoxide and Physical Properties of Liquid Dimethyl Sulfoxide Calculated Using Molecular Dynamics Simulation, *J. Am. Chem. Soc.*, **117**, 4363-4366 (1995) [dimethyl sulfoxide]

J. Gao, X. Xia and T. F. George, Importance of Bimolecular Interactions in Developing Empirical Potential Functions for Liquid Ammonia, *J. Phys. Chem.*, **97**, 9241-9246 (1993) [ammonia]

J. Aqvist, Ion-Water Interaction Potentials Derived from Free Energy Perturbation Simulations, *J. Phys. Chem.*, **94**, 8021-8024 (1990) [metal ions]

W. S. Ross and C. C. Hardin, Ion-Induced Stabilization of the G-DNA Quadruplex: Free Energy Perturbation Studies, *J. Am. Chem. Soc.*, **116**, 4363-4366 (1994) [alkali metal ions]

J. Chandrasekhar, D. C. Spellmeyer and W. L. Jorgensen, Energy Component Analysis for Dilute Aqueous Solutions of Li⁺, Na⁺, F⁻, and Cl⁻ Ions, *J. Am. Chem. Soc.*, **106**, 903-910 (1984) [halide ions]

Most parameters distributed with TINKER are from "OPLS and OPLS-AA Parameters for Organic Molecules, Ions, and Nucleic Acids" as provided by W. L. Jorgensen, Yale University, October 1997

OPLSAA.PRM

OPLS-AA with all-atom parameters for proteins and many general classes of organic molecules.

W. L. Jorgensen, D. S. Maxwell and J. Tirado-Rives, Development and Testing of the OPLS All-Atom Force Field on Conformational Energetics and Properties of Organic Liquids, *J. Am. Chem. Soc.*, **117**, 11225-11236 (1996)

W. L. Jorgensen and N. A. McDonald, Development of an All-Atom Force Field for Heterocycles. Properties of Liquid Pyridine and Diazenes, *THEOCHEM-J. Mol. Struct.*, **424**, 145-155 (1998)

N. A. McDonald and W. L. Jorgensen, Development of an All-Atom Force Field for Heterocycles. Properties of Liquid Pyrrole, Furan, Diazoles, and Oxazoles, *J. Phys. Chem. B*, **102**, 8049-8059 (1998)

All parameters distributed with TINKER are from "OPLS and OPLS-AA Parameters for Organic Molecules, Ions, and Nucleic Acids" as provided by W. L. Jorgensen, Yale University, October 1997

SMOOTH.PRM

Version of OPLS-UA for use with potential smoothing. Largely adapted largely from standard OPLS-UA parameters with modifications to the vdw and improper torsion terms.

R. V. Pappu, R. K. Hart and J. W. Ponder, Analysis and Application of Potential Energy Smoothing and Search Methods for Global Optimization, *J. Phys. Chem. B*, **102**, 9725-9742 (1998) [smoothing modifications]

SMOOTHAA.PRM

Version of OPLS-AA for use with potential smoothing. Largely adapted largely from standard OPLS-AA parameters with modifications to the vdw and improper torsion terms.

R. V. Pappu, R. K. Hart and J. W. Ponder, Analysis and Application of Potential Energy Smoothing and Search Methods for Global Optimization, *J. Phys. Chem. B*, 102, 9725-9742 (1998) [smoothing modifications]

TINKER.PRM

Preliminary parameters for the TINKER polarizable multipole force field. As the release of TINKER 3.9 we have completed parametrization for over 50 small molecule systems, aimed toward the first version of a TINKER force field for proteins and peptides. For further information, or if you are interested in testing a beta parameter set, please contact the TINKER developers.

WATER.PRM

The current TINKER water parameters for a polarizable multipole electrostatics model. This model is equal or better to the best available water models for many bulk and cluster properties.

Y. Kong and J. W. Ponder, Calculation of the Reaction Field Due to Off-Center Point Multipoles, *J. Chem. Phys.*, 107, 481-492 (1997)

The parameters distributed with TINKER are modified values based on the work of Pengyu Ren starting from the Ph.D. thesis of Yong Kong, "Multipole Electrostatic Methods for Protein Modeling with Reaction Field Treatment", Biochemistry & Molecular Biophysics, Washington University, St. Louis, August, 1997

7. Use of the Keyword Control File

This section contains a description of the keyword parameters which may be used to define or alter the course of a TINKER calculation. The keyword control file is optional in the sense that all of the TINKER programs will run in the absence of a keyfile and will simply use default values or query the user for needed information. However, the keywords allow use of a wide variety of algorithmic and procedural options, many of which are unavailable interactively.

Keywords are read from the keyword control file. All programs look first for a keyfile with the same base name as the input molecular system and ending in the extension **.key**. If this file does not exist, then TINKER tries to use a generic keyfile with the name **tinker.key** and located in the same directory as the input system. If neither a system specific nor a generic keyfile is present, TINKER will continue by using default values for keyword options and asking interactive questions as necessary.

TINKER searches the keyfile during the course of a calculation for relevant keywords that may be present. All keywords must appear as the first word on the line. Any blank space to the left of the keyword is ignored, and all contents of the keyfiles are case insensitive. Some keywords take modifiers; *i.e.*, TINKER looks further on the same line for additional information, such as the value of some parameter related to the keyword. Modifier information is read in free format, but must be completely contained on the same line as the original keyword. Any lines contained in the keyfile which do not qualify as valid keyword lines are treated as comments and are simply ignored.

Several keywords take a list of integer values (atom numbers, for example) as modifiers. For these keywords the integers can simply be listed explicitly and separated by spaces, commas or tabs. If a range of numbers is desired, it can be specified by listing the negative of the first number of the range, followed by a separator and the last number of the range. For example, the keyword line **ACTIVE 4 -9 17 23** could be used to add atoms 4, 9 through 17, and 23 to the set of active atoms during a TINKER calculation.

Listed below are the valid TINKER keywords sorted into groups by general function. The section ends with an alphabetical listing of the individual keywords along with brief descriptions of their action and possible modifiers, and examples of usage.

Keywords Grouped by Functionality

OUTPUT CONTROL KEYWORDS

| | | |
|--------------|---------------|------------|
| ARCHIVE | DEBUG | DIGITS |
| ECHO | EXIT-PAUSE | NOVERSION |
| OVERWRITE | PRINTOUT | SAVE-CYCLE |
| SAVE-INDUCED | SAVE-VELOCITY | VERBOSE |
| WRITEOUT | | |

FORCE FIELD SELECTION KEYWORDS

| | |
|------------|------------|
| FORCEFIELD | PARAMETERS |
|------------|------------|

POTENTIAL FUNCTION SELECTION KEYWORDS

| | | |
|--------------|--------------|-------------|
| ANGANGTERM | ANGLETERM | BONDTERM |
| CHARGETERM | CHGDPLTERM | DIPOLETERM |
| EXTRATERM | IMPROPTERM | IMPTORSTERM |
| METALTERM | MPOLETERM | OPBENDTERM |
| OPDISTTERM | POLARIZETERM | RESTRANTERM |
| RXNFIELDTERM | SOLVATETERM | STRBNDTERM |
| STRTORTERM | TORSIONTERM | TORTORTERM |
| UREYTERM | VDWTERM | |

POTENTIAL FUNCTION PARAMETER KEYWORDS

| | | |
|----------|-----------|----------|
| ANGANG | ANGLE | ANGLE3 |
| ANGLE4 | ANGLE5 | ANGLEF |
| ATOM | BIOTYPE | BOND |
| BOND3 | BOND4 | BOND5 |
| CHARGE | DIPOLE | DIPOLE3 |
| DIPOLE4 | DIPOLE5 | ELECTNEG |
| HBOND | IMPROPER | IMPTORS |
| METAL | MULTIPOLE | OPBEND |
| OPDIST | PIATOM | PIBOND |
| POLARIZE | SOLVATE | STRBND |
| STRTORS | TORSION | TORSION4 |
| TORSION5 | UREYBRAD | VDW |
| VDW14 | VDWPR | |

ENERGY UNIT CONVERSION KEYWORDS

| | | |
|-------------|------------|------------|
| ANGLEUNIT | ANGANGUNIT | BONDUNIT |
| IMPROPUNIT | IMPTORUNIT | OPBENDUNIT |
| OPDISTUNIT | STRBNDUNIT | STRTORUNIT |
| TORSIONUNIT | UREYUNIT | |

LOCAL GEOMETRY FUNCTIONAL FORM KEYWORDS

| | | |
|--------------|---------------|--------------|
| ANGLE-CUBIC | ANGLE-QUARTIC | ANGLE-PENTIC |
| ANGLE-SEXTIC | BOND-CUBIC | BOND-QUARTIC |
| BONDTYPE | MM2-STRBND | PISYSTEM |
| UREY-CUBIC | UREY-QUARTIC | |

VAN DER WAALS FUNCTIONAL FORM KEYWORDS

| | | |
|---------------|--------------|---------------|
| A-EXPTERM | B-EXPTERM | C-EXPTERM |
| DELTA-HALGREN | EPSILONRULE | GAMMA-HALGREN |
| GAUSSTYPE | RADIUSRULE | RADIUSSIZE |
| RADIUSTYPE | VDW-12-SCALE | VDW-13-SCALE |
| VDW-14-SCALE | VDW-15-SCALE | VDWTYPE |

ELECTROSTATICS FUNCTIONAL FORM KEYWORDS

| | | |
|--------------|--------------|--------------|
| CHG-12-SCALE | CHG-13-SCALE | CHG-14-SCALE |
|--------------|--------------|--------------|

CHG-15-SCALE
 DIRECT-12-SCALE
 MUTUAL-11-SCALE
 MUTUAL-14-SCALE
 POLAR-13-SCALE
 POLAR-EPS
 POLARIZATION

DIELECTRIC
 DIRECT-13-SCALE
 MUTUAL-12-SCALE
 POLAR-11-SCALE
 POLAR-14-SCALE
 POLAR-OLD
 REACTIONFIELD

DIRECT-11-SCALE
 DIRECT-14-SCALE
 MUTUAL-13-SCALE
 POLAR-12-SCALE
 POLAR-DAMP
 POLAR-SOR

NONBONDED CUTOFF KEYWORDS

CHG-CUTOFF
 DPL-CUTOFF
 LIGHTS
 NEIGHBOR-GROUPS
 TAPER
 VDW-TAPER

CHG-TAPER
 DPL-TAPER
 MPOLE-CUTOFF
 NEUTRAL-GROUPS
 TRUNCATE

CUTOFF
 HESS-CUTOFF
 MPOLE-TAPER
 POLYMER-CUTOFF
 VDW-CUTOFF

EWALD SUMMATION KEYWORDS

EWALD
 EWALD-CUTOFF
 PME-ORDER

EWALD-ALPHA
 EWALD-FRACTION

EWALD-BOUNDARY
 PME-GRID

CRYSTAL LATTICE & PERIODIC BOUNDARY KEYWORDS

A-AXIS
 ALPHA
 OCTAHEDRON

B-AXIS
 BETA
 SPACEGROUP

C-AXIS
 GAMMA

OPTIMIZATION KEYWORDS

ANGMAX
 HGUSS
 MAXITER
 STEPMAX

CAPPA
 INTMAX
 NEWHESS
 STPMIN

FCTMIN
 LBFGS-VECTORS
 NEXTITER

DYNAMICS KEYWORDS

COLLISION
 FRICTION-SCALING
 TAU-TEMPERATURE

COMPRESS
 INTEGRATE
 THERMOSTAT

FRICTION
 TAU-PRESSURE

TRANSITION STATE KEYWORDS

DIVERGE
 SADDLEPOINT

GAMMAMIN

REDUCE

DISTANCE GEOMETRY KEYWORDS

TRIAL-DISTANCE

TRIAL-DISTRIBUTION

RANDOM NUMBER KEYWORDS

RANDOMSEED

FREE ENERGY PERTURBATION KEYWORDS

LAMBDA

MUTATE

PARTIAL STRUCTURE KEYWORDS

ACTIVE

GROUP-INTRA

INACTIVE

GROUP

GROUP-MOLECULE

GROUP-INTER

GROUP-SELECT

CONSTRAINT & RESTRAINT KEYWORDS

BASIN

RESTRAIN-DIHEDRAL

SPHERE

RATTLE

RESTRAIN-DISTANCE

WALL

RATTLE-BOND

RESTRAIN-POSITION

POTENTIAL SMOOTHING KEYWORDS

DEFORM

DIFFUSE-VDW

DIFFUSE-CHARGE

DIFFUSE-TORSION

Description of Individual Keywords

The following is an alphabetical list of the TINKER keywords along with a brief description of the action of each keyword and required or optional parameters that can be used to extend or modify each keyword. The form of possible modifiers, if any, is shown in brackets following each keyword.

A-AXIS [real] Sets the value of the a-axis length for a crystal unit cell, or, equivalently, the X-axis length for a periodic box. The length value in Angstroms is listed after the keyword.

A-EXPTERM [real] Sets the value of the “A” premultiplier term in the Buckingham van der Waals function, *i.e.*, the value of A in the formula $E_{\text{vdw}} = \epsilon \{ A \exp[-B(R_0/R)] - C (R_0/R)^6 \}$.

ACTIVE [integer list] Sets the list of active atoms during a TINKER computation. Individual potential energy terms are computed when at least one atom involved in the term is active. For Cartesian space calculations, active atoms are those allowed to move. For torsional space calculations, rotations are allowed when all atoms on one side of the rotated bond are active. Multiple ACTIVE lines can be present in the keyfile and are treated cumulatively. On each line the keyword can be followed by one or more atom numbers or atom ranges. The presence of any ACTIVE keyword overrides any INACTIVE keywords in the keyfile.

ALPHA [real] Sets the value of the α angle of a crystal unit cell, *i.e.*, the angle between the b-axis and c-axis of a unit cell, or, equivalently, the angle between the Y-axis and Z-axis of a periodic box. The default value in the absence of the ALPHA keyword is 90 degrees.

ANGANG [1 integer & 3 reals] This keyword provides the values for a single angle-angle cross term potential parameter.

ANGANGTERM [NONE/ONLY] This keyword controls use of the angle-angle cross term potential energy. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

ANGANGUNIT [real] Sets the scale factor needed to convert the energy value computed by the angle-angle cross term potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default of $(\pi/180)^2 = 0.0003046$ is used, if the ANGANGUNIT keyword is not given in the force field parameter file or the keyfile.

ANGLE [3 integers & 4 reals] This keyword provides the values for a single bond angle bending parameter. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle which is to be defined. The real number modifiers give the force constant value for the angle and up to three ideal bond angles in degrees. In most cases only one ideal bond angle is given, and that value is used for all occurrences of the specified bond angle. If all three ideal angles are given, the values apply when the central atom of the angle is attached to 0, 1 or 2 additional hydrogen atoms, respectively. This “hydrogen environment” option is provided to implement the corresponding feature of Allinger’s MM force fields. The default units for the force constant are kcal/mole/radian², but this can be controlled via the ANGLEUNIT keyword.

ANGLE-CUBIC [real] Sets the value of the cubic term in the Taylor series expansion form of the bond angle bending potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the angle bending energy unit conversion factor, the force constant, and the cube of the deviation of the bond angle from its ideal value gives the cubic contribution to the angle bending energy. The default value in the absence of the ANGLE-CUBIC keyword is zero; *i.e.*, the cubic angle bending term is omitted.

ANGLE-PENTIC [real] Sets the value of the fifth power term in the Taylor series expansion form of the bond angle bending potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the angle bending energy unit conversion factor, the force constant, and the fifth power of the deviation of the bond angle from its ideal value gives the pentic contribution to the angle bending energy. The default value in the absence of the ANGLE-PENTIC keyword is zero; *i.e.*, the pentic angle bending term is omitted.

ANGLE-QUARTIC [real] Sets the value of the quartic term in the Taylor series expansion form of the bond angle bending potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the angle bending energy unit conversion factor, the force constant, and the fourth power of the deviation of the bond angle from its ideal value gives the quartic contribution to the angle bending energy. The default value in the absence of the ANGLE-QUARTIC keyword is zero; *i.e.*, the quartic angle bending term is omitted.

ANGLE-SEXTIC [real] Sets the value of the sixth power term in the Taylor series expansion form of the bond angle bending potential energy. The real number modifier gives

the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the angle bending energy unit conversion factor, the force constant, and the sixth power of the deviation of the bond angle from its ideal value gives the sextic contribution to the angle bending energy. The default value in the absence of the ANGLE-SEXTIC keyword is zero; *i.e.*, the sextic angle bending term is omitted.

ANGLE3 [3 integers & 4 reals] This keyword provides the values for a single bond angle bending parameter specific to atoms in 3-membered rings. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle which is to be defined. The real number modifiers give the force constant value for the angle and up to three ideal bond angles in degrees. If all three ideal angles are given, the values apply when the central atom of the angle is attached to 0, 1 or 2 additional hydrogen atoms, respectively. The default units for the force constant are kcal/mole/radian², but this can be controlled via the ANGLEUNIT keyword. If any ANGLE3 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special ANGLE3 parameters be given for all angles in 3-membered rings. In the absence of any ANGLE3 keywords, standard ANGLE parameters will be used for bonds in 3-membered rings.

ANGLE4 [3 integers & 4 reals] This keyword provides the values for a single bond angle bending parameter specific to atoms in 4-membered rings. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle which is to be defined. The real number modifiers give the force constant value for the angle and up to three ideal bond angles in degrees. If all three ideal angles are given, the values apply when the central atom of the angle is attached to 0, 1 or 2 additional hydrogen atoms, respectively. The default units for the force constant are kcal/mole/radian², but this can be controlled via the ANGLEUNIT keyword. If any ANGLE4 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special ANGLE4 parameters be given for all angles in 4-membered rings. In the absence of any ANGLE4 keywords, standard ANGLE parameters will be used for bonds in 4-membered rings.

ANGLE5 [3 integers & 4 reals] This keyword provides the values for a single bond angle bending parameter specific to atoms in 5-membered rings. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle which is to be defined. The real number modifiers give the force constant value for the angle and up to three ideal bond angles in degrees. If all three ideal angles are given, the values apply when the central atom of the angle is attached to 0, 1 or 2 additional hydrogen atoms, respectively. The default units for the force constant are kcal/mole/radian², but this can be controlled via the ANGLEUNIT keyword. If any ANGLE5 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special ANGLE5 parameters be given for all angles in 5-membered rings. In the absence of any ANGLE5 keywords, standard ANGLE parameters will be used for bonds in 5-membered rings.

ANGLEF [3 integers & 3 reals] This keyword provides the values for a single bond angle bending parameter for a SHAPES-style Fourier potential function. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle which is to be defined. The real number modifiers give the force constant value for the angle, the angle shift in degrees, and the periodicity value. Note that the force constant should be given as the “harmonic” value and not the native Fourier value. The default units for the force constant are kcal/mole/radian², but this can be controlled via the ANGLEUNIT keyword.

ANGLETERM [NONE/ONLY] This keyword controls use of the bond angle bending potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

ANGLEUNIT [real] Sets the scale factor needed to convert the energy value computed by the bond angle bending potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of $(\pi/180)^2 = 0.0003046$ is used, if the ANGLEUNIT keyword is not given in the force field parameter file or the keyfile.

ANGMAX [real] Set the maximum permissible angle between the current optimization search direction and the negative of the gradient direction. If this maximum angle value is exceeded, the optimization routine will note an error condition and may restart from the steepest descent direction. The default value in the absence of the ANGMAX keyword is usually 88 degrees for conjugate gradient methods and 180 degrees (*i.e.*, disabled) for variable metric optimizations.

ARCHIVE Causes TINKER molecular dynamics-based programs to write trajectories directly to a single plain-text archive file with the `.arc` format. If an archive file already exists at the start of the calculation, then the newly generated trajectory is appended to the end of the existing file. The default in the absence of this keyword is to write the trajectory snapshots to consecutively numbered cycle files.

ATOM [2 integers, name, quoted string, integer, real & integer] This keyword provides the values needed to define a single force field atom type.

B-AXIS [real] Sets the value of the b-axis length for a crystal unit cell, or, equivalently, the Y-axis length for a periodic box. The length value in Angstroms is listed after the keyword. If the keyword is absent, the b-axis length is set equal to the a-axis length.

B-EXPTERM [real] Sets the value of the “B” exponential factor in the Buckingham van der Waals function, *i.e.*, the value of B in the formula $E_{\text{vdw}} = \epsilon \{ A \exp[-B(R_0/R)] - C (R_0/R)^6 \}$.

BASIN [2 reals] Presence of this keyword turns on a “basin” restraint potential function that serves to drive the system toward a compact structure. The actual function is a Gaussian of the form $E_{\text{basin}} = \sum A \exp[-B R^2]$, summed over all pairs of atoms where R is the distance between atoms. The A and B values are the depth and width parameters given as modifiers to the BASIN keyword. This potential is currently used to control the degree of expansion during potential energy smooth procedures through the use of shallow, broad basins.

BETA [real] Sets the value of the β angle of a crystal unit cell, *i.e.*, the angle between the a-axis and c-axis of a unit cell, or, equivalently, the angle between the X-axis and Z-axis of a periodic box. The default value in the absence of the BETA keyword is to set the β angle equal to the α angle as given by the keyword ALPHA.

BIOTYPE [integer, name, quoted string & integer] This keyword provides the values to define the correspondence between a single biopolymer atom type and its force field atom type.

BOND [2 integers & 2 reals] This keyword provides the values for a single bond stretching parameter. The integer modifiers give the atom class numbers for the two kinds of atoms involved in the bond which is to be defined. The real number modifiers give the force constant value for the bond and the ideal bond length in Å. The default units for the force constant are kcal/mole/Å², but this can be controlled via the BONDUNIT keyword.

BOND-CUBIC [real] Sets the value of the cubic term in the Taylor series expansion form of the bond stretching potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the bond stretching energy unit conversion factor, the force constant, and the cube of the deviation of the bond length from its ideal value gives the cubic contribution to the bond stretching energy. The default value in the absence of the BOND-CUBIC keyword is zero; *i.e.*, the cubic bond stretching term is omitted.

BOND-QUARTIC [real] Sets the value of the quartic term in the Taylor series expansion form of the bond stretching potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. This term multiplied by the bond stretching energy unit conversion factor, the force constant, and the fourth power of the deviation of the bond length from its ideal value gives the quartic contribution to the bond stretching energy. The default value in the absence of the BOND-QUARTIC keyword is zero; *i.e.*, the quartic bond stretching term is omitted.

BOND3 [2 integers & 2 reals] This keyword provides the values for a single bond stretching parameter specific to atoms in 3-membered rings. The integer modifiers give the atom class numbers for the two kinds of atoms involved in the bond which is to be defined. The real number modifiers give the force constant value for the bond and the ideal bond length in Å. The default units for the force constant are kcal/mole/Å², but this can be controlled via the BONDUNIT keyword. If any BOND3 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special BOND3 parameters be given for all bonds in 3-membered rings. In the absence of any BOND3 keywords, standard BOND parameters will be used for bonds in 3-membered rings.

BOND4 [2 integers & 2 reals] This keyword provides the values for a single bond stretching parameter specific to atoms in 4-membered rings. The integer modifiers give the atom class numbers for the two kinds of atoms involved in the bond which is to be defined. The real number modifiers give the force constant value for the bond and the ideal bond length in Å. The default units for the force constant are kcal/mole/Å², but this can be controlled via the BONDUNIT keyword. If any BOND4 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special BOND4 parameters be given for all bonds in 4-membered rings. In the absence of any BOND4 keywords, standard BOND parameters will be used for bonds in 4-membered rings.

BOND5 [2 integers & 2 reals] This keyword provides the values for a single bond stretching parameter specific to atoms in 5-membered rings. The integer modifiers give the atom class numbers for the two kinds of atoms involved in the bond which is to be defined. The real number modifiers give the force constant value for the bond and the ideal bond length in Å. The default units for the force constant are kcal/mole/Å², but this can be controlled via the BONDUNIT keyword. If any BOND5 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special BOND5 parameters be given for all bonds in 5-membered rings. In the absence of any BOND5 keywords, standard BOND parameters will be used for bonds in 5-membered rings.

BONDTerm [NONE/ONLY] This keyword controls use of the bond stretching potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

BONDType [TAYLOR/MORSE/GAUSSIAN] Chooses the functional form of the bond stretching potential. The TAYLOR option selects a Taylor series expansion containing terms from harmonic through quartic. The MORSE option selects a Morse potential fit to the ideal bond length and stretching force constant parameter values. The GAUSSIAN option uses an inverted Gaussian with amplitude equal to the Morse bond dissociation energy and width set to reproduce the vibrational frequency of a harmonic potential. The default is to use the TAYLOR potential.

BONDUnit [real] Sets the scale factor needed to convert the energy value computed by the bond stretching potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the BONDUNIT keyword is not given in the force field parameter file or the keyfile.

C-AXIS [real] Sets the value of the C-axis length for a crystal unit cell, or, equivalently, the Z-axis length for a periodic box. The length value in Angstroms is listed after the keyword. If the keyword is absent, the C-axis length is set equal to the A-axis length.

C-EXPTerm [real] Sets the value of the “C” dispersion multiplier in the Buckingham van der Waals function, *i.e.*, the value of C in the formula $E_{\text{vdw}} = \epsilon \{ A \exp[-B(R_0/R)] - C (R_0/R)^6 \}$.

CAPPA [real] This keyword is used to set the normal termination criterion for the line search phase of TINKER optimization routines. The line search exits successfully if the ratio of the current gradient projection on the line to the projection at the start of the line search falls below the value of CAPPA. A default value of 0.1 is used in the absence of the CAPPA keyword.

CHARGE [1 integer & 1 real] This keyword provides a value for a single atomic partial charge electrostatic parameter. The integer modifier, if positive, gives the atom type number for which the charge parameter is to be defined. Note that vdw parameters are given for atom types, not atom classes. If the integer modifier is negative, then the parameter value to follow applies only to the individual atom whose atom number is the negative of the modifier. The real number modifier gives the values of the atomic partial charge in electrons.

CHARGEterm [NONE/ONLY] This keyword controls use of the charge-charge potential energy term between pairs of atomic partial charges. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

CHG-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to charge-charge electrostatic interactions between 1-2 connected atoms, *i.e.*, atoms that are directly bonded. The default value of 0.0 is used, if the CHG-12-SCALE keyword is not given in either the parameter file or the keyfile.

CHG-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to charge-charge electrostatic interactions between 1-3 connected atoms, *i.e.*, atoms separated by two covalent bonds. The default value of 0.0 is used, if the CHG-13-SCALE keyword is not given in either the parameter file or the keyfile.

CHG-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to charge-charge electrostatic interactions between 1-4 connected atoms, *i.e.*, atoms separated by three covalent bonds. The default value of 1.0 is used, if the CHG-14-SCALE keyword is not given in either the parameter file or the keyfile.

CHG-15-SCALE [real] This keyword provides a multiplicative scale factor that is applied to charge-charge electrostatic interactions between 1-5 connected atoms, *i.e.*, atoms separated by four covalent bonds. The default value of 1.0 is used, if the CHG-15-SCALE keyword is not given in either the parameter file or the keyfile.

CHG-CUTOFF [real] Sets the cutoff distance value in Angstroms for charge-charge electrostatic potential energy interactions. The energy for any pair of sites beyond the cutoff distance will be set to zero. Other keywords can be used to select a smoothing scheme near the cutoff distance. The default cutoff distance in the absence of the CHG-CUTOFF keyword is infinite for nonperiodic systems and 9.0 for periodic systems.

CHG-TAPER [real] This keyword allows modification of the cutoff window for charge-charge electrostatic potential energy interactions. It is similar in form and action to the TAPER keyword, except that its value applies only to the charge-charge potential. The default value in the absence of the CHG-TAPER keyword is to begin the cutoff window at 0.65 of the corresponding cutoff distance.

CHGDPLTERM [NONE/ONLY] This keyword controls use of the charge-dipole potential energy term between atomic partial charges and bond dipoles. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

COLLISION [real] Sets the value of the random collision frequency used in the Andersen stochastic collision dynamics thermostat. The supplied value has units of $\text{fs}^{-1} \text{atom}^{-1}$ and is multiplied internal to TINKER by the time step in fs and $N^{-2/3}$ where N is the number of atoms. The default value used in the absence of the COLLISION keyword is 0.1 which is appropriate for many systems but may need adjustment to achieve adequate temperature control without perturbing the dynamics.

COMPRESS [real] Sets the value of the bulk solvent isothermal compressibility in Atm^{-1} for use during pressure computation and scaling in molecular dynamics computations. The default value used in the absence of the COMPRESS keyword is 0.000046, appropriate for water. This parameter serves as a scale factor for the Groningen-style pressure bath coupling time, and its exact value should not be of critical importance.

CUTOFF [real] Sets the cutoff distance value for all nonbonded potential energy interactions. The energy for any of the nonbonded potentials of a pair of sites beyond the cutoff distance will be set to zero. Other keywords can be used to select a smoothing scheme near the cutoff distance, or to apply different cutoff distances to various nonbonded energy terms.

DEBUG Turns on printing of detailed information and intermediate values throughout the progress of a TINKER computation; not recommended for use with large structures or full potential energy functions since a summary of every individual interaction will usually be output.

DEFORM [real] Sets the amount of diffusion equation-style smoothing that will be applied to the potential energy surface when using the SMOOTH force field. The real number option is equivalent to the “time” value in the original Piela, *et al.* formalism; the larger the value, the greater the smoothing. The default value is zero, meaning that no smoothing will be applied.

DELTA-HALGREN [real] Sets the value of the δ parameter in Halgren’s buffered 14-7 vdw potential energy functional form. In the absence of the DELTA-HALGREN keyword, a default value of 0.07 is used.

DIELECTRIC [real] Sets the value of the bulk dielectric constant used to damp all electrostatic interaction energies for any of the TINKER electrostatic potential functions. The default value is force field dependent, but is usually equal to 1.0 (for Allinger’s MM force fields the default is 1.5).

DIFFUSE-CHARGE [real] This keyword is used during potential function smoothing procedures to specify the effective diffusion coefficient to be applied to the smoothed form of the Coulomb’s Law charge-charge potential function. In the absence of the DIFFUSE-CHARGE keyword, a default value of 3.5 is used.

DIFFUSE-TORSION [real] This keyword is used during potential function smoothing procedures to specify the effective diffusion coefficient to be applied to the smoothed form of the torsion angle potential function. In the absence of the DIFFUSE-TORSION keyword, a default value of 0.0225 is used.

DIFFUSE-VDW [real] This keyword is used during potential function smoothing procedures to specify the effective diffusion coefficient to be applied to the smoothed Gaussian approximation to the Lennard-Jones van der Waals potential function. In the absence of the DIFFUSE-VDW keyword, a default value of 1.0 is used.

DIGITS [integer] This keyword controls the number of digits of precision output by TINKER in reporting potential energies and atomic coordinates. The allowed values for the integer modifier are 4, 6 and 8. Input values less than 4 will be set to 4, and those greater than 8 will be set to 8. Final energy values reported by most TINKER programs will contain the specified number of digits to the right of the decimal point. The number of decimal places to be output for atomic coordinates is generally two larger than the value of DIGITS. In the absence of the DIGITS keyword a default value of 4 is used, and energies will be reported to 4 decimal places with coordinates to 6 decimal places.

DIPOLE [2 integers & 2 reals] This keyword provides the values for a single bond dipole electrostatic parameter. The integer modifiers give the atom type numbers for the two kinds of atoms involved in the bond dipole which is to be defined. The real number modifiers give the value of the bond dipole in Debyes and the position of the dipole site along the bond. If the bond dipole value is positive, then the first of the two atom types is the positive end of the dipole. For a negative bond dipole value, the first atom type listed is negative. The position along the bond is an optional modifier that gives the position of the dipole site as a fraction between the first atom type (position=0) and the second atom type (position=1). The

default for the dipole position in the absence of a specified value is 0.5, placing the dipole at the midpoint of the bond.

DIPOLE3 [2 integers & 2 reals] This keyword provides the values for a single bond dipole electrostatic parameter specific to atoms in 3-membered rings. The integer modifiers give the atom type numbers for the two kinds of atoms involved in the bond dipole which is to be defined. The real number modifiers give the value of the bond dipole in Debyes and the position of the dipole site along the bond. The default for the dipole position in the absence of a specified value is 0.5, placing the dipole at the midpoint of the bond. If any DIPOLE3 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special DIPOLE3 parameters be given for all bond dipoles in 3-membered rings. In the absence of any DIPOLE3 keywords, standard DIPOLE parameters will be used for bonds in 3-membered rings.

DIPOLE4 [2 integers & 2 reals] This keyword provides the values for a single bond dipole electrostatic parameter specific to atoms in 4-membered rings. The integer modifiers give the atom type numbers for the two kinds of atoms involved in the bond dipole which is to be defined. The real number modifiers give the value of the bond dipole in Debyes and the position of the dipole site along the bond. The default for the dipole position in the absence of a specified value is 0.5, placing the dipole at the midpoint of the bond. If any DIPOLE4 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special DIPOLE4 parameters be given for all bond dipoles in 4-membered rings. In the absence of any DIPOLE4 keywords, standard DIPOLE parameters will be used for bonds in 4-membered rings.

DIPOLE5 [2 integers & 2 reals] This keyword provides the values for a single bond dipole electrostatic parameter specific to atoms in 5-membered rings. The integer modifiers give the atom type numbers for the two kinds of atoms involved in the bond dipole which is to be defined. The real number modifiers give the value of the bond dipole in Debyes and the position of the dipole site along the bond. The default for the dipole position in the absence of a specified value is 0.5, placing the dipole at the midpoint of the bond. If any DIPOLE5 keywords are present, either in the master force field parameter file or the keyfile, then TINKER requires that special DIPOLE5 parameters be given for all bond dipoles in 5-membered rings. In the absence of any DIPOLE5 keywords, standard DIPOLE parameters will be used for bonds in 5-membered rings.

DIPOLETERM [NONE/ONLY] This keyword controls use of the dipole-dipole potential energy term between pairs of bond dipoles. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

DIRECT-11-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the permanent (direct) field due to atoms within a polarization group during an induced dipole calculation, *i.e.*, atoms that are in the same polarization group as the atom being polarized. The default value of 0.0 is used, if the DIRECT-11-SCALE keyword is not given in either the parameter file or the keyfile.

DIRECT-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the permanent (direct) field due to atoms in 1-2 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups directly connected to the group containing the atom being polarized. The default value of 0.0 is used, if the DIRECT-12-SCALE keyword is not given in either the parameter file or the keyfile.

DIRECT-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the permanent (direct) field due to atoms in 1-3 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups separated by one group from the group containing the atom being polarized. The default value of 0.0 is used, if the DIRECT-13-SCALE keyword is not given in either the parameter file or the keyfile.

DIRECT-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the permanent (direct) field due to atoms in 1-4 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups separated by two groups from the group containing the atom being polarized. The default value of 1.0 is used, if the DIRECT-14-SCALE keyword is not given in either the parameter file or the keyfile.

DIVERGE [real] This keyword is used by the SADDLE program to set the maximum allowed value of the ratio of the gradient length along the path to the total gradient norm at the end of a cycle of minimization perpendicular to the path. If the value provided by the DIVERGE keyword is exceeded, then another cycle of maximization along the path is required. A default value of 0.005 is used in the absence of the DIVERGE keyword.

DPL-CUTOFF [real] Sets the cutoff distance value in Angstroms for bond dipole-bond dipole electrostatic potential energy interactions. The energy for any pair of bond dipole sites beyond the cutoff distance will be set to zero. Other keywords can be used to select a smoothing scheme near the cutoff distance. The default cutoff distance in the absence of the DPL-CUTOFF keyword is essentially infinite for nonperiodic systems and 10.0 for periodic systems.

DPL-TAPER [real] This keyword allows modification of the cutoff windows for bond dipole-bond dipole electrostatic potential energy interactions. It is similar in form and action to the TAPER keyword, except that its value applies only to the vdw potential. The default value in the absence of the DPL-TAPER keyword is to begin the cutoff window at 0.75 of the dipole cutoff distance.

ECHO [text string] The presence of this keyword causes whatever text follows it on the line to be copied directly to the output file. This keyword is also active in parameter files. It has no default value; if no text follows the ECHO keyword, a blank line is placed in the output file.

ELECTNEG [3 integers & 1 real] This keyword provides the values for a single electronegativity bond length correction parameter. The first two integer modifiers give the atom class numbers of the atoms involved in the bond to be corrected. The third integer modifier is the atom class of an electronegative atom. In the case of a primary correction, an atom of this third class must be directly bonded to an atom of the second atom class. For a secondary correction, the third class is one atom removed from an atom of the second class. The real number modifier is the value in Å by which the original ideal bond length is to be corrected.

EPSILONRULE [GEOMETRIC/ARITHMETIC/HARMONIC/HHG] This keyword selects the combining rule used to derive the ϵ value for van der Waals interactions. The default in the absence of the EPSILONRULE keyword is to use the GEOMETRIC mean of the individual ϵ values of the two atoms involved in the van der Waals interaction.

EWALD This keyword turns on the use of Ewald summation during computation of electrostatic interactions in periodic systems. In the current version of TINKER, regular Ewald is used for polarizable atomic multipoles, and smooth particle mesh Ewald (PME) is used for charge-charge interactions. Ewald summation is not available for interactions involving bond-centered dipoles. By default, in the absence of the EWALD keyword, distance-based cutoffs are used for electrostatic interactions.

EWALD-ALPHA [real] Sets the value of the Ewald coefficient which controls the width of the Gaussian screening charges during particle mesh Ewald summation. In the absence of the EWALD-ALPHA keyword, a value is chosen which causes interactions outside the real-space cutoff to be below a fixed tolerance. For most standard applications of Ewald summation, the program default should be used.

EWALD-BOUNDARY This keyword invokes the use of “vacuum” boundary conditions during Ewald summation, corresponding to the media surrounding the system having a dielectric value of 1. The default in the absence of the EWALD-BOUNDARY keyword is to use “tinfoil” boundary conditions where the surrounding media is assumed to have an infinite dielectric value.

EWALD-CUTOFF [real] Sets the value in Angstroms of the real-space distance cutoff for use during Ewald summation. By default, in the absence of the EWALD-CUTOFF keyword, a value of 9.0 is used.

EWALD-FRACTION [real] Sets the fraction between 0 and 1 of reciprocal space included in the reciprocal sum when using regular Ewald summation. The keyword has no effect on PME calculations. A default value of 0.5 is used in the absence of the EWALD-FRACTION keyword.

EXIT-PAUSE This keyword causes TINKER programs to pause and wait for a carriage return at the end of execution prior to returning control to the operating system. This is useful to keep the execution window open following termination on machines running Microsoft Windows or Apple MacOS. The default in the absence of the EXIT-PAUSE keyword, is to return control to the operating system immediately at program termination.

EXTRATERM [NONE/ONLY] This keyword controls use of the user defined extra potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

FCTMIN [real] This keyword sets a convergence criterion for successful completion of a TINKER optimization. If the value of the optimization objective function, typically the potential energy, falls below the value set by FCTMIN, then the optimization is deemed to have converged. The default value in the absence of the FCTMIN keyword is -1000000, effectively removing this criterion as a possible agent for termination.

FORCEFIELD [name] This keyword provides a name for the force field to be used in the current calculation. Its value is usually set in the master force field parameter file for the calculation (see the PARAMETERS keyword) instead of in the keyfile.

FRICTION [real] Sets the value of the frictional coefficient in ps^{-1} for use with stochastic dynamics. The default value used in the absence of the FRICTION keyword is 91.0, which is generally appropriate for water.

GAMMA [real] Sets the value of the γ angle of a crystal unit cell, *i.e.*, the angle between the a-axis and b-axis of a unit cell, or, equivalently, the angle between the X-axis and Z-axis of a periodic box. The default value in the absence of the GAMMA keyword is to set the γ angle equal to the α angle as given by the keyword ALPHA.

GAMMA-HALGREN [real] Sets the value of the γ parameter in Halgren's buffered 14-7 vdw potential energy functional form. In the absence of the DELTA-HALGREN keyword, a default value of 0.12 is used.

GAMMAMIN [real] Sets the convergence target value for γ during searches for maxima along the quadratic synchronous transit used by the SADDLE program. The value of γ is the square of the ratio of the gradient projection along the path to the total gradient. A default value of 0.00001 is used in the absence of the GAMMAMIN keyword.

GAUSSTYPE [LJ-2/LJ-4/MM2-2/MM3-2/IN-PLACE] This keyword specifies the underlying vdw form that a Gaussian vdw approximation will attempt to fit. number of terms to be used in a Gaussian approximation of the Lennard-Jones van der Waals potential. The text modifier gives the name of the functional form to be used. Thus LJ-2 as a modifier will result in a 2-Gaussian fit to a Lennard-Jones vdw potential. The GAUSSTYPE keyword only takes effect when VDWTYPE is set to GAUSSIAN. This keyword has no default value.

GROUP [integer, integer list] This keyword defines an atom group as a substructure within the full input molecular structure. The value of the first integer is the group number which must be in the range from 1 to the maximum number of allowed groups. The remaining integers give the atom or atoms contained in this group as one or more atom numbers or ranges. Multiple keyword lines can be used to specify additional atoms in the same group. Note that an atom can only be in one group, the last group to which it is assigned is the one used.

GROUP-INTER This keyword assigns a value of 1.0 to all inter-group interactions and a value of 0.0 to all intra-group interactions. For example, combination with the GROUP-MOLECULE keyword provides for rigid-body calculations.

GROUP-INTRA This keyword assigns a value of 1.0 to all intra-group interactions and a value of 0.0 to all inter-group interactions.

GROUP-MOLECULE This keyword sets each individual molecule in the system to be a separate atom group, but does not assign weights to group-group interactions.

GROUP-SELECT [2 integers, real] This keyword gives the weight in the final potential energy of a specified set of intra- or intergroup interactions. The integer modifiers give the group numbers of the groups involved. If the two numbers are the same, then an intragroup set of interactions is specified. The real modifier gives the weight by which all energetic interactions in this set will be multiplied before incorporation into the final potential energy. If omitted as a keyword modifier, the weight will be set to 1.0 by default. If any SELECT-GROUP keywords are present, then any set of interactions not specified in a SELECT-GROUP keyword is given a zero weight. The default when no SELECT-GROUP keywords are specified is to use all intergroup interactions with a weight of 1.0 and to set all intragroup interactions to zero.

HBOND [2 integers & 2 reals] This keyword provides the values for the MM3-style directional hydrogen bonding parameters for a single pair of atoms. The integer modifiers give the pair of atom class numbers for which hydrogen bonding parameters are to be defined. The two real number modifiers give the values of the minimum energy contact distance in Å and the well depth at the minimum distance in kcal/mole.

HESS-CUTOFF [real] This keyword defines a lower limit for significant Hessian matrix elements. During computation of the Hessian matrix of partial second derivatives, any matrix elements with absolute value below HESS-CUTOFF will be set to zero and omitted from the sparse matrix Hessian storage scheme used by TINKER. For most calculations, the default in the absence of this keyword is zero, i.e., all elements will be stored. For most Truncated Newton optimizations the Hessian cutoff will be chosen dynamically by the optimizer.

HGUESS [real] Sets an initial guess for the average value of the diagonal elements of the scaled inverse Hessian matrix used by the optimally conditioned variable metric optimization routine. A default value of 0.4 is used in the absence of the HGUESS keyword.

IMPROPER [4 integers & 2 reals] This keyword provides the values for a single CHARMM-style improper dihedral angle parameter.

IMPROPTERM [NONE/ONLY] This keyword controls use of the CHARMM-style improper dihedral angle potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

IMPROPUNIT [real] Sets the scale factor needed to convert the energy value computed by the CHARMM-style improper dihedral angle potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the IMPROPUNIT keyword is not given in the force field parameter file or the keyfile.

IMPTORS [4 integers & up to 3 real/real/integer triples] This keyword provides the values for a single AMBER-style improper torsional angle parameter. The first four integer modifiers give the atom class numbers for the atoms involved in the improper torsional angle to be defined. By convention, the third atom class of the four is the trigonal atom on which the improper torsion is centered. The torsional angle computed is literally that defined by the four atom classes in the order specified by the keyword. Each of the remaining triples of real/real/integer modifiers give the half-amplitude, phase offset in degrees and periodicity of a particular improper torsional term, respectively. Periodicities through 3-fold are allowed for improper torsional parameters.

IMPTORSTERM [NONE/ONLY] This keyword controls use of the AMBER-style improper torsional angle potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

IMPTORSUNIT [real] Sets the scale factor needed to convert the energy value computed by the AMBER-style improper torsional angle potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the IMPTORSUNIT keyword is not given in the force field parameter file or the keyfile.

INACTIVE [integer list] Sets the list of inactive atoms during a TINKER computation. Individual potential energy terms are not computed when all atoms involved in the term are inactive. For Cartesian space calculations, inactive atoms are not allowed to move. For torsional space calculations, rotations are not allowed when there are inactive atoms on both sides of the rotated bond. Multiple INACTIVE lines can be present in the keyfile, and on each line the keyword can be followed by one or more atom numbers or ranges. If any INACTIVE keys are found, all atoms are set to active except those listed on the INACTIVE lines. The ACTIVE keyword overrides all INACTIVE keywords found in the keyfile.

INTEGRATE [VERLET/BEEMAN/STOCHASTIC/RIGIDBODY] Chooses the integration method for propagation of dynamics trajectories. The keyword is followed on the same line by the name of the option. Standard Newtonian MD can be run using either VERLET for the Velocity Verlet method, or BEEMAN for the velocity form of Bernie Brook's "Better Beeman" method. A Velocity Verlet-based stochastic dynamics trajectory is selected by the STOCHASTIC modifier. A rigid-body dynamics method is selected by the RIGIDBODY modifier. The default integration scheme is MD using the BEEMAN method. *Note the RIGIDBODY option is still under development and should be treated as experimental code in the current version of TINKER.*

INTMAX [integer] Sets the maximum number of interpolation cycles that will be allowed during the line search phase of an optimization. All gradient-based TINKER optimization routines use a common line search routine involving quadratic extrapolation and cubic interpolation. If the value of INTMAX is reached, an error status is set for the line search and the search is repeated with a much smaller initial step size. The default value in the absence of this keyword is optimization routine dependent, but is usually in the range 5 to 10.

LAMBDA [real] This keyword sets the value of the λ path parameter for free energy perturbation calculations. The real number modifier specifies the position along the mutation path and must be a number in the range from 0 (initial state) to 1 (final state). The actual atoms involved in the mutation are given separately in individual MUTATE keyword lines.

LBFGS-VECTORS [integer] Sets the number of correction vectors used by the limited-memory L-BFGS optimization routine. The current maximum allowable value, and the default in the absence of the LBFGS-VECTORS keyword is 15.

LIGHTS This keyword turns on Method of Lights neighbor generation for the charge-charge potential and any of the van der Waals potentials. This method will yield identical energetic results to the standard double loop method. Method of Lights will be faster when the volume of a sphere with radius equal to the nonbond cutoff distance is significantly less than half the volume of the total system (*i.e.*, the full molecular system, the crystal unit cell or the periodic box).

MAXITER [integer] Sets the maximum number of minimization iterations that will be allowed for any TINKER program that uses any of the nonlinear optimization routines. The default value in the absence of this keyword is program dependent, but is always set to a very large number.

METAL This keyword provides the values for a single transition metal ligand field parameter. *Note this keyword is present in the code, but not active in the current version of TINKER.*

METALTERM [NONE/ONLY] This keyword controls use of the transition metal ligand field potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one. ***Note this keyword is present in the code, but not active in the current version of TINKER.***

MM2-STRBND This keyword switches the behavior of the stretch-bend potential function to match the formulation used by the MM2 force field. In MM2, stretching of bonds to attached hydrogen atoms is not including in computing the stretch-bend cross term energy. The default behavior in the absence of this keyword is to include stretching of attached hydrogen atoms as in the MM3 force field.

MPOLE-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to permanent atomic multipole electrostatic interactions between 1-2 connected atoms, *i.e.*, atoms that are directly bonded. The default value of 0.0 is used, if the MPOLE-12-SCALE keyword is not given in either the parameter file or the keyfile.

MPOLE-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to permanent atomic multipole electrostatic interactions between 1-3 connected atoms, *i.e.*, atoms separated by two covalent bonds. The default value of 0.0 is used, if the MPOLE-13-SCALE keyword is not given in either the parameter file or the keyfile.

MPOLE-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to permanent atomic multipole electrostatic interactions between 1-4 connected atoms, *i.e.*, atoms separated by three covalent bonds. The default value of 1.0 is used, if the MPOLE-14-SCALE keyword is not given in either the parameter file or the keyfile.

MPOLE-15-SCALE [real] This keyword provides a multiplicative scale factor that is applied to permanent atomic multipole electrostatic interactions between 1-5 connected atoms, *i.e.*, atoms separated by four covalent bonds. The default value of 1.0 is used, if the MPOLE-15-SCALE keyword is not given in either the parameter file or the keyfile.

MPOLE-CUTOFF [real] Sets the cutoff distance value in Angstroms for atomic multipole potential energy interactions. The energy for any pair of sites beyond the cutoff distance will be set to zero. Other keywords can be used to select a smoothing scheme near the cutoff distance. The default cutoff distance in the absence of the MPOLE-CUTOFF keyword is infinite for nonperiodic systems and 9.0 for periodic systems.

MPOLE-TAPER [real] This keyword allows modification of the cutoff window for atomic multipole potential energy interactions. It is similar in form and action to the TAPER keyword, except that its value applies only to the atomic multipole potential. The default value in the absence of the MPOLE-TAPER keyword is to begin the cutoff window at 0.65 of the corresponding cutoff distance.

MPOLETERM [NONE/ONLY] This keyword controls use of the atomic multipole electrostatics potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

MULTIPOLE [5 lines with: 3 integers & 1 real; 3 reals; 1 real; 2 reals; 3 reals] This keyword provides the values for a set of atomic multipole parameters at a single site. A

complete keyword entry consists of three consecutive lines, the first line containing the MULTIPOLE keyword and the two following lines. The first line contains three integers which define the atom type on which the multipoles are centered, and the Z-axis and X-axis defining atom types for this center. The real number on the first line gives the monopole (atomic charge) in electrons. The second line contains three real numbers which give the X-, Y- and Z-components of the atomic dipole in electron-Å. The final three lines, consisting of one, two and three real numbers give the upper triangle of the traceless atomic quadrupole tensor in electron-Å².

MUTATE [3 integers] This keyword is used to specify atoms to be mutated during free energy perturbation calculations. The first integer modifier gives the atom number of an atom in the current system. The final two modifier values give the atom types corresponding to the $\lambda=0$ and $\lambda=1$ states of the specified atom.

MUTUAL-11-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the induced (mutual) field due to atoms within a polarization group during an induced dipole calculation, *i.e.*, atoms that are in the same polarization group as the atom being polarized. The default value of 1.0 is used, if the MUTUAL-11-SCALE keyword is not given in either the parameter file or the keyfile.

MUTUAL-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the induced (mutual) field due to atoms in 1-2 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups directly connected to the group containing the atom being polarized. The default value of 1.0 is used, if the MUTUAL-12-SCALE keyword is not given in either the parameter file or the keyfile.

MUTUAL-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the induced (mutual) field due to atoms in 1-3 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups separated by one group from the group containing the atom being polarized. The default value of 1.0 is used, if the MUTUAL-13-SCALE keyword is not given in either the parameter file or the keyfile.

MUTUAL-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to the induced (mutual) field due to atoms in 1-4 polarization groups during an induced dipole calculation, *i.e.*, atoms that are in polarization groups separated by two groups from the group containing the atom being polarized. The default value of 1.0 is used, if the MUTUAL-14-SCALE keyword is not given in either the parameter file or the keyfile.

NEIGHBOR-GROUPS This keyword causes the attached atom to be used in determining the charge-charge neighbor distance for all monovalent atoms in the molecular system. Its use causes all monovalent atoms to be treated the same as their attached atoms for purposes of including or scaling 1-2, 1-3 and 1-4 interactions. This option works only for the simple charge-charge electrostatic potential; it does not affect bond dipole or atomic multipole potentials. The NEIGHBOR-GROUPS scheme is similar to that used by some common force fields such as ENCAD.

NEUTRAL-GROUPS The keyword causes the attached atom to be used in determining the charge-charge interaction cutoff distance for all monovalent atoms in the molecular system. Its use reduces cutoff discontinuities by avoiding splitting many of the largest charge separations found in typical molecules. Note that this keyword does not rigorously implement the usual concept of a “neutral group” as used in the literature with

AMBER/OPLS and other force fields. This option works only for the simple charge-charge electrostatic potential; it does not affect bond dipole or atomic multipole potentials.

NEWHESS [integer] Sets the number of algorithmic iterations between recomputation of the Hessian matrix. At present this keyword applies exclusively to optimizations using the Truncated Newton method. The default value in the absence of this keyword is 1, *i.e.*, the Hessian is computed on every iteration.

NEXTITER [integer] Sets the iteration number to be used for the first iteration of the current computation. At present this keyword applies to optimization procedures where its use can effect convergence criteria, timing of restarts, and so forth. The default in the absence of this keyword is to take the initial iteration as iteration 1.

NOVERSION Turns off the use of version numbers appended to the end of filenames as the method for generating filenames for updated copies of an existing file. The presence of this keyword results in direct use of input file names without a search for the highest available version, and requires the entry of specific output file names in many additional cases. By default, in the absence of this keyword, TINKER generates and attaches version numbers in a manner similar to the Digital OpenVMS operating system. For example, subsequent new versions of the file `molecule.xyz` would be written first to the file `molecule.xyz_2`, then to `molecule.xyz_3`, *etc.*

OCTAHEDRON Specifies that the periodic “box” is a truncated octahedron with maximal distance across the truncated octahedron as given by the A-AXIS keyword. All other unit cell and periodic box size-defining keywords are ignored if the OCTAHEDRON keyword is present.

OPBEND [2 integers & 1 real] This keyword provides the values for a single Allinger MM-style out-of-plane angle bending potential parameter. The first integer modifier is the atom class of the central trigonal atom and the second integer is the atom class of the out-of-plane atom. The real number modifier gives the force constant value for the out-of-plane angle. The default units for the force constant are kcal/mole/radian², but this can be controlled via the OPBENDUNIT keyword.

OPBENDTERM [NONE/ONLY] This keyword controls use of the Allinger MM-style out-of-plane bending potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

OPBENDUNIT [real] Sets the scale factor needed to convert the energy value computed by the Allinger MM-style out-of-plane bending potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default of $(\pi/180)^2 = 0.0003046$ is used, if the OPBENDUNIT keyword is not given in the force field parameter file or the keyfile.

OPDIST [4 integers & 1 real] This keyword provides the values for a single out-of-plane distance potential parameter. The first integer modifier is the atom class of the central trigonal atom and the three following integer modifiers are the atom classes of the three attached atoms. The real number modifier is the force constant for the harmonic function of the out-of-plane distance of the central atom. The default units for the force constant are kcal/mole/Å², but this can be controlled via the OPDISTUNIT keyword.

OPDISTTERM [NONE/ONLY] This keyword controls use of the out-of-plane distance potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

OPDISTUNIT [real] Sets the scale factor needed to convert the energy value computed by the out-of-plane distance potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the OPDISTUNIT keyword is not given in the force field parameter file or the keyfile.

OVERWRITE Causes TINKER programs, such as minimizations, that output intermediate coordinate sets to create a single disk file for the intermediate results which is successively overwritten with the new intermediate coordinates as they become available. This keyword is essentially the opposite of the SAVECYCLE keyword.

PARAMETERS [file name] Provides the name of the force field parameter file to be used for the current TINKER calculation. The standard file name extension for parameter files, `.prm`, is an optional part of the file name modifier. The default in the absence of the PARAMETERS keyword is to look for a parameter file with the same base name as the molecular system and ending in the `.prm` extension. If a valid parameter file is not found, the user will be asked to provide a file name interactively.

PIATOM [1 integer & 3 reals] This keyword provides the values for the pisystem MO potential parameters for a single atom class belonging to a pisystem.

PIBOND [2 integers & 2 reals] This keyword provides the values for the pisystem MO potential parameters for a single type of pisystem bond.

PISYSTEM [integer list] This keyword sets the atoms within a molecule that are part of a conjugated π -system. The keyword is followed on the same line by a list of atom numbers and/or atom ranges that constitute the π -system. The Allinger MM force fields use this information to set up an MO calculation used to scale bond and torsion parameters involving π -system atoms.

PME-GRID [3 integers] This keyword sets the dimensions of the charge grid used during particle mesh Ewald summation. The three modifiers give the size along the X-, Y- and Z-axes, respectively. If either the Y- or Z-axis dimensions are omitted, then they are set equal to the X-axis dimension. The default in the absence of the PME-GRID keyword is to set the grid size along each axis to the smallest power of 2, 3 and/or 5 which is at least as large as 1.5 times the axis length in Angstroms. Note that the FFT used by PME is not restricted to, but is most efficient for, grid sizes which are powers of 2, 3 and/or 5.

PME-ORDER [integer] This keyword sets the order of the B-spline interpolation used during particle mesh Ewald summation. A default value of 8 is used in the absence of the PME-ORDER keyword.

POLAR-11-SCALE [real] This keyword provides a multiplicative scale factor that is applied to polarization interactions within a polarization group, *i.e.*, pairs of atoms that are in the same polarization group. The default value of 0.0 is used, if the POLAR-11-SCALE keyword is not given in either the parameter file or the keyfile.

POLAR-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to polarization interactions between 1-2 polarization groups, *i.e.*, pairs of atoms that are in directly connected polarization groups. The default value of 0.0 is used, if the POLAR-12-SCALE keyword is not given in either the parameter file or the keyfile.

POLAR-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to polarization interactions between 1-3 polarization groups, *i.e.*, pairs of atoms that are in polarization groups separated by one other group. The default value of 0.0 is used, if the POLAR-13-SCALE keyword is not given in either the parameter file or the keyfile.

POLAR-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to polarization interactions between 1-4 polarization groups, *i.e.*, pairs of atoms that are in polarization groups separated by two other groups. The default value of 1.0 is used, if the POLAR-14-SCALE keyword is not given in either the parameter file or the keyfile.

POLAR-DAMP [2 reals] Controls the strength of the damping function applied to induced dipoles and dipole polarization interaction energies. The first modifier sets the radius in Angstroms of a hypothetical atom with unit polarizability, while the second modifier sets the scale factor for the exponent of the damping function. The default values for the radius and the scale factor are 1.662 and 1.0, respectively. Damping is eliminated entirely by using this keyword to set the radius value to zero.

POLAR-EPS [real] This keyword sets the convergence criterion applied during computation of self-consistent induced dipoles. The calculation is deemed to have converged when the *rms* change (in Debyes) of the induced dipoles at all polarizable sites is less than the value specified with this keyword. The default value in the absence of the keyword is 10^{-6} Debyes.

POLAR-OLD This keyword selects the polarization damping scheme used in TINKER 3.8 and earlier. Beginning with the 3.9 release, TINKER implements a short range polarization damping method due to Thole. This option is included primarily to allow continued use of the early TINKER polarizable water model based on the originally implemented flat multiplicative damping.

POLAR-SOR [real] Sets a successive overrelaxation (SOR) factor for use in computation of induced atomic dipoles. Optimal values for this keyword will speed the induced dipole calculation, and poor values can result in convergence failure. The default value in the absence of the POLAR-SOR keyword is 0.7 which often a reasonable value when short-range intramolecular polarization is present. For models lacking intramolecular polarization, keyword values closer to 1.0 may be optimal.

POLARIZATION [DIRECT/MUTUAL] Selects between the use of direct and mutual dipole polarization for force fields that incorporate the polarization term. The DIRECT modifier avoids an iterative calculation by using only the permanent electric field in computation of induced dipoles. The MUTUAL option, which is the default in the absence of the POLARIZATION keyword, iterates the induced dipoles to self-consistency.

POLARIZE [1 integer, 1 real & up to 4 integers] This keyword provides the values for a single atomic dipole polarizability parameter. The integer modifier, if positive, gives the atom type number for which a polarizability parameter is to be defined. If the first integer modifier is negative, then the parameter value to follow applies only to the individual atom whose atom number is the negative of the modifier. The real number modifier gives the value

of the dipole polarizability in \AA^3 . The final integer modifiers list the atom type numbers of atoms directly bonded to the current atom and which will be considered to be part of the current atom's polarization group.

POLARIZETERM [NONE/ONLY] This keyword controls use of the atomic dipole polarization potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

POLYMER-CUTOFF [real] Sets the value of an additional cutoff parameter needed for infinite polymer systems. This value must be set to less than half the minimal periodic box dimension and should be greater than the largest possible interatomic distance that can be subject to scaling or exclusion as a local electrostatic or van der Waals interaction. The default in the absence of the POLYMER-CUTOFF keyword is 5.5 Angstroms.

PRINTOUT [integer] A general parameter for iterative procedures such as minimizations that sets the number of iterations between writes of status information to the standard output. The default value in the absence of the keyword is 1, *i.e.*, the calculation status is given every iteration.

RADIUSRULE [ARITHMETIC/GEOMETRIC/CUBIC-MEAN] Sets the functional form of the radius combining rule for heteroatomic van der Waals potential energy interactions. The default in the absence of the RADIUSRULE keyword is to use the arithmetic mean combining rule to get radii for heteroatomic interactions.

RADIUSSIZE [RADIUS/DIAMETER] Determines whether the atom size values given in van der Waals parameters read from VDW keyword statements are interpreted as atomic radius or diameter values. The default in the absence of the RADIUSSIZE keyword is to assume that vdw size parameters are given as radius values.

RADIUSTYPE [R-MIN/SIGMA] Determines whether atom size values given in van der Waals parameters read from VDW keyword statements are interpreted as potential minimum (R_{\min}) or LJ-style sigma (σ) values. The default in the absence of the RADIUSTYPE keyword is to assume that vdw size parameters are given as R_{\min} values.

RANDOMSEED [integer] Followed by an integer value, this keyword sets the initial seed value for the random number generator used by TINKER. Setting RANDOMSEED to the same value as an earlier run will allow exact reproduction of the earlier calculation. (Note that this will not hold across different machine types.) RANDOMSEED should be set to a positive integer less than about 2 billion. In the absence of the RANDOMSEED keyword the seed is chosen "randomly" based upon the number of seconds that have elapsed in the current decade.

RATTLE [BONDS/ANGLES/DIATOMIC/TRIATOMIC/WATER] Invokes the rattle algorithm, a velocity version of shake, on portions of a molecular system during a molecular dynamic calculation. The RATTLE keyword can be followed by any of the modifiers shown, in which case all occurrences of the modifier species are constrained at ideal values taken from the bond and angle parameters of the force field in use. In the absence of any modifier, RATTLE constrains all bonds to hydrogen atoms at ideal bond lengths.

RATTLE-BOND [2 integers] This keyword allows the use of rattle (see above) on a the bond between the two atoms whose numbers are specified on the keyword line. If the two

atoms are involved in a covalent bond, then their distance is constrained to the ideal bond length from the force field. For nonbonded atoms, the rattle constraint fixes their distance at the distance in the input coordinate file.

REACTIONFIELD [2 reals & 1 integer] This keyword provides parameters needed for the reaction field potential energy calculation. The two real modifiers give the radius of the dielectric cavity and the ratio of the bulk dielectric outside the cavity to that inside the cavity. The integer modifier gives the number of terms in the reaction field summation to be used. In the absence of the REACTIONFIELD keyword, the default values are a cavity of radius 1000000 Å, a dielectric ratio of 80 and use of only the first term of the reaction field summation.

REDUCE [real] Specifies the fraction between zero and one by which the path between starting and final conformational state will be shortened at each major cycle of the transition state location algorithm implemented by the SADDLE program. This causes the path endpoints to move up and out of the terminal structures toward the transition state region. In favorable cases, a nonzero value of the REDUCE modifier can speed convergence to the transition state. The default value in the absence of the REDUCE keyword is zero.

RESTRAIN-ANGLE [3 integers & 3 reals] This keyword implements a flat-welled harmonic potential that can be used to restrain the angle between three atoms to lie within a specified angle range. The initial integer modifiers contains the atom numbers of the three atoms whose angle is to be restrained. The first two real number modifiers give the lower and upper bounds in degrees on the allowed angle values. If the angle lies between the lower and upper bounds, the restraint potential is zero. Outside the bounds, a harmonic potential with force constant in kcal/degree² given by the final real modifier is applied. If the force constant is omitted, a default value of 10.0 is used. If all the real modifiers are omitted, then the atoms are restrained to an angle of zero with the default force constant.

RESTRAIN-DISTANCE [2 integers & 3 reals] This keyword implements a flat-welled harmonic potential that can be used to restrain two atoms to lie within a specified distance range. The initial integer modifiers contains the atom numbers of the two atoms to be restrained. The first two real number modifiers give the lower and upper bounds in Angstroms on the allowed distance values. If the interatomic distance lies between the lower and upper bounds, the restraint potential is zero. Outside the bounds, a harmonic potential with force constant in kcal/Å² given by the final real modifier is applied. If the force constant is omitted, a default value of 100.0 is used. If all the real modifiers are omitted, then the atoms are restrained to an interatomic distance of zero with the default force constant.

RESTRAIN-POSITION [1 integer & 4 reals] This keyword provides the ability to restrain an individual atom to a specified coordinate position. The initial integer modifier contains the atom number of the atom to be restrained. The first three real number modifiers give the X-, Y- and Z-coordinates to which the atom is tethered. The final real modifier sets the force constant in kcal/Å² for the harmonic restraint potential. If the force constant is omitted, a default value of 100.0 is used. If all the real modifiers are omitted, then the atom is restrained to the origin with the default force constant.

RESTRAIN-TORSION [4 integers & 3 reals] This keyword implements a flat-welled harmonic potential that can be used to restrain the torsional angle between four atoms to lie within a specified angle range. The initial integer modifiers contains the atom numbers of the four atoms whose torsional angle, computed in the atom order listed, is to be restrained. The first two real number modifiers give the lower and upper bounds in degrees on the

allowed torsional angle values. The angle values given can wrap around across -180 and +180 degrees. Outside the allowed angle range, a harmonic potential with force constant in kcal/degree² given by the final real modifier is applied. If the force constant is omitted, a default value of 1.0 is used. If all the real modifiers are omitted, then the atoms are restrained to a torsional angle of zero with the default force constant.

RESTRAINTERM [NONE/ONLY] This keyword controls use of the restraint potential energy terms. In the absence of a modifying option, this keyword turns on use of these potentials. The NONE option turns off use of these potential energy terms. The ONLY option turns off all potential energy terms except for these terms.

RXNFIELDTERM [NONE/ONLY] This keyword controls use of the reaction field continuum solvation potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

SADDLEPOINT The presence of this keyword allows Newton-style second derivative-based optimization routine used by NEWTON, NEWTROT and other programs to converge to saddlepoints as well as minima on the potential surface. By default, in the absence of the SADDLEPOINT keyword, checks are applied that prevent convergence to stationary points having directions of negative curvature.

SAVE-CYCLE This keyword causes TINKER programs, such as minimizations, that output intermediate coordinate sets to save each successive set to the next consecutively numbered cycle file. The SAVE-CYCLE keyword is the opposite of the OVERWRITE keyword.

SAVE-INDUCED This keyword causes TINKER molecular dynamics calculations that involve polarizable atomic multipoles to save the values of the induced dipole components on each polarizable atom to a separate cycle file. These files are written whenever the atomic coordinate snapshots are written during the dynamics run. Each induced dipole file name contains as a suffix the cycle number followed by the letter u.

SAVE-VELOCITY This keyword causes TINKER molecular dynamics calculations to save the values of the velocity components on each atom to a separate cycle file. These files are written whenever the atomic coordinate snapshots are written during the dynamics run. Each velocity file name contains as a suffix the cycle number followed by the letter v.

SOLVATE [ASP/SASA/ONION/STILL/HCT/ACE/GBSA] Use of this keyword during energy calculations with any of the standard force fields turns on a continuum solvation free energy term. Several algorithms are available based on the modifier used: ASP= Eisenberg-McLachlan ASP method using the Wesson-Eisenberg vacuum-to-water parameters; SASA= the Ooi-Scheraga SASA method; ONION= the original 1990 Still "Onion-shell" GB/SA method; STILL= the 1997 analytical GB/SA method from Still's group; HCT= the pairwise descreening method of Hawkins, Cramer and Truhlar; ACE= the analytical continuum solvation method from the Karplus group; GBSA= equivalent to the STILL modifier. At present, GB/SA-style methods are only valid for force fields that use simple partial charge electrostatics. *The ACE method is not recommended for use in the current version of TINKER; the algorithm is fully implemented in the source code, but is not yet completely parametrized.*

SOLVATETERM [NONE/ONLY] This keyword controls use of the macroscopic solvation potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

SPACEGROUP [name] This keyword selects the space group to be used in manipulation of crystal unit cells and asymmetric units. The name option must be chosen from one of the following currently implemented space groups: P1, P1(-), P21, Cc, P21/a, P21/n, P21/c, C2/c, P212121, Pna21, Pn21a, Cmc21, Pccn, Pbcn, Pbca, P41, I41/a, P4(-)21c, P4(-)m2, R3c, P6(3)/mcm, Fm3(-)m, Im3(-)m.

SPHERE [4 reals, or 1 integer & 1 real] This keyword provides an alternative to the ACTIVE and INACTIVE keywords for specification of subsets of active atoms. If four real number modifiers are provided, the first three are taken as X-, Y- and Z-coordinates and the fourth is the radius of a sphere centered at these coordinates. In this case, all atoms within the sphere at the *start* of the calculation are active *throughout* the calculation, while all other atoms are inactive. Similarly if one integer and real number are given, an “active” sphere with radius set by the real is centered on the system atom with atom number given by the integer modifier. Multiple SPHERE keyword lines can be present in a single keyfile, and the list of active atoms specified by the spheres is cumulative.

STEPMAX [real] This keyword and its modifying value set the maximum size of an individual step during the line search phase of conjugate gradient or truncated Newton optimizations. The step size is computed as the norm of the vector of changes in parameters being optimized. The default value depends on the particular TINKER program, but is usually in the range from 1.0 to 5.0 when not specified via the STEPMAX keyword.

STEPMIN [real] This keyword and its modifying value set the minimum size of an individual step during the line search phase of conjugate gradient or truncated Newton optimizations. The step size is computed as the norm of the vector of changes in parameters being optimized. The default value is usually set to about 10^{-16} when not specified via the STEPMIN keyword.

STRBND [1 integer & 3 reals] This keyword provides the values for a single stretch-bend cross term potential parameter. The integer modifier gives the atom class number for the central atom of the bond angle involved in stretch-bend interactions. The real number modifiers give the force constant values to be used when the central atom of the angle is attached to 0, 1 or 2 additional hydrogen atoms, respectively. The default units for the stretch-bend force constant are kcal/mole/Å-degree, but this can be controlled via the STRBNDUNIT keyword.

STRBNDTERM [NONE/ONLY] This keyword controls use of the bond stretching-angle bending cross term potential energy. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

STRBNDUNIT [real] Sets the scale factor needed to convert the energy value computed by the bond stretching-angle bending cross term potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the STRBNDUNIT keyword is not given in the force field parameter file or the keyfile.

STRTORS [2 integers & 1 real] This keyword provides the values for a single stretch-torsion cross term potential parameter. The two integer modifiers give the atom class numbers for the atoms involved in the central bond of the torsional angles to be parameterized. The real modifier gives the value of the stretch-torsion force constant for all torsional angles with the defined central bond atom classes. The default units for the stretch-torsion force constant can be controlled via the STRTORUNIT keyword.

STRTORTERM [NONE/ONLY] This keyword controls use of the bond stretching-torsional angle cross term potential energy. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

STRTORUNIT [real] Sets the scale factor needed to convert the energy value computed by the bond stretching-torsional angle cross term potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the STRTORUNIT keyword is not given in the force field parameter file or the keyfile.

TAPER [real] This keyword allows modification of the cutoff windows for nonbonded potential energy interactions. The nonbonded terms are smoothly reduced from their standard value at the beginning of the cutoff window to zero at the far end of the window. The far end of the window is specified via the CUTOFF keyword or its potential function specific variants. The modifier value supplied with the TAPER keyword sets the beginning of the cutoff window. The modifier can be given either as an absolute distance value in Angstroms, or as a fraction between zero and one of the CUTOFF distance. The default value in the absence of the TAPER keyword ranges from 0.65 to 0.9 of the CUTOFF distance depending on the type of potential function. The windows are implemented via polynomial-based switching functions, in some cases combined with energy shifting.

TAU-PRESSURE [real] Sets the coupling time in picoseconds for the Groningen-style pressure bath coupling used to control the system pressure during molecular dynamics calculations. A default value of 2.0 is used for TAU-PRESSURE in the absence of the keyword.

TAU-TEMPERATURE [real] Sets the coupling time in picoseconds for the Groningen-style temperature bath coupling used to control the system temperature during molecular dynamics calculations. A default value of 0.1 is used for TAU-TEMPERATURE in the absence of the keyword.

THERMOSTAT [BERENDSEN/ANDERSEN] This keyword selects a thermostat algorithm for use during molecular dynamics. Two modifiers are available, a Berendsen bath coupling method, and an Andersen stochastic collision method. The default in the absence of the THERMOSTAT keyword is to use the BERENDSEN algorithm.

TORSION [4 integers & up to 6 real/real/integer triples] This keyword provides the values for a single torsional angle parameter. The first four integer modifiers give the atom class numbers for the atoms involved in the torsional angle to be defined. Each of the remaining triples of real/real/integer modifiers give the half-amplitude, phase offset in degrees and periodicity of a particular torsional function term, respectively. Periodicities through 6-fold are allowed for torsional parameters.

TORSION4 [4 integers & up to 6 real/real/integer triples] This keyword provides the values for a single torsional angle parameter specific to atoms in 4-membered rings. The first four integer modifiers give the atom class numbers for the atoms involved in the torsional angle to be defined. The remaining triples of real number and integer modifiers operate as described above for the TORSION keyword.

TORSION5 [4 integers & up to 6 real/real/integer triples] This keyword provides the values for a single torsional angle parameter specific to atoms in 5-membered rings. The first four integer modifiers give the atom class numbers for the atoms involved in the torsional angle to be defined. The remaining triples of real number and integer modifiers operate as described above for the TORSION keyword.

TORSIONTERM [NONE/ONLY] This keyword controls use of the torsional angle potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

TORSIONUNIT [real] Sets the scale factor needed to convert the energy value computed by the torsional angle potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the TORSIONUNIT keyword is not given in the force field parameter file or the keyfile.

TORTORTERM [NONE/ONLY] This keyword controls use of the torsion-torsion potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one. This energy term is not implemented in the current version of TINKER.

TRIAL-DISTANCE [CLASSIC/RANDOM/TRICOR/HAVEL integer/PAIRWISE integer] Sets the method for selection of a trial distance matrix during distance geometry computations. The keyword takes a modifier that selects the method to be used. The HAVEL and PAIRWISE modifiers also require an additional integer value that specifies the number of atoms used in metrization and the percentage of metrization, respectively. The default in the absence of this keyword is to use the PAIRWISE method with 100 percent metrization. Further information on the various methods is given with the description of the TINKER distance geometry program.

TRIAL-DISTRIBUTION [real] Sets the initial value for the mean of the Gaussian distribution used to select trial distances between the lower and upper bounds during distance geometry computations. The value given must be between 0 and 1 which represent the lower and upper bounds respectively. This keyword is rarely needed since TINKER will usually be able to choose a reasonable value by default.

TRUNCATE Causes all distance-based nonbond energy cutoffs to be sharply truncated to an energy of zero at distances greater than the value set by the cutoff keyword(s) without use of any shifting, switching or smoothing schemes. At all distances within the cutoff sphere, the full interaction energy is computed.

UREY-CUBIC [real] Sets the value of the cubic term in the Taylor series expansion form of the Urey-Bradley potential energy. The real number modifier gives the value of the

coefficient as a multiple of the quadratic coefficient. The default value in the absence of the UREY-CUBIC keyword is zero; *i.e.*, the cubic Urey-Bradley term is omitted.

UREY-QUARTIC [real] Sets the value of the quartic term in the Taylor series expansion form of the Urey-Bradley potential energy. The real number modifier gives the value of the coefficient as a multiple of the quadratic coefficient. The default value in the absence of the UREY-QUARTIC keyword is zero; *i.e.*, the quartic Urey-Bradley term is omitted.

UREYBRAD [3 integers & 2 reals] This keyword provides the values for a single Urey-Bradley cross term potential parameter. The integer modifiers give the atom class numbers for the three kinds of atoms involved in the angle for which a Urey-Bradley term is to be defined. The real number modifiers give the force constant value for the term and the target value for the 1-3 distance in Å. The default units for the force constant are kcal/mole/Å², but this can be controlled via the UREYUNIT keyword.

UREYTERM [NONE/ONLY] This keyword controls use of the Urey-Bradley potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

UREYUNIT [real] Sets the scale factor needed to convert the energy value computed by the Urey-Bradley potential into units of kcal/mole. The correct value is force field dependent and typically provided in the header of the master force field parameter file. The default value of 1.0 is used, if the UREYUNIT keyword is not given in the force field parameter file or the keyfile.

VDW [1 integer & 3 reals] This keyword provides values for a single van der Waals parameter. The integer modifier, if positive, gives the atom class number for which vdw parameters are to be defined. Note that vdw parameters are given for atom classes, not atom types. The three real number modifiers give the values of the atom size in Å, homoatomic well depth in kcal/mole, and an optional reduction factor for univalent atoms.

VDW-12-SCALE [real] This keyword provides a multiplicative scale factor that is applied to van der Waals potential interactions between 1-2 connected atoms, *i.e.*, atoms that are directly bonded. The default value of 0.0 is used, if the VDW-12-SCALE keyword is not given in either the parameter file or the keyfile.

VDW-13-SCALE [real] This keyword provides a multiplicative scale factor that is applied to van der Waals potential interactions between 1-3 connected atoms, *i.e.*, atoms separated by two covalent bonds. The default value of 0.0 is used, if the VDW-13-SCALE keyword is not given in either the parameter file or the keyfile.

VDW-14-SCALE [real] This keyword provides a multiplicative scale factor that is applied to van der Waals potential interactions between 1-4 connected atoms, *i.e.*, atoms separated by three covalent bonds. The default value of 1.0 is used, if the VDW-14-SCALE keyword is not given in either the parameter file or the keyfile.

VDW-15-SCALE [real] This keyword provides a multiplicative scale factor that is applied to van der Waals potential interactions between 1-5 connected atoms, *i.e.*, atoms separated by four covalent bonds. The default value of 1.0 is used, if the VDW-15-SCALE keyword is not given in either the parameter file or the keyfile.

VDW-CUTOFF [real] Sets the cutoff distance value in Angstroms for van der Waals potential energy interactions. The energy for any pair of van der Waals sites beyond the cutoff distance will be set to zero. Other keywords can be used to select a smoothing scheme near the cutoff distance. The default cutoff distance in the absence of the VDW-CUTOFF keyword is infinite for nonperiodic systems and 9.0 for periodic systems.

VDW-TAPER [real] This keyword allows modification of the cutoff windows for van der Waals potential energy interactions. It is similar in form and action to the TAPER keyword, except that its value applies only to the vdw potential. The default value in the absence of the VDW-TAPER keyword is to begin the cutoff window at 0.9 of the vdw cutoff distance.

VDW14 [1 integer & 2 reals] This keyword provides values for a single van der Waals parameter for use in 1-4 nonbonded interactions. The integer modifier, if positive, gives the atom class number for which vdw parameters are to be defined. Note that vdw parameters are given for atom classes, not atom types. The two real number modifiers give the values of the atom size in Å and the homoatomic well depth in kcal/mole. Reduction factors, if used, are carried over from the VDW keyword for the same atom class.

VDWPR [2 integers & 2 reals] This keyword provides the values for the vdw parameters for a single special heteroatomic pair of atoms. The integer modifiers give the pair of atom class numbers for which special vdw parameters are to be defined. The two real number modifiers give the values of the minimum energy contact distance in Å and the well depth at the minimum distance in kcal/mole.

VDWTERM [NONE/ONLY] This keyword controls use of the van der Waals repulsion-dispersion potential energy term. In the absence of a modifying option, this keyword turns on use of the potential. The NONE option turns off use of this potential energy term. The ONLY option turns off all potential energy terms except for this one.

VDWTYPE [LENNARD-JONES/BUCKINGHAM/BUFFERED-14-7/MM3-HBOND/GAUSSIAN] Sets the functional form for the van der Waals potential energy term. The text modifier gives the name of the functional form to be used. The GAUSSIAN modifier value implements a two or four Gaussian fit to the corresponding Lennard-Jones function for use with potential energy smoothing schemes. The default in the absence of the VDWTYPE keyword is to use the standard two parameter Lennard-Jones function.

VERBOSE Turns on printing of secondary and informational output during a variety of TINKER computations; a subset of the more extensive output provided by the DEBUG keyword.

WALL [real] Sets the radius of a spherical boundary used to maintain droplet boundary conditions. The real modifier specifies the desired approximate radius of the droplet. In practice, an artificial van der Waals wall is constructed at a fixed buffer distance of 2.5 Å outside the specified radius. The effect is that atoms which attempt to move outside the region defined by the droplet radius will be forced toward the center.

WRITEOUT [integer] A general parameter for iterative procedures such as minimizations that sets the number of iterations between writes of intermediate results (such as the current coordinates) to disk file(s). The default value in the absence of the keyword is 1, *i.e.*, the intermediate results are written to file on every iteration. Whether successive results are saved to new files or replace previously written intermediate results is controlled by the OVERWRITE and SAVECYCLE keywords.

8. Notes on Special Features & Methods

This section contains several short notes with further information about TINKER methodology, algorithms and special features. The discussion is not intended to be exhaustive, but rather to explain features and capabilities so that users can make more complete use of the package.

FILE VERSION NUMBERS

All of the input and output file types routinely used by the TINKER package are capable of existing as multiple versions of a base file name. For example, if the program XYZINT is run on the input file `molecule.xyz`, the output internal coordinates file will be written to `molecule.int`. If a file named `molecule.int` is already present prior to running XYZINT, then the output will be written instead to the next available version, in this case to `molecule.int_2`. In fact the output is generally written to the lowest available, previously unused version number (`molecule.int_3`, `molecule.int_4`, *etc.*, as high as needed). Input file names are handled similarly. If simply `molecule` or `molecule.xyz` is entered as the input file name upon running XYZINT, then the highest version of `molecule.xyz` will be used as the actual input file. If an explicit version number is entered as part of the input file name, then the specified version will be used as the input file.

The version number scheme will be recognized by many older users as a holdover from the VMS origins of the first version of the TINKER software. It has been maintained to make it easier to chain together multiple calculations that may create several new versions of a given file, and to make it more difficult to accidentally overwrite a needed result. The version scheme applies to most uses of many common TINKER file types such as `.xyz`, `.int`, `.key`, `.arc`. It is not used when an overwritten file "update" is obviously the correct action, for example, the `.dyn` molecular dynamics restart files. For those users who prefer a more Unix-like operation, and do not desire use of file versions, this feature can be turned off by adding the NOVERSION keyword to the applicable TINKER keyfile.

The version scheme as implemented in TINKER does have two known quirks. First, it becomes impossible to directly use the original unversioned copy of a file if higher version numbers are present. For example, if the files `molecule.xyz` and `molecule.xyz_2` both exist, then `molecule.xyz` cannot be accessed as input by XYZINT. If `molecule.xyz` is entered in response to the input file name question, `molecule.xyz_2` (or the highest present version number) will be used as input. The only workaround is to copy or rename `molecule.xyz` to something else, say `molecule.new`, and use that name for the input file. Secondly, missing version numbers always end the search for the highest available version number; *i.e.*, version numbers are assumed to be consecutive and without gaps. For example, if `molecule.xyz`, `molecule.xyz_2` and `molecule.xyz_4` are present, but not `molecule.xyz_3`, then `molecule.xyz_2` will be used as input to XYZINT if `molecule` is given as the input file name. Similarly, output files will fill in gaps in an already existing set of file versions.

COMMAND LINE OPTIONS

Many operating systems or compiler supplied-libraries make available something like the standard Unix `iargc` and `getarg` routines for capturing command line arguments. On these machines most of the TINKER programs support a selection of command line arguments and options. The name of the keyfile to be used for a calculation is read from the

argument following a **-k** (equivalent to either **-key** or **-keyfile**, case insensitive) command line argument. Note that the **-k** options can appear anywhere on the command line following the executable name. All other command line arguments, excepting the name of the executable program itself, are treated as input arguments. These input arguments are read from left to right and interpreted in order as the answers to questions that would be asked by an interactive invocation of the same TINKER program. For example, the following command line:

```
newton molecule -k test a a 0.01
```

will invoke the NEWTON program on the structure file `molecule.xyz` using the keyfile `test.key`, automatic mode [**a**] for both the method and preconditioning, and 0.01 for the RMS gradient per atom termination criterion in kcal/mole/Å. Provided that the force field parameter set, *etc.* is provided in `test.key`, the above computation will proceed directly from the command line invocation without further interactive input.

USE ON MICROSOFT WINDOWS SYSTEMS

TINKER executables for Microsoft PC systems should be run from the DOS Prompt window available under the various versions of Windows. The TINKER executable directory should be added to your path via the `autoexec.bat` file or similar. If using Win2000, set the number of scrollable lines in the DOS Prompt window to a very large number, so that you will be able to inspect screen output after it flies by. With Win95/98, these DOS Prompt windows are only able to scroll a small number of lines (*amazing!*), so TINKER programs which generate large amounts of screen output should be run such that output will be redirected to a file. This can be accomplished by running the TINKER program in batch mode or by using the Unix-like output redirection build into DOS. For example, the command:

```
dynamic < molecule.inp > molecule.log
```

will run the TINKER dynamic program taking input from the file `molecule.inp` and sending output to `molecule.log`. Also note that command line options as described above are available with the distributed TINKER executables.

Another alternative, particularly attractive to those already familiar with Linux or Unix systems, is to download the Cygwin package currently available under GPL license from the site <http://source.redhat.com/cygwin/>. The cygwin tools provide many of the GNU tools, including a bash shell window from which TINKER programs can be run.

If the distributed TINKER executables are directly from Windows by double clicking on the program icon, then the program will run in its own window. However, upon completion of the program the window will close and screen output will be lost. Any output files written by the program will, of course, still be available. The Windows behavior can be changed by adding the EXIT-PAUSE keyword to the keyfile.

USE ON APPLE MACINTOSH SYSTEMS

The TINKER executables can be run under MacOS by double clicking on a program icon. The program will run in its own window to which all "screen" output will be directed. Upon program termination the window will remain active pending a final return entered by the user which will close the window. Prior to the final return, the contents of the screen

window can be saved to a file via the clipboard for permanent storage. Note that Macintosh uses a colon instead of a forward- or back-slash as the directory separator, so keyfiles transferred from other machines will need to be altered accordingly.

ATOM TYPES VS. ATOM CLASSES

Manipulation of atom types and the proliferation of parameters as atoms are further subdivided into new types is the bane of force field calculation. For example, if each topologically distinct atom arising from the 20 natural amino acids is given a different atom type, then about 300 separate type are required (this ignores the different N- and C-terminal forms of the residues, diastereotopic hydrogens, *etc.*). However, all these types lead to literally thousands of different force field parameters. In fact, there are many thousands of distinct torsional parameters alone. It is impossible at present to fully optimize each of these parameters; and even if we could, a great many of the parameters would be nearly identical. Two somewhat complimentary solutions are available to handle the proliferation of parameters. The first is to specify the molecular fragments to which a given parameter can be applied in terms of a chemical structure language, SMILES strings for example. Some commercial systems, such as the TRIPOS Sybyl software, make use of such a scheme to parse structures and assign force field parameters.

A second general approach is to use hierarchical cascades of parameter groups. TINKER uses a simple version of this scheme. Each TINKER force field atom has both an atom *type* number and an atom *class* number. The types are subsets of the atom classes, *i.e.*, several different atom types can belong to the same atom class. Force field parameters that are somewhat less sensitive to local environment, such as local geometry terms, are then provided and assigned based on atom class. Other energy parameters, such as electrostatic parameters, that are very environment dependent are assigned over the atom types. This greatly reduces the number of independent multiple-atom parameters like the four-atom torsional parameters.

CALCULATIONS ON PARTIAL STRUCTURES

Two methods are available for performing energetic calculations on portions or substructures within a full molecular system. TINKER allows division of the entire system into *active* and *inactive* parts which can be defined via keywords. In subsequent calculations, such as minimization or dynamics, only the active portions of the system are allowed to move. The force field engine responds to the active/inactive division by computing all energetic interactions involving at least one active atom; *i.e.*, any interaction whose energy can change with the motion of one or more active atoms is computed.

The second method for partial structure computation involves dividing the original system into a set of atom *groups*. As before, the groups can be specified via appropriate keywords. The current TINKER implementation allows specification of up to a maximum number of groups as given in the `sizes.i` dimensioning file. The groups must be disjoint in that no atom can belong to more than one group. Further keywords allow the user to specify which intra- and intergroup sets of energetic interactions will contribute to the total force field energy. Weights for each set of interactions in the total energy can also be input. A specific energetic interaction is assigned to a particular intra- or intergroup set if all the atoms involved in the interaction belong to the group (intra-) or pair of groups (inter-). Interactions involving atoms from more than two groups are not computed.

Note that the groups method and active/inactive method use different assignment procedures for individual interactions. The active/inactive scheme is intended for situations where only a portion of a system is allowed to move, but the total energy needs to reflect the presence of the remaining inactive portion of the structure. The groups method is intended for use in rigid body calculations, and is needed for certain kinds of free energy perturbation calculations.

METAL COMPLEXES AND HYPERVALENT SPECIES

The distribution version of TINKER comes dimensioned for a maximum atomic coordination number of four as needed for standard organic compounds. In order to use TINKER for calculations on species containing higher coordination numbers, simply change the value of the parameter `maxval` in the master dimensioning file `sizes.i` and rebuilt the package. Note that this parameter value should not be set larger than necessary since large values can slow the execution of portions of some TINKER programs.

Many molecular mechanics approaches to inorganic and metal structures use an angle bending term which is softer than the usual harmonic bending potential. TINKER implements a Fourier bending term similar to that used by the Landis group's SHAPES force field. The parameters for specific Fourier angle terms are supplied via the `ANGLEF` parameter and keyword format. Note that a Fourier term will only be used for a particular angle if a corresponding harmonic angle term is not present in the parameter file.

We are now collaborating with Anders Carlsson's group in St. Louis to add his transition metal ligand field term to TINKER. Support for this additional potential functional form is already in the TINKER source code, and we plan to release the energy routines after further testing and parameterization.

NEIGHBOR METHODS FOR NONBONDED TERMS

In addition to standard double loop methods, the Method of Lights is available to speed neighbor searching. This method based on taking intersections of sorted atom lists can be much faster for problems where the cutoff distance is significantly smaller than half the maximal cell dimension. The current version of TINKER does not implement the "neighbor list" schemes common to many other simulation packages.

PERIODIC BOUNDARY CONDITIONS

Both spherical cutoff images or replicates of a cell are supported by all TINKER programs that implement periodic boundary conditions. Whenever the cutoff distance is too large for the minimum image to be the only relevant neighbor (*i.e.*, half the minimum box dimension for orthogonal cells), TINKER will automatically switch from the image formalism to use of replicated cells.

DISTANCE CUTOFFS FOR ENERGY FUNCTIONS

Polynomial energy switching over a window is used for terms whose energy is small near the cutoff distance. For monopole electrostatic interactions, which are quite large in typical cutoff ranges, a two polynomial multiplicative-additive shifted energy switch unique to TINKER is applied. The TINKER method is similar in spirit to the force switching methods of Steinbach and Brooks, *J. Comput. Chem.*, **15**, 667-683 (1994). While the particle mesh Ewald method is preferred when periodic boundary conditions are present, TINKER's

shifted energy switch with reasonable switching windows is quite satisfactory for most routine modeling problems. The shifted energy switch minimizes the perturbation of the energy and the gradient at the cutoff to acceptable levels. Problems should arise only if the property you wish to monitor is known to require explicit inclusion of long range components (*i.e.*, calculation of the dielectric constant, *etc.*).

EWALD SUMMATION METHODS

TINKER contains a versions of the Ewald summation technique for inclusion of long range electrostatic interactions via periodic boundaries. The particle mesh Ewald (PME) method is available for simple charge-charge potentials, while regular Ewald is provided for polarizable atomic multipole interactions. The accuracy and speed of the regular and PME calculations is dependent on several interrelated parameters. For both methods, the Ewald coefficient and real-space cutoff distance must be set to reasonable and complementary values. Additional control variables for regular Ewald are the fractional coverage and number of vectors used in reciprocal space. For PME the additional control values are the B-spline order and charge grid dimensions. Complete control over all of these parameters is available via the TINKER keyfile mechanism. By default TINKER will select a set of parameters which provide a reasonable compromise between accuracy and speed, but these should be checked and modified as necessary for each individual system.

CONTINUUM SOLVATION MODELS

Several alternative continuum solvation algorithms are contained within TINKER. All of these are accessed via the SOLVATE keyword and its modifiers. Two simple surface area methods are implemented: the ASP method of Eisenberg and McLachlan, and the SASA method from Scheraga's group. These methods are applicable to any of the standard TINKER force fields. Various schemes based on the generalized Born formalism are also available: the original 1990 numerical "Onion-shell" GB/SA method from Still's group, the 1997 analytical GB/SA method also due to Still, a pairwise descreening algorithm originally proposed by Hawkins, Cramer and Truhlar, and the analytical continuum solvation (ACE) method of Schaefer and Karplus. At present, the generalized Born methods should only be used with force fields having simple partial charge electrostatic interactions.

Some further comments are in order regarding the GB/SA-style solvation models. The "Onion-shell" model is provided mostly for comparison purposes. It uses an exact, analytical surface area calculation for the cavity term and the numerical scheme described in the original paper for the polarization term. This method is very slow, especially for large systems, and does not contain the contribution of the Born radii chain rule term to the first derivatives. We recommend its use only for single-point energy calculations. The other GB/SA methods ("analytical" Still, H-C-T pairwise descreening, and ACE) use an approximate cavity term based on Born radii, and do contain fully correct derivatives including the Born radii chain rule contribution. These methods all scale in CPU time with the square of the size of the system, and can be used with minimization, molecular dynamics and large molecules.

Finally, we note that the ACE solvation model should not be used with the current version of TINKER. The algorithm is fully implemented in the source code, but parameterization is not complete. As of late 2000, parameter values are only available in the literature for use of ACE with the older CHARMM19 force field. We plan to develop values for use with more modern all-atom force fields, and these will be incorporated into TINKER sometime in the future.

POLARIZABLE MULTIPOLE ELECTROSTATICS

Atomic multipole electrostatics through the quadrupole moment is supported by the current version of TINKER, as is either mutual or direct dipole polarization. Ewald summation is available for inclusion of long range interactions. Calculations are implemented via a mixture of the CCP5 algorithms of W. Smith and the Applequist-Dykstra Cartesian polytensor method. At present analytical energy and Cartesian gradient code is provided.

The TINKER package allows intramolecular polarization to be treated via a version of the interaction damping scheme of Thole. To implement the Thole scheme, it is necessary to set all the `mutual-1x-scale` keywords to a value of one. The other polarization scaling keyword series, `direct-1x-scale` and `polar-1x-scale`, can be set independently to enable a wide variety of polarization models. In order to use an Applequist-style model without polarization damping, simply set the `polar-damp` keyword to zero.

POTENTIAL ENERGY SMOOTHING

Versions of our Potential Smoothing and Search (PSS) methodology have been implemented within TINKER. This methods belong to the same general family as Scheraga's Diffusion Equation Method, Straub's Gaussian Density Annealing, Shalloway's Packet Annealing and Verschelde's Effective Diffused Potential, but our algorithms reflect our own ongoing research in this area. In many ways the TINKER potential smoothing methods are the deterministic analog of stochastic simulated annealing. The PSS algorithms are very powerful, but are relatively new and are still undergoing modification, testing and calibration within our research group. This version of TINKER also includes a basin-hopping conformational scanning algorithm in the program SCAN which is particularly effective on smoothed potential surfaces.

DISTANCE GEOMETRY METRIZATION

A much improved and very fast random pairwise metrization scheme is available which allows good sampling during trial distance matrix generation without the usual structural anomalies and CPU constraints of other metrization procedures. An outline of the methodology and its application to NMR NOE-based structure refinement is described in the paper by Hodsdon, *et al.* in *J. Mol. Biol.*, **264**, 585-602 (1996). We have obtained good results with something like the keyword phrase `trial-distribution pairwise 5`, which performs 5% partial random pairwise metrization. For structures over several hundred atoms, a value less than 5 for the percentage of metrization should be fine.

9. Descriptions of TINKER Routines

The distribution version of the TINKER package contains over 600 separate programs, subroutines and functions. This section contains a brief description of the purpose of most of these code units. Further information can be found in the comments located at the top of each source code file.

ACTIVE Subroutine

"active" sets the list of atoms that are used during each potential energy function calculation

ADDBASE Subroutine

"addbase" builds the Cartesian coordinates for a single nucleic acid base; coordinates are read from the Protein Data Bank file or found from internal coordinates, then atom types are assigned and connectivity data generated

ADDBOND Subroutine

"addbond" adds entries to the attached atoms list in order to generate a direct connection between two atoms

ADDSIDE Subroutine

"addside" builds the Cartesian coordinates for a single amino acid side chain; coordinates are read from the Protein Data Bank file or found from internal coordinates, then atom types are assigned and connectivity data generated

ADJACENT Function

"adjacent" finds an atom connected to atom "i1" other than atom "i2"; if no such atom exists, then the closest atom in space is returned

ALCHEMY Program

"alchemy" computes the free energy difference corresponding to a small perturbation by Boltzmann weighting the potential energy difference over a number of sample states; current version (incorrectly) considers the charge energy to be intermolecular in finding the perturbation energies

ANALYSIS Subroutine

"analysis" calls the series of routines needed to calculate the potential energy and perform energy partitioning analysis in terms of type of interaction or atom number

ANALYZ4 Subroutine

"analyz4" prints the energy to 4 decimal places and number of interactions for each component of the potential energy

ANALYZ6 Subroutine

"analyze6" prints the energy to 6 decimal places and number of interactions for each component of the potential energy

ANALYZ8 Subroutine

"analyze8" prints the energy to 8 decimal places and number of interactions for each component of the potential energy

ANALYZE Program

"analyze" computes and displays the total potential; options are provided to partition the energy by atom or by potential function type; parameters used in computing interactions can also be displayed by atom; output of large energy interactions and of electrostatic and inertial properties is available

ANGLES Subroutine

"angles" finds the total number of bond angles and stores the atom numbers of the atoms defining each angle; for each angle to a tricoordinate central atom, the third bonded atom is stored for use in out-of-plane bending

ANNEAL Program

"anneal" performs a simulated annealing protocol by means of variable temperature molecular dynamics using either linear, exponential or sigmoidal cooling schedules

ANORM Function

"anorm" finds the norm (length) of a vector; used as a service routine by the Connolly surface area and volume computation

ARCHIVE Program

"archive" is a utility program for coordinate files which concatenates multiple coordinate sets into a single archive file, or extracts individual coordinate sets from an archive

ASET Subroutine

ATOMYZE Subroutine

"atomyze" prints the potential energy components broken down by atom and to a choice of precision

ATTACH Subroutine

"attach" generates lists of 1-3, 1-4 and 1-5 connectivities starting from the previously determined list of attached atoms (ie, 1-2 connectivity)

BASEFILE Subroutine

"basefile" extracts from an input filename the portion consisting of any directory name and the base filename

BEEMAN Subroutine

"beeman" performs a single molecular dynamics time step by means of a Beeman multistep recursion formula; the actual coefficients are Brooks' "Better Beeman" values

BETACF Function

"betacf" computes a rapidly convergent continued fraction needed by routine "betai" to evaluate the cumulative Beta distribution

BETAI Function

"betai" evaluates the cumulative Beta distribution function as the probability that a random variable from a distribution with Beta parameters "a" and "b" will be less than "x"

BIGBLOCK Subroutine

"bigblock" replicates the coordinates of a single unit cell to give a larger block of repeated units

BMAX Function

BNDERR Function

"bnderr" is the distance bound error function and derivatives; this version implements the original and Havel's normalized lower bound penalty, the normalized version is preferred when lower bounds are small (as with NMR NOE restraints), the original penalty is needed if large lower bounds are present

BONDS Subroutine

"bonds" finds the total number of covalent bonds and stores the atom numbers of the atoms defining each bond

BORN Subroutine

"born" computes the Born radius of each atom for use with the various GB/SA solvation models

BORN1 Subroutine

"born1" computes derivatives of the Born radii with respect to atomic coordinates and increments total energy derivatives and virial components for potentials involving Born radii

BOUNDS Subroutine

"bounds" finds the center of mass of each molecule, translates any stray molecules back into the periodic box, and saves the offset of each atom relative to the molecular center of mass

BSET Subroutine

BSPLINE Subroutine

"bspline" calculates the coefficients for an n-th order B-spline approximation

BSPLINE1 Subroutine

"bspline1" calculates the coefficients and derivative coefficients for an n-th order B-spline approximation

BSSSTEP Subroutine

CALENDAR Subroutine

"calendar" returns the current time as a set of integer values representing the year, month, day, hour, minute and second

CELLATOM Subroutine

"cellatom" completes the addition of a symmetry related atom to a unit cell by updating the atom type and attachment arrays

CENTER Subroutine

"center" moves the weighted centroid of each coordinate set to the origin during least squares superposition

CERROR Subroutine

"cerror" is the error handling routine for the Connolly surface area and volume computation

CFFTB Subroutine

"cftb" computes the backward complex discrete Fourier transform, the Fourier synthesis

CFFTB1 Subroutine

CFFTF Subroutine

"cftf" computes the forward complex discrete Fourier transform, the Fourier analysis

CFFTF1 Subroutine

CFFTI Subroutine

"cfti" initializes the array "wsave" which is used in both forward and backward transforms; the prime factorization of "n" together with a tabulation of the trigonometric functions are computed and stored in "wsave"

CFFTI1 Subroutine

CHIRER Function

"chirer" computes the chirality error and its derivatives with respect to atomic Cartesian coordinates as a sum the squares of deviations of chiral volumes from target values

CHKSIZE Subroutine

"chksize" computes a measure of overall global structural expansion or compaction from the number of excess upper or lower bounds matrix violations

CHKTREE Subroutine

"chktree" tests a minimum energy structure to see if it belongs to the correct progenitor in the existing map

CHKXYZ Subroutine

"chkxyz" finds any pairs of atoms with identical Cartesian coordinates, and prints a warning message

CHOLESKY Subroutine

"cholesky" uses a modified Cholesky method to solve the linear system $Ax = b$, returning "x" in "b"; "A" is assumed to be a real symmetric positive definite matrix with diagonal and upper triangle stored by rows in "A"; thus the actual size of the passed portion of "A" is $nvar*(nvar+1)/2$

CIRPLN Subroutine

CJKM Function

CLIMBER Subroutine

CLIMBRGD Subroutine

CLIMBROT Subroutine

CLIMBTOR Subroutine

CLIMBXYZ Subroutine

CLOCK Subroutine

"clock" determines elapsed CPU time in seconds since the start of the job

CLUSTER Subroutine

"cluster" gets the partitioning of the system into groups and stores a list of the group to which each atom belongs

COLUMN Subroutine

"column" takes the off-diagonal Hessian elements stored as sparse rows and sets up indices to allow column access

COMMAND Subroutine

"command" uses the standard Unix-like `iargc/getarg` routines to get the number and values of arguments specified on the command line at program runtime

COMPRESS Subroutine

"compress" transfers only the non-buried tori from the temporary tori arrays to the final tori arrays

CONNECT Subroutine

"connect" sets up the attached atom arrays starting from a set of internal coordinates

CONNOLLY Subroutine

"connolly" uses the algorithms from the AMS/VAM programs of Michael Connolly to compute the analytical molecular surface area and volume of a collection of spherical atoms; thus it implements Fred Richards' molecular surface definition as a set of analytically defined spherical and toroidal polygons

CONTACT Subroutine

"contact" constructs the contact surface, cycles and convex faces

CONTROL Subroutine

"control" gets initial values for parameters that determine the output style and information level provided by TINKER

COORDS Subroutine

"coords" converts the three principal eigenvalues/vectors from the metric matrix into atomic coordinates, and calls a routine to compute the rms deviation from the bounds

CORRELATE Program

"correlate" computes the time correlation function of some user-supplied property from individual snapshot frames taken from a molecular dynamics or other trajectory

CRYSTAL Program

"crystal" is a utility program which converts between fractional and Cartesian coordinates, and can generate full unit cells from asymmetric units

CUTOFFS Subroutine

"cutoffs" initializes and stores spherical energy cutoff distance windows, Hessian element and Ewald sum cutoffs, and the pairwise neighbor generation method

D1D2 Function

DELETE Subroutine

"delete" removes a specified atom from the Cartesian coordinates list and shifts the remaining atoms

DEPTH Function

DFTMOD Subroutine

"dftmod" computes the modulus of the discrete Fourier transform of "bsarray", storing it into "bsmod"

DIAGQ Subroutine

"diagq" is a matrix diagonalization routine which is derived from the classical given, housec, and eigen algorithms with several modifications to increase the efficiency and accuracy

DIFFEQ Subroutine

"diffeq" performs the numerical integration of an ordinary differential equation using an adaptive stepsize method to solve the corresponding coupled first-order equations of the general form $dy_i/dx = f(x, y_1, \dots, y_n)$ for $y_i = y_1, \dots, y_n$

DIFFUSE Program

"diffuse" finds the self-diffusion constant for a homogeneous liquid via the Einstein relation from a set of stored molecular dynamics frames; molecular centers of mass are unfolded and mean squared displacements are computed versus time separation

DIST2 Function

"dist2" finds the distance squared between two points; used as a service routine by the Connolly surface area and volume computation

DISTGEOM Program

"distgeom" uses a metric matrix distance geometry procedure to generate structures with interpoint distances that lie within specified bounds, with chiral centers that maintain chirality, and with torsional angles restrained to desired values; the user also has the ability to interactively inspect and alter the triangle smoothed bounds matrix prior to embedding

DMDUMP Subroutine

"dmdump" puts the distance matrix of the final structure into the upper half of a matrix, the distance of each atom to the centroid on the diagonal, and the individual terms of the bounds errors into the lower half of the matrix

DOCUMENT Program

"document" generates a formatted description of all the code modules or common blocks, a listing of all valid keywords, a list of include file dependencies as needed by a Unix-style Makefile, or a formatted force field parameter set summary

DOT Function

"dot" finds the dot product of two vectors

DSTMAT Subroutine

"dstmat" selects a distance matrix containing values between the previously smoothed upper and lower bounds; the distance values are chosen from uniform distributions, in a triangle correlated fashion, or using random partial metrization

DYNAMIC Program

"dynamic" computes a molecular dynamics trajectory in any of several statistical mechanical ensembles with optional periodic boundaries and optional coupling to temperature and pressure baths alternatively a stochastic dynamics trajectory can be generated

EANGANG Subroutine

"eangang" calculates the angle-angle potential energy

EANGANG1 Subroutine

"eangang1" calculates the angle-angle potential energy and first derivatives with respect to Cartesian coordinates

EANGANG2 Subroutine

"eangang2" calculates the angle-angle potential energy second derivatives with respect to Cartesian coordinates using finite difference methods

EANGANG2A Subroutine

"eangang2a" calculates the angle-angle first derivatives for a single interaction with respect to Cartesian coordinates; used in computation of finite difference second derivatives

EANGANG3 Subroutine

"eangang3" calculates the angle-angle potential energy; also partitions the energy among the atoms

EANGLE Subroutine

"eangle" calculates the angle bending potential energy; projected in-plane angles at trigonal centers or Fourier angle bending terms are optionally used

EANGLE1 Subroutine

"eangle1" calculates the angle bending potential energy and the first derivatives with respect to Cartesian coordinates; projected in-plane angles at trigonal centers or Fourier angle bending terms are optionally used

EANGLE2 Subroutine

"eangle2" calculates second derivatives of the angle bending energy for a single atom using a mixture of analytical and finite difference methods; projected in-plane angles at trigonal centers or Fourier angle bending terms are optionally used

EANGLE2A Subroutine

"eangle2a" calculates bond angle bending potential energy second derivatives with respect to Cartesian coordinates

EANGLE2B Subroutine

"eangle2b" computes projected in-plane bending first derivatives for a single angle with respect to Cartesian coordinates; used in computation of finite difference second derivatives

EANGLE3 Subroutine

"eangle3" calculates the angle bending potential energy, also partitions the energy among the atoms; projected in-plane angles at trigonal centers or Fourier angle bending terms are optionally used

EBOND Subroutine

"ebond" calculates the bond stretching energy

EBOND1 Subroutine

"ebond1" calculates the bond stretching energy and first derivatives with respect to Cartesian coordinates

EBOND2 Subroutine

"ebond2" calculates second derivatives of the bond stretching energy for a single atom at a time

EBOND3 Subroutine

"ebond3" calculates the bond stretching energy; also partitions the energy among the atoms

EBUCK Subroutine

"ebuck" calculates the Buckingham exp-6 van der Waals energy

EBUCK0A Subroutine

"ebuck0a" calculates the Buckingham exp-6 van der Waals energy using a pairwise double loop

EBUCK0B Subroutine

"ebuck0b" calculates the Buckingham exp-6 van der Waals energy using the method of lights to locate neighboring atoms

EBUCK1 Subroutine

"ebuck1" calculates the Buckingham exp-6 van der Waals energy and its first derivatives with respect to Cartesian coordinates

EBUCK1A Subroutine

"ebuck1a" calculates the Buckingham exp-6 van der Waals energy and its first derivatives with respect to Cartesian coordinates using a pairwise double loop

EBUCK1B Subroutine

"ebuck1b" calculates the Buckingham exp-6 van der Waals energy and its first derivatives with respect to Cartesian coordinates using the method of lights to locate neighboring atoms

EBUCK2 Subroutine

"ebuck2" calculates the Buckingham exp-6 van der Waals second derivatives for a single atom at a time

EBUCK3 Subroutine

"ebuck3" calculates the Buckingham exp-6 van der Waals energy and partitions the energy among the atoms

EBUCK3A Subroutine

"ebuck3a" calculates the Buckingham exp-6 van der Waals energy and partitions the energy among the atoms using a pairwise double loop

EBUCK3B Subroutine

"ebuck3b" calculates the Buckingham exp-6 van der Waals energy and also partitions the energy among the atoms using the method of lights to locate neighboring atoms

ECHARGE Subroutine

"echarge" calculates the charge-charge interaction energy

ECHARGE0A Subroutine

"echarge0a" calculates the charge-charge interaction energy using a pairwise double loop

ECHARGE0B Subroutine

"echarge0b" calculates the charge-charge interaction energy using the method of lights to locate neighboring atoms

ECHARGE0C Subroutine

"echarge0c" calculates the charge-charge interaction energy for use with potential smoothing methods

ECHARGE0D Subroutine

"echarge0d" calculates the charge-charge interaction energy using a particle mesh Ewald summation

ECHARGE1 Subroutine

"echarge1" calculates the charge-charge interaction energy and first derivatives with respect to Cartesian coordinates

ECHARGE1A Subroutine

"echarge1a" calculates the charge-charge interaction energy and first derivatives with respect to Cartesian coordinates using a pairwise double loop

ECHARGE1B Subroutine

"echarge1b" calculates the charge-charge interaction energy and first derivatives with respect to Cartesian coordinates using the method of lights to locate neighboring atoms

ECHARGE1C Subroutine

"echarge1c" calculates the charge-charge interaction energy and first derivatives with respect to Cartesian coordinates for use with potential smoothing methods

ECHARGE1D Subroutine

"echarge1d" calculates the charge-charge interaction energy and first derivatives with respect to Cartesian coordinates using a particle mesh Ewald summation

ECHARGE2 Subroutine

"echarge2" calculates second derivatives of the charge-charge interaction energy for a single atom

ECHARGE2A Subroutine

"echarge2a" calculates second derivatives of the charge-charge interaction energy for a single atom using a pairwise double loop

ECHARGE2B Subroutine

"echarge2b" calculates second derivatives of the charge-charge interaction energy for a single atom for use with potential smoothing methods

ECHARGE2C Subroutine

"echarge2c" calculates second derivatives of the charge-charge interaction energy for a single atom using a particle mesh Ewald summation

ECHARGE3 Subroutine

"echarge3" calculates the charge-charge interaction energy and partitions the energy among the atoms

ECHARGE3A Subroutine

"echarge3a" calculates the charge-charge interaction energy and partitions the energy among the atoms using a pairwise double loop

ECHARGE3B Subroutine

"echarge3b" calculates the charge-charge interaction energy and partitions the energy among the atoms using the method of lights to locate neighboring atoms

ECHARGE3C Subroutine

"echarge3c" calculates the charge-charge interaction energy and partitions the energy among the atoms for use with potential smoothing methods

ECHARGE3D Subroutine

"echarge3d" calculates the charge-charge interaction energy and partitions the energy among the atoms using a particle mesh Ewald summation

ECHGDPL Subroutine

"echgdpl" calculates the charge-dipole interaction energy

ECHGDPL1 Subroutine

"echgdpl1" calculates the charge-dipole interaction energy and first derivatives with respect to Cartesian coordinates

ECHGDPL2 Subroutine

"echgdpl2" calculates second derivatives of the charge-dipole interaction energy for a single atom

ECHGDPL3 Subroutine

"echgdpl3" calculates the charge-dipole interaction energy; also partitions the energy among the atoms

EDIPOLE Subroutine

"edipole" calculates the dipole-dipole interaction energy

EDIPOLE1 Subroutine

"edipole1" calculates the dipole-dipole interaction energy and first derivatives with respect to Cartesian coordinates

EDIPOLE2 Subroutine

"edipole2" calculates second derivatives of the dipole-dipole interaction energy for a single atom

EDIPOLE3 Subroutine

"edipole3" calculates the dipole-dipole interaction energy; also partitions the energy among the atoms

EGAUSS Subroutine

"egauss" calculates the Gaussian expansion van der Waals interaction energy

EGAUSS1 Subroutine

"egauss1" calculates the Gaussian expansion van der Waals interaction energy and its first derivatives with respect to Cartesian coordinates

EGAUSS2 Subroutine

"egauss2" calculates the Gaussian expansion van der Waals second derivatives for a single atom at a time

EGAUSS3 Subroutine

"egauss3" calculates the Gaussian expansion van der Waals interaction energy and partitions the energy among the atoms

EGBSA0A Subroutine

"egbsa0a" calculates the generalized Born polarization energy for the GB/SA and ACE solvation models

EGBSA0B Subroutine

"egbsa0b" calculates the generalized Born polarization energy for the GB/SA and ACE solvation models for use with potential smoothing methods via analogy to the smoothing of Coulomb's law

EGBSA1A Subroutine

"egbsa1a" calculates the generalized Born energy and first derivatives with respect to Cartesian coordinates for the GB/SA solvation models

EGBSA1B Subroutine

"egbsa1b" calculates the generalized Born energy and first derivatives with respect to Cartesian coordinates for the GB/SA solvation models

EGBSA2A Subroutine

"egbsa2a" calculates second derivatives of the generalized Born energy term for the GB/SA solvation models

EGBSA2B Subroutine

"egbsa2b" calculates second derivatives of the generalized Born energy term for the GB/SA solvation models via analogy to the smoothing of Coulomb's law

EGBSA3A Subroutine

"egbsa3a" calculates the generalized Born energy term for the GB/SA and ACE solvation models; also partitions the energy among the atoms

EGBSA3B Subroutine

"egbsa3b" calculates the generalized Born polarization energy for the GB/SA and ACE solvation models for use with potential smoothing methods via analogy to the smoothing of Coulomb's law; also partitions the energy among the atoms

EGEOM Subroutine

"egeom" calculates the energy due to restraints on positions, distances, angles and torsions as well as Gaussian basin and spherical droplet restraints

EGEOM1 Subroutine

"egeom1" calculates the energy and first derivatives with respect to Cartesian coordinates due to restraints on positions, distances, angles and torsions as well as Gaussian basin and spherical droplet restraints

EGEOM2 Subroutine

"egeom2" calculates second derivatives of restraints on positions, distances, angles and torsions as well as Gaussian basin and spherical droplet restraints

EGEOM3 Subroutine

"egeom3" calculates the energy due to restraints on positions, distances, angles and torsions as well as Gaussian basin and droplet restraints; also partitions energy among the atoms

EHAL Subroutine

"ehal" calculates the buffered 14-7 van der Waals energy

EHAL0A Subroutine

"ehal0a" calculates the buffered 14-7 van der Waals energy using a pairwise double loop

EHAL0B Subroutine

"ehal0a" calculates the buffered 14-7 van der Waals energy using the method of lights to locate neighboring atoms

EHAL1 Subroutine

"ehal1" calculates the buffered 14-7 van der Waals energy and its first derivatives with respect to Cartesian coordinates

EHAL1A Subroutine

"ehal1a" calculates the buffered 14-7 van der Waals energy and its first derivatives with respect to Cartesian coordinates using a pairwise double loop

EHAL1B Subroutine

"ehal1b" calculates the buffered 14-7 van der Waals energy and its first derivatives with respect to Cartesian coordinates using the method of lights to locate neighboring atoms

EHAL2 Subroutine

"ehal2" calculates the buffered 14-7 van der Waals second derivatives for a single atom at a time

EHAL3 Subroutine

"ehal3" calculates the buffered 14-7 van der Waals energy and partitions the energy among the atoms

EHAL3A Subroutine

"ehal3a" calculates the buffered 14-7 van der Waals energy and partitions the energy among the atoms using a pairwise double loop

EHAL3B Subroutine

"ehal3b" calculates the buffered 14-7 van der Waals energy and also partitions the energy among the atoms using the method of lights to locate neighboring atoms

EIGEN Subroutine

"eigen" uses the power method to compute the largest eigenvalues and eigenvectors of the metric matrix, "valid" is set true if the first three eigenvalues are positive

EIGENRGD Subroutine

EIGENROT Subroutine

EIGENROT Subroutine

EIGENTOR Subroutine

EIGENXYZ Subroutine

EIMPROP Subroutine

"eimprop" calculates the improper dihedral potential energy

EIMPROP1 Subroutine

"eimprop1" calculates improper dihedral energy and its first derivatives with respect to Cartesian coordinates

EIMPROP2 Subroutine

"eimprop2" calculates second derivatives of the improper dihedral angle energy for a single atom

EIMPROP3 Subroutine

"eimprop3" calculates the improper dihedral potential energy; also partitions the energy terms among the atoms

EIMPTOR Subroutine

"eimptor" calculates the improper torsion potential energy

EIMPTOR1 Subroutine

"eimptor1" calculates improper torsion energy and its first derivatives with respect to Cartesian coordinates

EIMPTOR2 Subroutine

"eimptor2" calculates second derivatives of the improper torsion energy for a single atom

EIMPTOR3 Subroutine

"eimptor3" calculates the improper torsion potential energy; also partitions the energy terms among the atoms

ELJ Subroutine

"elj" calculates the Lennard-Jones 6-12 van der Waals energy

ELJOA Subroutine

"elj0a" calculates the Lennard-Jones 6-12 van der Waals energy using a pairwise double loop

ELJ0B Subroutine

"elj0b" calculates the Lennard-Jones 6-12 van der Waals energy using the method of lights to locate neighboring atoms

ELJ1 Subroutine

"elj1" calculates the Lennard-Jones 6-12 van der Waals energy and its first derivatives with respect to Cartesian coordinates

ELJ1A Subroutine

"elj1a" calculates the Lennard-Jones 6-12 van der Waals energy and its first derivatives with respect to Cartesian coordinates using a pairwise double loop

ELJ1B Subroutine

"elj1b" calculates the Lennard-Jones 6-12 van der Waals energy and its first derivatives with respect to Cartesian coordinates using the method of lights to locate neighboring atoms

ELJ2 Subroutine

"elj2" calculates the Lennard-Jones 6-12 van der Waals second derivatives for a single atom at a time

ELJ3 Subroutine

"elj3" calculates the Lennard-Jones 6-12 van der Waals energy and also partitions the energy among the atoms

ELJ3A Subroutine

"elj3a" calculates the Lennard-Jones 6-12 van der Waals energy and also partitions the energy among the atoms using a pairwise double loop

ELJ3B Subroutine

"elj3b" calculates the Lennard-Jones 6-12 van der Waals energy and also partitions the energy among the atoms using the method of lights to locate neighboring atoms

EMBED Subroutine

"embed" is a distance geometry routine patterned after the ideas of Gordon Crippen, Irwin Kuntz and Tim Havel; it takes as input a set of upper and lower bounds on the interpoint distances, chirality restraints and torsional restraints, and attempts to generate a set of coordinates that satisfy the input bounds and restraints

EMETAL Subroutine

"emetal" calculates the transition metal ligand field energy

EMETAL1 Subroutine

"emetal1" calculates the transition metal ligand field energy and its first derivatives with respect to Cartesian coordinates

EMETAL2 Subroutine

"emetal2" calculates the transition metal ligand field second derivatives for a single atom at a time

EMETAL3 Subroutine

"emetal3" calculates the transition metal ligand field energy and also partitions the energy among the atoms

EMM3HB Subroutine

"emm3hb" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy

EMM3HB0A Subroutine

"emm3hb0a" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy using a pairwise double loop

EMM3HB0B Subroutine

"emm3hb0b" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy using the method of lights to locate neighboring atoms

EMM3HB1 Subroutine

"emm3hb1" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy with respect to Cartesian coordinates

EMM3HB1A Subroutine

"emm3hb1a" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy with respect to Cartesian coordinates using a pairwise double loop

EMM3HB1B Subroutine

"emm3hb1b" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy with respect to Cartesian coordinates using the method of lights to locate neighboring atoms

EMM3HB2 Subroutine

"emm3hb2" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding second derivatives for a single atom at a time

EMM3HB3 Subroutine

"emm3hb3" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy, and partitions the energy among the atoms

EMM3HB3A Subroutine

"emm3hb3" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy, and partitions the energy among the atoms

EMM3HB3B Subroutine

"emm3hb3b" calculates the MM3 exp-6 van der Waals and directional charge transfer hydrogen bonding energy using the method of lights to locate neighboring atoms

EMPOLE Subroutine

"empole" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability

EMPOLE0A Subroutine

"empole0a" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability using a pairwise double loop

EMPOLE0B Subroutine

"empole0b" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability using a regular Ewald summation

EMPOLE1 Subroutine

"empole1" calculates the multipole and dipole polarization energy and derivatives with respect to Cartesian coordinates

EMPOLE1A Subroutine

"empole1a" calculates the multipole and dipole polarization energy and derivatives with respect to Cartesian coordinates using a pairwise double loop

EMPOLE1B Subroutine

"empole1b" calculates the multipole and dipole polarization energy and derivatives with respect to Cartesian coordinates using a regular Ewald summation

EMPOLE2 Subroutine

"empole2" calculates second derivatives of the multipole and dipole polarization energy for a single atom at a time

EMPOLE2A Subroutine

"empole2a" computes multipole and dipole polarization first derivatives for a single atom with respect to Cartesian coordinates; used to get finite difference second derivatives

EMPOLE3 Subroutine

"empole3" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability, and partitions the energy among the atoms

EMPOLE3A Subroutine

"empole3a" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability, and partitions the energy among the atoms using a double loop

EMPOLE3B Subroutine

"empole3b" calculates the electrostatic energy due to atomic multipole interactions and dipole polarizability, and partitions the energy among the atoms using a regular Ewald summation

ENERGY Function

"energy" calls the subroutines to calculate the potential energy terms and sums up to form the total energy

ENRGYZE Subroutine

"energyze" is an auxiliary routine for the analyze program that performs the energy analysis and prints the total and intermolecular energies

EOPBEND Subroutine

"eopbend" computes the out-of-plane bend potential energy at trigonal centers via a Wilson-Decius-Cross angle bend

EOPBEND1 Subroutine

"eopbend1" computes the out-of-plane bend potential energy and first derivatives at trigonal centers via a Wilson-Decius-Cross angle bend

EOPBEND2 Subroutine

"eopbend2" calculates second derivatives of the out-of-plane bend energy via a Wilson-Decius-Cross angle bend for a single atom using finite difference methods

EOPBEND2A Subroutine

"eopbend2a" calculates out-of-plane bending first derivatives at a trigonal center via a Wilson-Decius-Cross angle bend; used in computation of finite difference second derivatives

EOPBEND3 Subroutine

"eopbend3" computes the out-of-plane bend potential energy at trigonal centers via a Wilson-Decius-Cross angle bend; also partitions the energy among the atoms

EOPDIST Subroutine

"eopdist" computes the out-of-plane distance potential energy at trigonal centers via the central atom height

EOPDIST1 Subroutine

"eopdist1" computes the out-of-plane distance potential energy and first derivatives at trigonal centers via the central atom height

EOPDIST2 Subroutine

"eopdist2" calculates second derivatives of the out-of-plane distance energy for a single atom via the central atom height

EOPDIST3 Subroutine

"eopdist3" computes the out-of-plane distance potential energy at trigonal centers via the central atom height; also partitions the energy among the atoms

EPME Subroutine

"epme" computes the reciprocal space energy for a particle mesh Ewald summation over partial charges

EPME1 Subroutine

"epme1" computes the reciprocal space energy and first derivatives for a particle mesh Ewald summation

EPME3 Subroutine

"epme3" computes the reciprocal space energy for a particle mesh Ewald summation over partial charges and prints information about the energy over the charge grid points

EPUCLC Subroutine

EREAL Subroutine

"ereal" evaluates the real space portion of the regular Ewald summation energy due to atomic multipole interactions and dipole polarizability

EREAL1 Subroutine

"ereal1" evaluates the real space portion of the regular Ewald summation energy and gradient due to atomic multipole interactions and dipole polarizability

EREAL3 Subroutine

"ereal3" evaluates the real space portion of the regular Ewald summation energy due to atomic multipole interactions and dipole polarizability and partitions the energy among the atoms

ERECIP Subroutine

"erecip" evaluates the reciprocal space portion of the regular Ewald summation energy due to atomic multipole interactions and dipole polarizability

ERECIP1 Subroutine

"erecip1" evaluates the reciprocal space portion of the regular Ewald summation energy and gradient due to atomic multipole interactions and dipole polarizability

ERECIP3 Subroutine

"erecip3" evaluates the reciprocal space portion of the regular Ewald summation energy due to atomic multipole interactions and dipole polarizability, and prints information about the energy over the reciprocal lattice vectors

ERF Function

"erf" computes a numerical approximation to the value of the error function via a Chebyshev approximation

ERFC Function

"erfc" computes a numerical approximation to the value of the complementary error function via a Chebyshev approximation

ERFCORE Subroutine

"erfcore" evaluates erf(x) or erfc(x) for a real argument x; when called with mode set to 0 it returns erf, a mode of 1 returns erfc; uses rational functions that approximate erf(x) and erfc(x) to at least 18 significant decimal digits

ERFIK Subroutine

ERFINV Function

"erfinv" evaluates the inverse of the error function erf for a real argument in the range (-1,1) using a rational function approximation followed by cycles of Newton-Raphson correction

ERXNFLD Subroutine

"erxnfld" calculates the macroscopic reaction field energy

ERXNFLD1 Subroutine

"erxnfld1" calculates the macroscopic reaction field energy and derivatives with respect to Cartesian coordinates

ERXNFLD2 Subroutine

"erxnfld2" calculates second derivatives of the macroscopic reaction field energy for a single atom at a time

ERXNFLD3 Subroutine

"erxnfld3" calculates the macroscopic reaction field energy, and also partitions the energy among the atoms

ESOLV Subroutine

"esolv" calculates the continuum solvation energy via either the Eisenberg-McLachlan ASP model, Ooi-Scheraga SASA model, various GB/SA methods or the ACE model

ESOLV1 Subroutine

"esolv1" calculates the continuum solvation energy and first derivatives with respect to Cartesian coordinates using either the Eisenberg-McLachlan ASP, Ooi-Scheraga SASA or various GB/SA solvation models

ESOLV2 Subroutine

"esolv2" calculates second derivatives of the continuum solvation energy using either the Eisenberg-McLachlan ASP, Ooi-Scheraga SASA or various GB/SA solvation models

ESOLV3 Subroutine

"esolv3" calculates the continuum solvation energy using either the Eisenberg-McLachlan ASP model, Ooi-Scheraga SASA model, various GB/SA methods or the ACE model; also partitions the energy among the atoms

ESTRBND Subroutine

"estrbnd" calculates the stretch-bend potential energy

ESTRBND1 Subroutine

"estrbnd1" calculates the stretch-bend potential energy and first derivatives with respect to Cartesian coordinates

ESTRBND2 Subroutine

"estrbnd2" calculates the stretch-bend potential energy second derivatives with respect to Cartesian coordinates

ESTRBND3 Subroutine

"estrbnd3" calculates the stretch-bend potential energy; also partitions the energy among the atoms

ESTRTOR Subroutine

"estrtor" calculates the stretch-torsion potential energy

ESTRTOR1 Subroutine

"estrtor1" calculates the stretch-torsion energy and first derivatives with respect to Cartesian coordinates

ESTRTOR2 Subroutine

"estrtor2" calculates the stretch-torsion potential energy second derivatives with respect to Cartesian coordinates

ESTRTOR3 Subroutine

"estrtor3" calculates the stretch-torsion potential energy; also partitions the energy terms among the atoms

ETORS Subroutine

"etors" calculates the torsional potential energy

ETORS1 Subroutine

"etors1" calculates torsional potential energy and first derivatives with respect to Cartesian coordinates

ETORS2 Subroutine

"etors2" calculates second derivatives of the torsional energy for a single atom

ETORS3 Subroutine

"etors3" calculates the torsional potential energy; also partitions the energy terms among the atoms

EUREY Subroutine

"eurey" calculates the Urey-Bradley 1-3 interaction energy

EUREY1 Subroutine

"eurey1" calculates the Urey-Bradley interaction energy and its first derivatives with respect to Cartesian coordinates

EUREY2 Subroutine

"eurey2" calculates second derivatives of the Urey-Bradley interaction energy for a single atom at a time

EUREY3 Subroutine

"eurey3" calculates the Urey-Bradley energy; also partitions the energy among the atoms

EWALDCOF Subroutine

"ewaldcof" finds a value of the Ewald coefficient such that all terms beyond the specified cutoff distance will have an value less than a specified tolerance

EXPLORE Subroutine

"explore" uses simulated annealing on an initial crude embedded distance geometry structure to refine versus the bound, chirality, planarity and torsional error functions

EXTRA Subroutine

"extra" calculates any additional user defined potential energy contribution

EXTRA1 Subroutine

"extral" calculates any additional user defined potential energy contribution and its first derivatives

EXTRA2 Subroutine

"extra2" calculates second derivatives of any additional user defined potential energy contribution for a single atom at a time

EXTRA3 Subroutine

"extra3" calculates any additional user defined potential contribution and also partitions the energy among the atoms

FATAL Subroutine

"fatal" terminates execution due to a user request, a severe error or some other nonstandard condition

FFTBACK Subroutine

FFTFRONT Subroutine

FFTSETUP Subroutine

FIELD Subroutine

"field" sets the force field potential energy functions from a parameter file and modifications specified in a keyfile

FINAL Subroutine

"final" performs any final program actions, prints a status message, and then pauses if necessary to avoid closing the execution window

FINDATM Subroutine

"findatm" locates a specific PDB atom name type within a range of atoms from the PDB file, returns zero if the name type was not found

FIXPDB Subroutine

"fixpdb" corrects problems with PDB files by converting residue and atom names to the forms used by TINKER

FRACDIST Subroutine

"fracdist" computes a normalized distribution of the pairwise fractional distances between the smoothed upper and lower bounds

FREEUNIT Function

"freeunit" finds an unopened Fortran I/O unit and returns its numerical value from 1 to 99; the units already assigned to "input" and "iout" (usually 5 and 6) are skipped since they have special meaning as the default I/O units

GAMMLN Function

"gammln" uses a series expansion due to Lanczos to compute the natural logarithm of the Gamma function at "x" in [0,1]

GDA Program

"gda" implements Gaussian Density Annealing (GDA) algorithm for global optimization via simulated annealing

GDA1 Function

GDA2 Function

GDA3 Subroutine

GDASTAT Subroutine

GENDOT Subroutine

"gendot" finds the coordinates of a specified number of surface points for a sphere with the input radius and coordinate center

GEODESIC Subroutine

"geodesic" smooths the upper and lower distance bounds via the triangle inequality using a sparse matrix version of a shortest path algorithm

GEOMETRY Function

"geometry" finds the value of the interatomic distance, angle or dihedral angle defined by two to four input atoms

GETBASE Subroutine

"getbase" finds the base heavy atoms for a single nucleotide residue and copies the names and coordinates to the Protein Data Bank file

GETIME Subroutine

"getime" gets elapsed CPU time in seconds for an interval

GETINT Subroutine

"getint" asks for an internal coordinate file name, then reads the internal coordinates and computes Cartesian coordinates

GETKEY Subroutine

"getkey" finds a valid keyfile and stores its contents as line images for subsequent keyword parameter searching

GETMOL2 Subroutine

"getmol2" asks for a Sybyl MOL2 molecule file name, then reads the coordinates from the file

GETNUCH Subroutine

"getnuch" finds the nucleotide hydrogen atoms for a single residue and copies the names and coordinates to the Protein Data Bank file

GETNUMB Subroutine

"getnumb" searches an input string from left to right for an integer and puts the numeric value in "number"; returns zero with "next" unchanged if no integer value is found

GETPDB Subroutine

"getpdb" asks for a Protein Data Bank file name, then reads in the coordinates file

GETPRB Subroutine

"getprb" tests for a possible probe position at the interface between three neighboring atoms

GETPRM Subroutine

"getprm" finds the potential energy parameter file and then opens and reads the parameters

GETPROH Subroutine

"getproh" finds the hydrogen atoms for a single amino acid residue and copies the names and coordinates to the Protein Data Bank file

GETREF Subroutine

"getref" copies structure information from the reference area into the standard variables for the current system structure

GETSEQ Subroutine

"getseq" asks the user for the amino acid sequence and torsional angle values needed to define a peptide

GETSEQN Subroutine

"getseqn" asks the user for the nucleotide sequence and torsional angle values needed to define a nucleic acid

GETSIDE Subroutine

"getside" finds the side chain heavy atoms for a single amino acid residue and copies the names and coordinates to the Protein Data Bank file

GETSTRING Subroutine

"getstring" searches for a quoted text string within an input character string; the region between the first and second quotes is returned as the "text"; if the actual text is too long, only the first part is returned

GETTEXT Subroutine

"gettext" searches an input string for the first string of non-blank characters; the region from a non-blank character to the first blank space is returned as "text"; if the actual text is too long, only the first part is returned

GETTOR Subroutine

"gettor" tests for a possible torus position at the interface between two atoms, and finds the torus radius, center and axis

GETWORD Subroutine

"getword" searches an input string for the first alphabetic character (A-Z or a-z); the region from this first character to the first blank space or comma is returned as a "word"; if the actual word is too long, only the first part is returned

GETXYZ Subroutine

"getxyz" asks for a Cartesian coordinate file name, then reads in the coordinates file

GRADIENT Subroutine

"gradient" calls subroutines to calculate the potential energy and first derivatives with respect to Cartesian coordinates

GRADRGD Subroutine

"gradrgd" calls subroutines to calculate the potential energy and first derivatives with respect to rigid body coordinates

GRADROT Subroutine

"gradrot" calls subroutines to calculate the potential energy and its torsional first derivatives

GRAFIC Subroutine

"grafic" outputs the upper & lower triangles and diagonal of a square matrix in a schematic form for visual inspection

GROUPS Subroutine

"groups" tests a set of atoms to see if all are members of a single atom group or a pair of atom groups; if so, then the correct intra- or intergroup weight is assigned

GRPLINE Subroutine

"grpline" tests each atom group for linearity of the sites contained in the group

GYRATE Subroutine

"gyrate" computes the radius of gyration of a molecular system from its atomic coordinates

HANGLE Subroutine

"hangle" constructs hybrid angle bending parameters given an initial state, final state and "lambda" value

HATOM Subroutine

"hatom" assigns a new atom type to each hybrid site

HBOND Subroutine

"hbond" constructs hybrid bond stretch parameters given an initial state, final state and "lambda" value

HCHARGE Subroutine

"hcharge" constructs hybrid charge interaction parameters given an initial state, final state and "lambda" value

HDIPOLE Subroutine

"hdipole" constructs hybrid dipole interaction parameters given an initial state, final state and "lambda" value

HESSIAN Subroutine

"hessian" calls subroutines to calculate the Hessian elements for each atom in turn with respect to Cartesian coordinates

HESSRGD Subroutine

"hessrgd" computes the numerical Hessian elements with respect to rigid body coordinates; either the full matrix or just the diagonal can be calculated; the full matrix needs $6 \times n_{\text{group}} + 1$ gradient evaluations while the diagonal requires just two gradient calls

HESSROT Subroutine

"hessrot" computes the numerical Hessian elements with respect to torsional angles; either the full matrix or just the diagonal can be calculated; the full matrix needs $n_{\text{omega}} + 1$ gradient evaluations while the diagonal requires just two gradient calls

HIMPTOR Subroutine

"himptor" constructs hybrid improper torsional parameters given an initial state, final state and "lambda" value

HSTRBND Subroutine

"hstrbnd" constructs hybrid stretch-bend parameters given an initial state, final state and "lambda" value

HSTRTOR Subroutine

"hstrtor" constructs hybrid stretch-torsion parameters given an initial state, final state and "lambda" value

HTORS Subroutine

"htors" constructs hybrid torsional parameters for a given initial state, final state and "lambda" value

HVDW Subroutine

"hvdw" constructs hybrid van der Waals parameters given an initial state, final state and "lambda" value

HYBRID Subroutine

"hybrid" constructs the hybrid hamiltonian for a specified initial state, final state and mutation parameter "lambda"

IJK_PT Subroutine

IMAGE Subroutine

"image" takes the components of pairwise distance between two points in the same or neighboring periodic boxes and converts to the components of the minimum image distance

IMPOSE Subroutine

"impose" performs the least squares best superposition of two atomic coordinate sets via a quaternion method; upon return, the first coordinate set is unchanged while the second set is translated and rotated to give best fit; the final root mean square fit is returned in "rmsvalue"

INDUCE Subroutine

"induce" computes the induced dipole moment at each polarizable site due to direct or mutual polarization

INDUCE0A Subroutine

"induce0a" computes the induced dipole moment at each polarizable site using a pairwise double loop

INDUCE0B Subroutine

"induce0b" computes the induced dipole moment at each polarizable site using a regular Ewald summation

INEDGE Subroutine

"inedge" inserts a concave edge into the linked list for its temporary torus

INERTIA Subroutine

"inertia" computes the principal moments of inertia for the system, and optionally translates the center of mass to the origin and rotates the principal axes onto the global axes

INITERR Function

"initerr" is the initial error function and derivatives for a distance geometry embedding; it includes components from the local geometry and torsional restraint errors

INITIAL Subroutine

"initial" sets up original values for some parameters and variables that might not otherwise get initialized

INITPRM Subroutine

"initprm" completely initializes a force field by setting all parameters to zero and using defaults for control values

INITRES Subroutine

"initres" sets names for biopolymer residue types used in PDB file conversion and automated generation of structures

INITROT Subroutine

"initrot" asks for torsional angles which are to be rotated in subsequent computation, it will automatically locate all rotatable single bonds if desired; assumes that an appropriate internal coordinates file has already been read in

INSERT Subroutine

"insert" adds the specified atom to the Cartesian coordinates list and shifts the remaining atoms

INTEDIT Program

"intedit" allows the user to extract information from or alter the values within an internal coordinates file

INTXYZ Program

"intxyz" takes as input an internal coordinates file, converts to and then writes out Cartesian coordinates

INVBETA Function

"invbeta" computes the inverse Beta distribution function via a combination of Newton iteration and bisection search

INVERT Subroutine

"invert" inverts a matrix using the Gauss-Jordan method

IPEDGE Subroutine

"ipedge" inserts convex edge into linked list for atom

JACOBI Subroutine

"jacobi" performs a matrix diagonalization of a real symmetric matrix by the method of Jacobi rotations

KANGANG Subroutine

"kangang" assigns the parameters for angle-angle cross term interactions and processes new or changed parameter values

KANGLE Subroutine

"kangle" assigns the force constants and ideal angles for the bond angles; also processes new or changed parameters

KATOM Subroutine

"katom" assigns an atom type definitions to each atom in the structure and processes any new or changed values

KBOND Subroutine

"kbond" assigns a force constant and ideal bond length to each bond in the structure and processes any new or changed parameter values

KCHARGE Subroutine

"kcharge" assigns partial charges to the atoms within the structure and processes any new or changed values

KCHIRAL Subroutine

"kchiral" determines the target value for each chirality and planarity restraint as the signed volume of the parallelepiped spanned by vectors from a common atom to each of three other atoms

KDIPOLE Subroutine

"kdipole" assigns bond dipoles to the bonds within the structure and processes any new or changed values

KENEG Subroutine

"keneg" applies primary and secondary electronegativity bond length corrections to applicable bond parameters

KEWALD Subroutine

"kewald" assigns both regular Ewald summation and particle mesh Ewald parameters for a periodic box

KGEOM Subroutine

"kgeom" assigns parameters for geometric restraint terms to be included in the potential energy calculation

KIMPROP Subroutine

"kimprop" assigns potential parameters to each improper dihedral in the structure and processes any changed values

KIMPTOR Subroutine

"kimptor" assigns torsional parameters to each improper torsion in the structure and processes any changed values

KMETAL Subroutine

"kmetal" assigns ligand field parameters to transition metal atoms and processes any new or changed parameter values

KMPOLE Subroutine

"kmpole" assigns atomic multipole moments to the atoms of the structure and processes any new or changed values

KOPBEND Subroutine

"kopbend" assigns the force constants for out-of-plane bending at trigonal centers via Wilson-Decius-Cross angle bends; also processes any new or changed parameter values

KOPDIST Subroutine

"kopdist" assigns the force constants for out-of-plane distance at trigonal centers via the central atom height; also processes any new or changed parameter values

KORBIT Subroutine

"korbit" assigns pi-orbital parameters to conjugated systems and processes any new or changed parameters

KPOLAR Subroutine

"kpolar" assigns atomic dipole polarizabilities to the atoms within the structure and processes any new or changed values

KSOLV Subroutine

"ksolv" assigns continuum solvation energy parameters for the Eisenberg-McLachlan ASP, Ooi-Scheraga SASA or various GB/SA solvation models

KSTRBND Subroutine

"kstrbnd" assigns the parameters for the stretch-bend interactions and processes new or changed parameter values

KSTRTOR Subroutine

"kstrtor" assigns stretch-torsion parameters to torsions needing them, and processes any new or changed values

KTORS Subroutine

"ktors" assigns torsional parameters to each torsion in the structure and processes any new or changed values

KUREY Subroutine

"kurey" assigns the force constants and ideal distances for the Urey-Bradley 1-3 interactions; also processes any new or changed parameter values

KVDW Subroutine

"kvdw" assigns the parameters to be used in computing the van der Waals interactions and processes any new or changed values for these parameters

LATTICE Subroutine

"lattice" stores the periodic box dimensions and sets angle values to be used in computing fractional coordinates

LBFGS Subroutine

"lbfgs" is a limited memory BFGS quasi-newton nonlinear optimization routine

LIGASE Subroutine

"ligase" translates a nucleic acid structure in Protein Data Bank format to a Cartesian coordinate file and sequence file

LIGHTS Subroutine

"lights" computes the set of nearest neighbor interactions using the method of lights algorithm

LINBODY Subroutine

"linbody" finds the angular velocity of a linear rigid body

LMSTEP Subroutine

"lmstep" computes the Levenberg-Marquardt step during a nonlinear least squares calculation; this version is based upon ideas from the Minpack routine LMPAR together with the internal doubling strategy of Dennis and Schnabel

LOCALMIN Subroutine

"localmin" is used during normal mode local search to perform a Cartesian coordinate energy minimization

LOCALRGD Subroutine

"localrgd" is used during the PSS local search procedure to perform a rigid body energy minimization

LOCALROT Subroutine

"localrot" is used during the PSS local search procedure to perform a torsional space energy minimization

LOCALXYZ Subroutine

"localxyz" is used during the potential smoothing and search procedure to perform a local optimization at the current smoothing level

LOCERR Function

"locerr" is the local geometry error function and derivatives including the 1-2, 1-3 and 1-4 distance bound restraints

LOWCASE Subroutine

"lowercase" converts a text string to all lower case letters

MAJORIZE Subroutine

"majorize" refines the projected coordinates by attempting to minimize the least square residual between the trial distance matrix and the distances computed from the coordinates

MAKEINT Subroutine

"makeint" converts Cartesian to internal coordinates where selection of internal coordinates is controlled by "mode"

MAKEPDB Subroutine

"makexyz" converts a set of Cartesian coordinates to Protein Data Bank format with special handling for systems consisting of polypeptide chains, ligands and water molecules

MAKEREf Subroutine

"makeref" copies the information contained in the "xyz" file of the current structure into corresponding reference areas

MAKEXYZ Subroutine

"makexyz" generates a complete set of Cartesian coordinates for a full structure from the internal coordinate values

MAPCHECK Subroutine

"mapcheck" checks the current minimum energy structure for possible addition to the master list of local minima

MAXWELL Function

"maxwell" returns a speed in Angstroms/picosecond randomly selected from a 3-D Maxwell-Boltzmann distribution for the specified particle mass and system temperature

MDINIT Subroutine

"mdinit" initializes the velocities and accelerations for a molecular dynamics trajectory, including restarts

MDREST Subroutine

"mdrest" finds and removes any translational or rotational kinetic energy of the overall system center of mass

MDSAVE Subroutine

"mdsave" writes molecular dynamics trajectory snapshots and auxiliary files with velocity and induced dipole information; also checks for user requested termination of a simulation

MDSTAT Subroutine

"mdstat" is called at each molecular dynamics time step to form statistics on various average values and fluctuations, and to periodically save the state of the trajectory

MEASFN Subroutine

MEASFP Subroutine

MEASFS Subroutine

MEASPM Subroutine

"measpm" computes the volume of a single prism section of the full interior polyhedron

MECHANIC Subroutine

"mechanic" sets up needed parameters for the potential energy calculation and reads in many of the user selectable options

MERGE Subroutine

"merge" combines the reference and current structures into a single new "current" structure containing the reference atoms followed by the atoms of the current structure

METRIC Subroutine

"metric" takes as input the trial distance matrix and computes the metric matrix of all possible dot products between the atomic vectors and the center of mass using the law of cosines and the following formula for the distances to the center of mass:

MIDERR Function

"miderr" is the secondary error function and derivatives for a distance geometry embedding; it includes components from the distance bounds, local geometry, chirality and torsional restraint errors

MINIMIZE1 Function

"minimize1" is a service routine that computes the energy and gradient for a low storage BFGS optimization in Cartesian coordinate space

MINIMIZE Program

"minimize" performs energy minimization in Cartesian coordinate space using a low storage BFGS nonlinear optimization

MINIROT Program

"minirot" performs an energy minimization in torsional angle space using a low storage BFGS nonlinear optimization

MINIROT1 Function

"minirot1" is a service routine that computes the energy and gradient for a low storage BFGS nonlinear optimization in torsional angle space

MINPATH Subroutine

"minpath" is a routine for finding the triangle smoothed upper and lower bounds of each atom to a specified root atom using a sparse variant of the Bellman-Ford shortest path algorithm

MINRIGID Program

"minrigid" performs an energy minimization of rigid body atom groups using a low storage BFGS nonlinear optimization

MINRIGID1 Function

"minrigid1" is a service routine that computes the energy and gradient for a low storage BFGS nonlinear optimization of rigid bodies

MMID Subroutine

"mmid" implements a modified midpoint method to advance the integration of a set of first order differential equations

MODECART Subroutine

MODEROT Subroutine

MODESRCH Subroutine

MODETORS Subroutine

MODULI Subroutine

"moduli" sets the moduli of the inverse discrete Fourier transform of the B-splines; bsmod[1-3] hold these values, nfft[1-3] are the grid dimensions, bsorder is the order of B-spline approximation

MOLECULE Subroutine

"molecule" counts the molecules, assigns each atom to its molecule and computes the mass of each molecule

MOLUIND Subroutine

"moluind" computes the molecular induced dipole components in the presence of an external electric field

MOMENTS Subroutine

"moments" computes the total electric charge, dipole and quadrupole moments for the entire system as a sum over the partial charges, bond dipoles and atomic multipole moments

MUTATE Subroutine

"mutate" constructs the hybrid hamiltonian for a specified initial state, final state and mutation parameter "lambda"

NEIGHBOR Subroutine

"neighbor" finds all of the neighbors of each atom

NEWATM Subroutine

"newatm" creates and defines an atom needed for the Cartesian coordinates file, but which may not present in the original Protein Data Bank file

NEWCRD Subroutine

"newcrd" computes updated atomic coordinates for a rigid body given the previous coordinates, the rotation matrix and shift in the center of mass corresponding to the motion

NEWTON Program

"newton" performs an energy minimization in Cartesian coordinate space using a truncated Newton method

NEWTON1 Function

"newton1" is a service routine that computes the energy and gradient for truncated Newton optimization in Cartesian coordinate space

NEWTON2 Subroutine

"newton2" is a service routine that computes the sparse matrix Hessian elements for truncated Newton optimization in Cartesian coordinate space

NEWTROT Program

"newtrot" performs an energy minimization in torsional angle space using a truncated Newton conjugate gradient method

NEWTROT1 Function

"newtrot1" is a service routine that computes the energy and gradient for truncated Newton conjugate gradient optimization in torsional angle space

NEWTROT2 Subroutine

"newtrot2" is a service routine that computes the sparse matrix Hessian elements for truncated Newton optimization in torsional angle space

NEXTARG Subroutine

"nextarg" finds the next unused command line argument and returns it in the input character string

NEXTTEXT Function

"nexttext" finds and returns the location of the first non-blank character within an input text string; zero is returned if no such character is found

NORMAL Function

"normal" generates a random number from a normal Gaussian distribution with a mean of zero and a variance of one

NUCBASE Subroutine

"nucbase" builds the side chain for a single nucleotide base in terms of internal coordinates

NUCCHAIN Subroutine

"nucchain" builds up the internal coordinates for a nucleic acid sequence from the sugar type, backbone and glycosidic torsional values

NUCLEIC Program

"nucleic" builds the internal and Cartesian coordinates of a polynucleotide from nucleic acid sequence and torsional angle values for the nucleic acid backbone and side chains

NUMBER Function

"number" converts a text numeral into an integer value; the input string must contain only numeric characters

NUMERAL Subroutine

"numeral" converts an input integer number into the corresponding right- or left-justified text numeral

NUMGRAD Subroutine

"numgrad" computes the gradient of the objective function "fvalue" with respect to Cartesian coordinates of the atoms via a two-sided numerical differentiation

OCVM Subroutine

"ocvm" is an optimally conditioned variable metric nonlinear optimization routine without line searches

OLDATM Subroutine

"oldatm" get the Cartesian coordinates for an atom from the Protein Data Bank file, then assigns the atom type and atomic connectivities

OPENEND Subroutine

"openend" opens a file on a Fortran unit such that the position is set to the bottom for appending to the end of the file

OPTIMIZ1 Function

"optimiz1" is a service routine that computes the energy and gradient for optimally conditioned variable metric optimization in Cartesian coordinate space

OPTIMIZE Program

"optimize" performs energy minimization in Cartesian coordinate space using an optimally conditioned variable metric method

OPTIROT Program

"optirot" performs an energy minimization in torsional angle space using an optimally conditioned variable metric method

OPTIROT1 Function

"optirot1" is a service routine that computes the energy and gradient for optimally conditioned variable metric optimization in torsional angle space

OPTRIGID Program

"optrigid" performs an energy minimization of rigid body atom groups using an optimally conditioned variable metric method

OPTRIGID1 Function

"optrigid1" is a service routine that computes the energy and gradient for optimally conditioned variable metric optimization of rigid bodies

ORBITAL Subroutine

"orbital" finds and organizes lists of atoms in a pisystem, bonds connecting pisystem atoms and torsions whose two central atoms are both pisystem atoms

ORIENT Subroutine

"orient" computes a set of reference Cartesian coordinates in standard orientation for each rigid body atom group

ORTHOG Subroutine

"orthog" performs an orthogonalization of an input matrix via the modified Gram-Schmidt algorithm

OVERLAP Subroutine

"overlap" computes the overlap for two parallel p-orbitals given the atomic numbers and distance of separation

PARAMYZE Subroutine

"paramyze" prints the force field parameters used in the computation of each of the potential energy terms

PASSB Subroutine

PASSB2 Subroutine

PASSB3 Subroutine

PASSB4 Subroutine

PASSB5 Subroutine

PASSF Subroutine

PASSF2 Subroutine

PASSF3 Subroutine

PASSF4 Subroutine

PASSF5 Subroutine

PATH Program

"path" locates a series of structures equally spaced along a conformational pathway connecting the input reactant and product structures; a series of constrained optimizations orthogonal to the path is done via Lagrangian multipliers

PATH1 Function

PATHPNT Subroutine

"pathpnt" finds a structure on the synchronous transit path with the specified path value "t"

PATHSCAN Subroutine

"pathscan" makes a scan of a synchronous transit pathway by computing structures and energies for specific path values

PATHVAL Subroutine

"pathval" computes the synchronous transit path value for the specified structure

PDBATM Subroutine

"pdbatm" adds an atom to the Protein Data Bank file

PDBXYZ Program

"pdbxyz" takes as input a Protein Data Bank file and then converts to and writes out a Cartesian coordinates file and, for polypeptides, a sequence file

PIALTER Subroutine

"pialter" first modifies bond lengths and force constants according to the standard bond slope parameters and the bond order values stored in "pnpl"; also alters some 2-fold torsional parameters based on the bond-order * beta matrix

PIMOVE Subroutine

"pimove" rotates the vector between atoms "list(1)" and "list(2)" so that atom 1 is at the origin and atom 2 along the x-axis; the atoms defining the respective planes are also moved and their bond lengths normalized

PIPLANE Subroutine

"piplane" selects the three atoms which specify the plane perpendicular to each p-orbital; the current version will fail in certain situations, including ketenes, allenes, and isolated or adjacent triple bonds

PISCF Subroutine

"piscf" performs an scf molecular orbital calculation for the pisystem using a modified Pariser-Parr-Pople method

PITILT Subroutine

"pitilt" calculates for each pibond the ratio of the actual p-orbital overlap integral to the ideal overlap if the same orbitals were perfectly parallel

PLACE Subroutine

"place" finds the probe sites by putting the probe sphere tangent to each triple of neighboring atoms

POLARGRP Subroutine

"polargrp" generates members of the polarization group of each atom and separate lists of the 1-2, 1-3 and 1-4 group connectivities

POLARIZE Program

"polarize" computes the molecular polarizability by applying an external field along each axis followed by diagonalization of the resulting polarizability tensor

POLYMER Subroutine

"polymer" tests for the presence of an infinite polymer extending across periodic boundaries

POLYP Subroutine

POTNRG Function

POTOFF Subroutine

"potoff" clears the forcefield definition by turning off the use of each of the potential energy functions

POWER Subroutine

"power" uses the power method with deflation to compute the few largest eigenvalues and eigenvectors of a symmetric matrix

PRECISE Function

"precise" finds a machine precision value as selected by the input argument: (1) the smallest positive floating point value, (2) the smallest relative floating point spacing, (3) the largest relative floating point spacing

PRECOND Subroutine

"precond" solves a simplified version of the Newton equations $M\mathbf{s} = \mathbf{r}$, and uses the result to precondition linear conjugate gradient iterations on the full Newton equations in "tnsolve"

PRESSURE Subroutine

"pressure" uses the internal virial to find the pressure in a periodic box and maintains a constant desired pressure by scaling the coordinates via coupling to an external constant pressure bath

PRMKEY Subroutine

"field" parses a text string to extract keywords related to force field potential energy functional forms and constants

PROCHAIN Subroutine

"prochain" builds up the internal coordinates for an amino acid sequence from the phi, psi, omega and chi values

PROJECT Subroutine

PROMO Subroutine

"promo" writes a short message containing information about the TINKER version number and the copyright notice

PROPERTY Function

"property" takes two input snapshot frames and computes the value of the property for which the correlation function is being accumulated

PROPYZE Subroutine

"propyze" finds and prints the total charge, dipole moment components, radius of gyration and moments of inertia

PROSIDE Subroutine

"proside" builds the side chain for a single amino acid residue in terms of internal coordinates

PROTEIN Program

"protein" builds the internal and Cartesian coordinates of a polypeptide from amino acid sequence and torsional angle values for the peptide backbone and side chains

PRTDYN Subroutine

"prtdyn" writes out the information needed to restart a molecular dynamics trajectory to an external disk file

PRTERR Subroutine

"prtterr" writes out a set of coordinates to a disk file prior to aborting on a serious error

PRTINT Subroutine

"prtint" writes out a set of Z-matrix internal coordinates to an external disk file

PRTMOL2 Program

"prtmol2" writes out a set of coordinates in Sybyl MOL2 format to an external disk file

PRTMSI Subroutine

"prtmsi" writes out a set of Cartesian coordinates for all active atoms in the MSI Insight II archive format

PRTPDB Subroutine

"prtpdb" writes out a set of Protein Data Bank coordinates to an external disk file

PRTPRM Subroutine

"prtprm" writes out a formatted listing of the default set of potential energy parameters for a force field

PRTSEQ Subroutine

"prtseq" writes out a biopolymer sequence to an external disk file with 15 residues per line and distinct chains separated by blank lines

PRTXMOL Subroutine

"prtxmol" writes out a set of Cartesian coordinates for all active atoms in a simple, generic XYZ format originally due to the XMOL program

PRTXYZ Subroutine

"prtxyz" writes out a set of Cartesian coordinates to an external disk file

PSS Program

"pss" implements the potential smoothing plus search method for global optimization in Cartesian coordinate space with local searches performed in Cartesian or torsional space

PSS1 Function

"pss1" is a service routine that computes the energy and gradient during PSS global optimization in Cartesian coordinate space

PSS2 Subroutine

"pss2" is a service routine that computes the sparse matrix Hessian elements during PSS global optimization in Cartesian coordinate space

PSSRGD1 Function

"pssrgd1" is a service routine that computes the energy and gradient during PSS global optimization over rigid bodies

PSSRIGID Program

"pssrigid" implements the potential smoothing plus search method for global optimization for a set of rigid bodies

PSSROT Program

"pssrot" implements the potential smoothing plus search method for global optimization in torsional space

PSSROT1 Function

"pssrot1" is a service routine that computes the energy and gradient during PSS global optimization in torsional space

PSSWRITE Subroutine

PTINCY Function

PZEXTR Subroutine

QRFACT Subroutine

"qract" performs Householder transformations with column pivoting (optional) to compute a QR factorization of the m by n matrix a; the routine determines an orthogonal matrix q, a permutation matrix p, and an upper trapezoidal matrix r with diagonal elements of nonincreasing magnitude, such that $a \cdot p = q \cdot r$; the Householder transformation for column k, $k = 1, 2, \dots, \min(m, n)$, is of the form

QRSOLVE Subroutine

"qrsolve" solves $a \cdot x = b$ and $d \cdot x = 0$ in the least squares sense; normally used in combination with routine "qract" to solve least squares problems

QUATFIT Subroutine

"quatfit" uses a quaternion-based method to achieve the best fit superposition of two sets of coordinates

RADIAL Program

"radial" finds the radial distribution function for a specified pair of atom types via analysis of a set of stored coordinate frames from a liquid simulation

RANDOM Function

"random" generates a random number on [0,1] via a long period generator due to L'Ecuyer with Bays-Durham shuffle

RANVEC Subroutine

"ranvec" generates a unit vector in 3-dimensional space with uniformly distributed random orientation

RATTLE Subroutine

"rattle" implements the first portion of the rattle algorithm by correcting atomic positions and half-step velocities to maintain constrained interatomic distances

RATTLE2 Subroutine

"rattle2" implements the second portion of the rattle algorithm by correcting the full-step velocities in order to maintain constrained interatomic distances

READBLK Subroutine

"readblk" reads in a set of snapshot frames and transfers the values to internal arrays for use in the computation of time correlation functions

READDYN Subroutine

"readdyn" get the positions, velocities and accelerations for a molecular dynamics restart from an external disk file

READINT Subroutine

"readint" gets a set of Z-matrix internal coordinates from an external file

READMOL2 Subroutine

"readmol2" gets a set of Sybyl MOL2 coordinates from an external disk file

READPDB Subroutine

"readpdb" gets a set of Protein Data Bank coordinates from an external disk file

READPRM Subroutine

"readprm" processes the potential energy parameter file in order to define the default force field parameters

READSEQ Subroutine

"readseq" gets a biopolymer sequence containing one or more separate chains from an external file; all lines containing sequence must begin with the starting sequence number, the actual sequence is read from subsequent nonblank characters

READXYZ Subroutine

"readxyz" gets a set of Cartesian coordinates from an external disk file

REFINE Subroutine

"refine" performs minimization of the atomic coordinates of an initial crude embedded distance geometry structure versus the bound, chirality, planarity and torsional error functions

REGBODY Subroutine

"regbody" finds the angular velocity of a regular, nonlinear rigid body

REPLICA Subroutine

"replica" decides between images and replicates for generation of periodic boundary conditions, and sets the cell replicate list if the replicates method is to be used

RFINDEX Subroutine

RGDSRCH Subroutine

RGDSTEP Subroutine

"rgdstep" performs a single molecular dynamics time step for a rigid-body calculation

RIBOSOME Subroutine

"ribosome" translates a polypeptide structure in Protein Data Bank format to a Cartesian coordinate file and sequence file

RIGIDXYZ Subroutine

"rigidxyz" computes Cartesian coordinates for a rigid body group via rotation and translation of reference coordinates

RINGS Subroutine

"rings" searches the structure for small rings and stores their component atoms; code to remove the reducible rings consisting of smaller rings is commented in this version since reducible rings are needed for parameter assignment

RMSERROR Subroutine

"rmserror" computes the maximum absolute deviation and the rms deviation from the distance bounds, and the number and rms value of the distance restraint violations

RMSFIT Function

"rmsfit" computes the rms fit of two coordinate sets

ROTANG Function

ROTCHECK Function

"rotcheck" tests a specified candidate rotatable bond for the disallowed case where inactive atoms are found on both sides of the candidate bond

ROTEULER Subroutine

"roteuler" computes a set of Euler angle values consistent with an input rotation matrix

ROTLIST Subroutine

"rotlist" generates the minimum list of all the atoms lying to one side of a pair of directly bonded atoms; optionally finds the minimal list by choosing the side with fewer atoms

ROTMAT Subroutine

"rotmat" find the rotation matrix that converts from the local coordinate system at each multipole site to the global system

ROTPOLE Subroutine

"rotpole" computes the atomic multipole values in the global coordinate frame by applying a rotation matrix to a set of locally defined multipoles

ROTRGD Subroutine

"rotrgd" finds the rotation matrix for a rigid body due to a single step of dynamics

SADDLE Program

"saddle" finds a transition state between two conformational minima using a combination of ideas from the synchronous transit (Halgren-Lipscomb) and quadratic path (Bell-Crighton) methods

SADDLE1 Function

SADDLES Subroutine

"saddles" constructs circles, convex edges and saddle faces

SCAN Program

"scan" attempts to find all the local minima on a potential energy surface via an iterative series of local searches

SCAN1 Function

"scan1" is a service routine that computes the energy and gradient during exploration of a potential energy surface via iterative local search

SCAN2 Subroutine

"scan2" is a service routine that computes the sparse matrix Hessian elements during exploration of a potential energy surface via iterative local search

SDAREA Subroutine

"sdarea" optionally scales the atomic friction coefficient of each atom based on its accessible surface area

SDSTEP Subroutine

"sdstep" performs a single stochastic dynamics time step via a velocity Verlet integration algorithm

SDTERM Subroutine

"sdterm" gets frictional and random force terms needed to update positions and velocities via stochastic dynamics

SEARCH Subroutine

"search" is a line search minimizer based upon parabolic extrapolation and cubic interpolation using both function and gradient values; if forced to search in an uphill direction, return is after the initial step

SETIME Subroutine

"setime" initializes the elapsed interval CPU timer

SHAKEUP Subroutine

"shakeup" initializes any holonomic constraints for use with the rattle algorithm during molecular dynamics

SIGMOID Function

"sigmoid" implements a normalized sigmoidal function on the interval [0,1]; the curves connect (0,0) to (1,1) and have a cooperativity controlled by beta, they approach a straight line as $\beta \rightarrow 0$ and get more nonlinear as β increases

SLATER Subroutine

"slater" is a general routine for computing the overlap integrals between two Slater-type orbitals

SMOOTH Subroutine

"smooth" sets extent of potential surface deformation for use with potential smoothing plus search, the diffusion equation method or Gaussian density annealing

SNIFFER Program

"sniffer" performs a global energy minimization using a discrete version of Griewank's global search trajectory

SNIFFER1 Function

"sniffer1" is a service routine that computes the energy and gradient for the Sniffer global optimization method

SOAK Subroutine

"soak" takes a currently defined solute system and places it into a solvent box, with removal of any solvent molecules that overlap the solute

SORT Subroutine

"sort" takes an input list of integers and sorts it into ascending order using the Heapsort algorithm

SORT2 Subroutine

"sort2" takes an input list of reals and sorts it into ascending order using the Heapsort algorithm; it also returns a key into the original ordering

SORT3 Subroutine

"sort3" takes an input list of integers and sorts it into ascending order using the Heapsort algorithm; it also returns a key into the original ordering

SORT4 Subroutine

"sort4" takes an input list of integers and sorts it into ascending absolute value using the Heapsort algorithm

SORT5 Subroutine

"sort5" takes an input list of integers and sorts it into ascending order based on each value modulo "m"

SORT6 Subroutine

"sort6" takes an input list of character strings and sorts it into alphabetical order using the Heapsort algorithm

SORT7 Subroutine

"sort7" takes an input list of character strings and sorts it into alphabetical order using the Heapsort algorithm; it also returns a key into the original ordering

SORT8 Subroutine

"sort8" takes an input list of integers and sorts it into ascending order using the Heapsort algorithm, duplicate values are removed from the final sorted list

SPACEFILL Program

"spacefill" computes the surface area and volume of a structure; the van der Waals, accessible-excluded, and contact-reentrant definitions are available

SPECTRUM Program

"spectrum" computes a power spectrum over a wavelength range from the velocity autocorrelation as a function of time

SQUARE Subroutine

"square" is a nonlinear least squares routine derived from the IMSL routine BCLSF and More's Minpack routine LMDER; the Jacobian is estimated by finite differences and bounds can be specified for the variables to be refined

SUFFIX Subroutine

"suffix" checks a filename for the presence of an extension, and appends an extension if none is found

SUPERPOSE Program

"superpose" takes pairs of structures and superimposes them in the optimal least squares sense; it will attempt to match all atom pairs or only those specified by the user

SURFACE Subroutine

"surface" performs an analytical computation of the weighted solvent accessible surface area of each atom and the first derivatives of the area with respect to Cartesian coordinates

SURFATOM Subroutine

"surfatom" performs an analytical computation of the surface area of a specified atom; a simplified version of "surface"

SWITCH Subroutine

"switch" sets the coefficients used by the fifth and seventh order polynomial switching functions for spherical cutoffs

SYBYLXYZ Program

"sybylxyz" takes as input a Sybyl MOL2 coordinates file, converts to and then writes out Cartesian coordinates

SYMMETRY Subroutine

"symmetry" applies symmetry operators to the fractional coordinates of the asymmetric unit in order to generate the symmetry related atoms of the full unit cell

TANGENT Subroutine

"tangent" finds the projected gradient on the synchronous transit path for a point along the transit pathway

TEMPER Subroutine

"temper" maintains a constant desired temperature via either Berendsen's velocity scaling coupled to an external temperature bath or Andersen's stochastic collision method

TESTGRAD Program

"testgrad" computes and compares the analytical and numerical gradient vectors of the potential energy function with respect to Cartesian coordinates

TESTHESS Program

"testhess" computes and compares the analytical and numerical Hessian matrices of the potential energy function with respect to Cartesian coordinates

TESTLIGHT Program

"testlight" performs a set of timing tests to compare the evaluation of potential energy and energy/gradient using the method of lights with a double loop over all atom pairs

TESTROT Program

"testrot" computes and compares the analytical and numerical gradient vectors of the potential energy function with respect to rotatable torsional angles

TIMER Program

"timer" measures the CPU time required for file reading and parameter assignment, potential energy computation, energy and gradient computation, and Hessian matrix evaluation

TIMEROT Program

"timerot" measures the CPU time required for file reading and parameter assignment, potential energy computation, energy and gradient over torsions, and torsional angle Hessian matrix evaluation

TNCG Subroutine

"tncg" implements a truncated Newton optimization algorithm in which a preconditioned linear conjugate gradient method is used to approximately solve Newton's equations; special features include use of an explicit sparse Hessian or finite-difference gradient-Hessian products within the PCG iteration; the exact Newton search directions can be used optionally; by default the algorithm checks for negative curvature to prevent convergence to a stationary point having negative eigenvalues; if a saddle point is desired this test can be removed by disabling "negtest"

TNSOLVE Subroutine

"tnsolve" uses a linear conjugate gradient method to find an approximate solution to the set of linear equations represented in matrix form by $H\mathbf{p} = -\mathbf{g}$ (Newton's equations)

TORPHASE Subroutine

"torphase" sets the n-fold amplitude and phase values for each torsion via sorting of the input parameters

TORQUE Subroutine

"torque" takes the torque values on sites defined by local coordinate frames and distributes them to convert to forces on the original sites and sites specifying the local frames

TORQUE1 Subroutine

"torque1" takes the torque value on a site defined by a local coordinate frame and distributes it to convert to forces on the original site and sites specifying the local frame

TORSER Function

"torse" computes the torsional error function and its first derivatives with respect to the atomic Cartesian coordinates based on the deviation of specified torsional angles from desired values, the contained bond angles are also restrained to avoid a numerical instability

TORSIONS Subroutine

"torsions" finds the total number of dihedral angles and the numbers of the four atoms defining each dihedral angle

TORUS Subroutine

"torus" sets a list of all of the temporary torus positions by testing for a torus between each atom and its neighbors

TOTERR Function

"toterr" is the error function and derivatives for a distance geometry embedding; it includes components from the distance bounds, hard sphere contacts, local geometry, chirality and torsional restraint errors

TRANSIT Function

"transit" evaluates the synchronous transit function and gradient; linear and quadratic transit paths are available

TRIANGLE Subroutine

"triangle" smooths the upper and lower distance bounds via the triangle inequality using a full-matrix variant of the Floyd-Warshall shortest path algorithm; this routine is usually much slower than the sparse matrix shortest path methods in "geodesic" and "trifix", and should be used only for comparison with answers generated by those routines

TRIFIX Subroutine

"trifix" rebuilds both the upper and lower distance bound matrices following tightening of one or both of the bounds between a specified pair of atoms, "p" and "q", using a modification of Murchland's shortest path update algorithm

TRIMTEXT Function

"trimtext" finds and returns the location of the last non-blank character before the first null character in an input text string; the function returns zero if no such character is found

TRIPLE Function

"triple" finds the triple product of three vectors; used as a service routine by the Connolly surface area and volume computation

TRUST Subroutine

"trust" updates the model trust region for a nonlinear least squares calculation; this version is based on the ideas found in NL2SOL and in Dennis and Schnabel's book

UDIRECT1 Subroutine

"udirect1" computes the reciprocal space contribution of the permanent atomic multipole moments to the electrostatic field for use in finding the direct induced dipole moments via a regular Ewald summation

UDIRECT2 Subroutine

"udirect2" computes the real space contribution of the permanent atomic multipole moments to the electrostatic field for use in finding the direct induced dipole moments via a regular Ewald summation

UFIELD Subroutine

"ufield" finds the field at each polarizable site due to the induced dipoles at the other sites using Thole's method to damp the field at close range

UMUTUAL1 Subroutine

"umutual1" computes the reciprocal space contribution of the induced atomic dipole moments to the electrostatic field for use in iterative calculation of induced dipole moments via a regular Ewald summation

UMUTUAL2 Subroutine

"umutual2" computes the real space contribution of the induced atomic dipole moments to the electrostatic field for use in iterative calculation of induced dipole moments via a regular Ewald summation

UNITCELL Subroutine

"unitcell" gets the periodic boundary box size and related values from an external keyword file

UPCASE Subroutine

"upcase" converts a text string to all upper case letters

VAM Subroutine

"vam" takes the analytical molecular surface defined as a collection of spherical and toroidal polygons and uses it to compute the volume and surface area

VCROSS Subroutine

"vcross" finds the cross product of two vectors

VDWERR Function

"vdwerr" is the hard sphere van der Waals bound error function and derivatives that penalizes close nonbonded contacts, pairwise neighbors are generated via the method of lights

VECANG Function

"vecang" finds the angle between two vectors handed with respect to a coordinate axis; returns an angle in the range $[0, 2\pi]$

VERLET Subroutine

"verlet" performs a single molecular dynamics time step by means of the velocity Verlet multistep recursion formula

VERSION Subroutine

"version" checks the name of a file about to be opened; if "old" status is passed, the name of the highest current version is returned; if "new" status is passed the filename of the next available unused version is generated

VIBRATE Program

"vibrate" performs a vibrational normal mode analysis; the Hessian matrix of second derivatives is determined and then diagonalized both directly and after mass weighting; output consists of the eigenvalues of the force constant matrix as well as the vibrational frequencies and displacements

VIBROT Program

VNORM Subroutine

"vnorm" normalizes a vector to unit length; used as a service routine by the Connolly surface area and volume computation

VOLUME Subroutine

"volume" calculates the excluded volume via the Connolly analytical volume and surface area algorithm

VOLUME1 Subroutine

"volumel" calculates first derivatives of the total excluded volume with respect to the Cartesian coordinates of each atom

VOLUME2 Subroutine

"volume2" calculates second derivatives of the total excluded volume with respect to the Cartesian coordinates of the atoms

WATSON Subroutine

"watson" uses a rigid-body optimization to approximately align the paired strands of a nucleic acid double helix

WATSON1 Function

"watson1" is a service routine that computes the energy and gradient for optimally conditioned variable metric optimization of rigid bodies

WRITEOUT Subroutine

"writeout" is used by each of the optimization routines to save intermediate atomic coordinates to a disk file

XTALERR Subroutine

XTALFIT Program

"xtalfit" computes an optimized set of potential energy parameters for user specified van der Waals and electrostatic interactions by fitting to crystal structure, lattice energy and monomer dipole moment data

XTALLAT1 Function

"xtalmol1" is a service routine that computes the energy and numerical gradient with respect to the six lattice lengths and angles for a crystal energy minimization

XTALMIN Program

"xtalmin" performs a full crystal energy minimization by alternating cycles of truncated Newton optimization over atomic coordinates with variable metric optimization over the six lattice dimensions and angles

XTALMOL1 Function

"xtalmol1" is a service routine that computes the energy and gradient with respect to the atomic Cartesian coordinates for a crystal energy minimization

XTALMOL2 Subroutine

"xtalmol2" is a service routine that computes the sparse matrix Hessian elements with respect to the atomic Cartesian coordinates for a crystal energy minimization

XTALMOVE Subroutine

XTALPRM Subroutine

"xtalprm" stores or retrieves a crystal structure; used to make a previously stored structure the currently active structure, or to store a structure for later use; only the intermolecular energy terms are provided for

XTALWRT Subroutine

XYZATM Subroutine

"xyzatm" computes the Cartesian coordinates of a single atom from its defining internal coordinate values

XYZEDIT Program

"xyzedit" provides for modification and manipulation of the contents of a Cartesian coordinates file

XYZINT Program

"xyzint" takes as input a Cartesian coordinates file, then converts to and writes out an internal coordinates file

XYZPDB Program

"xyzpdb" takes as input a Cartesian coordinates file, then converts to and writes out a Protein Data Bank file

XYZRIGID Subroutine

"xyzrigid" computes the center of mass and Euler angle rigid body coordinates for each atom group in the system

XYZSYBYL Program

"xyzsybyl" takes as input a Cartesian coordinates file, converts to and then writes out a Sybyl MOL2 file

ZATOM Subroutine

"zatom" adds an atom to the end of the current Z-matrix and then increments the atom counter; atom type, defining atoms and internal coordinates are passed as arguments

ZHELP Subroutine

"zhelp" prints the general information and instructions for the Z-matrix editing program

ZVALUE Subroutine

"zvalue" gets user supplied values for selected coordinates as needed by the internal coordinate editing program

10. Contents of Common Block Variables

The Fortran common blocks found in the TINKER package are listed below along with a brief description of the contents of each variable in each common block. Each individual common block is present as a separate ".i" file in the /source subdirectory. A source code listing containing each of the source code modules and each of the common blocks can be produced by running the "listing.make" script found in the distribution.

ACTION

total number of each energy term computed

| | |
|-------|---|
| neb | number of bond stretch energy terms computed |
| nea | number of angle bend energy terms computed |
| neba | number of stretch-bend energy terms computed |
| neub | number of Urey-Bradley energy terms computed |
| neaa | number of angle-angle energy terms computed |
| neopb | number of out-of-plane bend energy terms computed |
| neopd | number of out-of-plane distance energy terms computed |
| neid | number of improper dihedral energy terms computed |
| neit | number of improper torsion energy terms computed |
| net | number of torsional energy terms computed |
| nebt | number of stretch-torsion energy terms computed |
| nett | number of torsion-torsion energy terms computed |
| nev | number of van der Waals energy terms computed |
| nec | number of charge-charge energy terms computed |
| necd | number of charge-dipole energy terms computed |
| ned | number of dipole-dipole energy terms computed |
| nem | number of multipole energy terms computed |
| nep | number of polarization energy terms computed |
| new | number of Ewald summation energy terms computed |
| ner | number of reaction field energy terms computed |
| nes | number of solvation energy terms computed |
| nelf | number of metal ligand field energy terms computed |
| neg | number of geometric restraint energy terms computed |
| nex | number of extra energy terms computed |

ALIGN

information for superposition of structures

| | |
|------|--|
| wfit | weights assigned to atom pairs during superposition |
| nfit | number of atoms to use in superimposing two structures |
| ifit | atom numbers of pairs of atoms to be superimposed |

ANALYZ

energy components partitioned over atoms

| | |
|-------|---|
| aeb | bond stretch energy partitioned over atoms |
| aea | angle bend energy partitioned over atoms |
| aeba | stretch-bend energy partitioned over atoms |
| aeub | Urey-Bradley energy partitioned over atoms |
| aeaa | angle-angle energy partitioned over atoms |
| aeopb | out-of-plane bend energy partitioned over atoms |
| aeopd | out-of-plane distance energy partitioned over atoms |
| aeid | improper dihedral energy partitioned over atoms |

| | |
|------|---|
| aeit | improper torsion energy partitioned over atoms |
| aet | torsional energy partitioned over atoms |
| aebt | stretch-torsion energy partitioned over atoms |
| aett | torsion-torsion energy partitioned over atoms |
| aev | van der Waals energy partitioned over atoms |
| aec | charge-charge energy partitioned over atoms |
| aecd | charge-dipole energy partitioned over atoms |
| aed | dipole-dipole energy partitioned over atoms |
| aem | multipole energy partitioned over atoms |
| aep | polarization energy partitioned over atoms |
| aer | reaction field energy partitioned over atoms |
| aes | solvation energy partitioned over atoms |
| aelf | metal ligand field energy partitioned over atoms |
| age | geometric restraint energy partitioned over atoms |
| aex | extra energy term partitioned over atoms |

ANGANG

angle-angle terms in current structure

| | |
|---------|---|
| kaa | force constant for angle-angle cross terms |
| nangang | total number of angle-angle interactions |
| iaa | angle numbers used in each angle-angle term |

ANGLE

bond angles within the current structure

| | |
|--------|--|
| ak | harmonic angle force constant (kcal/mole/rad**2) |
| anat | ideal bond angle or phase shift angle (degrees) |
| afld | periodicity for Fourier bond angle term |
| nangle | total number of bond angles in the system |
| iang | numbers of the atoms in each bond angle |
| angtyp | potential energy function type for each bond angle |

ANGPOT

specifics of bond angle functional forms

| | |
|----------|--|
| cang | cubic coefficient in angle bending potential |
| qang | quartic coefficient in angle bending potential |
| pang | quintic coefficient in angle bending potential |
| sang | sextic coefficient in angle bending potential |
| angunit | convert angle force constant to kcal/mole/deg**2 |
| stbnunit | convert str-bend constant to kcal/mole/deg-Ang**2 |
| aaunit | convert angle-angle constant to kcal/mole/deg**2 |
| opbunit | convert out-of-plane bend force to kcal/mole/deg**2 |
| opdunit | convert out-of-plane distance to kcal/mole/Ang**2 |
| mm2stbn | logical flag governing use of MM2-style stretch-bend |

ARGUE

command line arguments at program startup

| | |
|---------|---|
| maxarg | maximum number of command line arguments |
| narg | number of command line arguments to the program |
| listarg | flag to mark available command line arguments |
| arg | strings containing the command line arguments |

ATMLST

local geometry terms involving each atom

| | |
|---------------|---|
| bndlist | list of the bond numbers involving each atom |
| anglist | list of the angle numbers centered on each atom |
| ATMTYP | atomic properties for each current atom |
| mass | atomic weight for each atom in the system |
| tag | integer atom labels from input coordinates file |
| class | atom class number for each atom in the system |
| atomic | atomic number for each atom in the system |
| valence | valence number for each atom in the system |
| name | atom name for each atom in the system |
| story | descriptive type for each atom in system |
| ATOMS | number, position and type of current atoms |
| x | current x-coordinate for each atom in the system |
| y | current y-coordinate for each atom in the system |
| z | current z-coordinate for each atom in the system |
| n | total number of atoms in the current system |
| type | atom type number for each atom in the system |
| BATH | temperature and pressure control parameters |
| kelvin | target value for the system temperature (K) |
| atmsph | target value for the system pressure (atm) |
| tautemp | time constant in psec for temperature bath coupling |
| taupres | time constant in psec for pressure bath coupling |
| compress | isothermal compressibility of medium (atm ⁻¹) |
| collide | collision frequency for Andersen thermostat |
| isothermal | logical flag governing use of temperature bath |
| isobaric | logical flag governing use of pressure bath |
| thermostat | type of thermostat, either Berendsen or Andersen |
| BNDPOT | specifics of bond stretch functional forms |
| cbnd | cubic coefficient in bond stretch potential |
| qbnd | quartic coefficient in bond stretch potential |
| bndunit | convert bond force constant to kcal/mole/Ang**2 |
| bndtyp | type of bond stretch potential energy function |
| BOND | covalent bonds in the current structure |
| bk | bond stretch force constants (kcal/mole/Ang**2) |
| bl | ideal bond length values in Angstroms |
| nbond | total number of bond stretches in the system |
| ibnd | numbers of the atoms in each bond stretch |
| BORDER | bond orders for a conjugated pisystem |
| pbpl | pi-bond orders for bonds in "planar" pisystem |
| pnpl | pi-bond orders for bonds in "nonplanar" pisystem |

BOUND

polycut
polycut2
use_bounds
use_image
use_replica
use_polymer

control of periodic boundary conditions

cutoff distance for infinite polymer nonbonds
square of infinite polymer nonbond cutoff
flag to use periodic boundary conditions
flag to use images for periodic system
flag to use replicates for periodic system
flag to mark presence of infinite polymer

BOXES

xbox
ybox
zbox
alpha
beta
gamma
xbox2
ybox2
zbox2
box34
recip
volbox
beta_sin
beta_cos
gamma_sin
gamma_cos
beta_term
gamma_term
orthogonal
monoclinic
triclinic
octahedron
spacegrp

parameters for periodic boundary conditions

length in Angs of a-axis of periodic box
length in Angs of b-axis of periodic box
length in Angs of c-axis of periodic box
angle in degrees between b- and c-axes of box
angle in degrees between a- and c-axes of box
angle in degrees between a- and b-axes of box
half of the a-axis length of periodic box
half of the b-axis length of periodic box
half of the c-axis length of periodic box
three-fourths axis length of truncated octahedron
reciprocal lattice vectors as matrix columns
volume in Ang**3 of the periodic box
sine of the beta periodic box angle
cosine of the beta periodic box angle
sine of the gamma periodic box angle
cosine of the gamma periodic box angle
term used in generating triclinic box
term used in generating triclinic box
flag to mark periodic box as orthogonal
flag to mark periodic box as monoclinic
flag to mark periodic box as triclinic
flag to mark box as truncated octahedron
space group symbol for the unitcell type

CELL

xcell
ycell
zcell
xccl2
yccl2
zccl2
ncell
icell

periodic boundaries using replicated cells

length of the a-axis of the complete replicated cell
length of the b-axis of the complete replicated cell
length of the c-axis of the complete replicated cell
half the length of the a-axis of the replicated cell
half the length of the b-axis of the replicated cell
half the length of the c-axis of the replicated cell
total number of cell replicates for periodic boundaries
offset along axes for each replicate periodic cell

CENTRE

xcm
ycm
zcm

atom coordinates relative to center of mass

offset of each atom from center of mass x-coordinate
offset of each atom from center of mass y-coordinate
offset of each atom from center of mass z-coordinate

CHARGE

pchg
nion
iion
jion
kion
chglist

partial charges for the current structure

magnitude of the partial charges (e-)
total number of partial charges in system
number of the atom site for each partial charge
neighbor generation site for each partial charge
cutoff switching site for each partial charge
partial charge site for each atom (0=no charge)

CHGPOT

dielec
c2scale
c3scale
c4scale
c5scale
neutnbr
neutcut

specifics of charge-charge functional form

dielectric constant for electrostatic interactions
factor by which 1-2 charge interactions are scaled
factor by which 1-3 charge interactions are scaled
factor by which 1-4 charge interactions are scaled
factor by which 1-5 charge interactions are scaled
logical flag governing use of neutral group neighbors
logical flag governing use of neutral group cutoffs

CHRONO

cputim

timing statistics for the current program

elapsed cpu time in seconds since start of program

COUPLE

maxn13
maxn14
maxn15
n12
il2
n13
il3
n14
il4
n15
il5

near-neighbor atom connectivity lists

maximum number of atoms 1-3 connected to an atom
maximum number of atoms 1-4 connected to an atom
maximum number of atoms 1-5 connected to an atom
number of atoms directly bonded to each atom
atom numbers of atoms 1-2 connected to each atom
number of atoms in a 1-3 relation to each atom
atom numbers of atoms 1-3 connected to each atom
number of atoms in a 1-4 relation to each atom
atom numbers of atoms 1-4 connected to each atom
number of atoms in a 1-5 relation to each atom
atom numbers of atoms 1-5 connected to each atom

CUTOFF

vdwcut
chgcut
dplcut
mpolecut
vdwtaper
chgtaper
dpltaper
mpoletaper
ewaldcut
use_ewald
use_lights

cutoff distances for energy interactions

cutoff distance for van der Waals interactions
cutoff distance for charge-charge interactions
cutoff distance for dipole-dipole interactions
cutoff distance for atomic multipole interactions
distance at which van der Waals switching begins
distance at which charge-charge switching begins
distance at which dipole-dipole switching begins
distance at which atomic multipole switching begins
cutoff distance for direct space Ewald summation
logical flag governing use of Ewald summation term
logical flag to use method of lights neighbors

DERIV

| | |
|-------|--|
| deb | bond stretch Cartesian coordinate derivatives |
| dea | angle bend Cartesian coordinate derivatives |
| deba | stretch-bend Cartesian coordinate derivatives |
| deub | Urey-Bradley Cartesian coordinate derivatives |
| deaa | angle-angle Cartesian coordinate derivatives |
| deopb | out-of-plane bend Cartesian coordinate derivatives |
| deopd | out-of-plane distance Cartesian coordinate derivatives |
| deid | improper dihedral Cartesian coordinate derivatives |
| deit | improper torsion Cartesian coordinate derivatives |
| det | torsional Cartesian coordinate derivatives |
| debt | stretch-torsion Cartesian coordinate derivatives |
| dett | torsion-torsion Cartesian coordinate derivatives |
| dev | van der Waals Cartesian coordinate derivatives |
| dec | charge-charge Cartesian coordinate derivatives |
| decdd | charge-dipole Cartesian coordinate derivatives |
| ded | dipole-dipole Cartesian coordinate derivatives |
| dem | multipole Cartesian coordinate derivatives |
| dep | polarization Cartesian coordinate derivatives |
| der | reaction field Cartesian coordinate derivatives |
| des | solvation Cartesian coordinate derivatives |
| delf | metal ligand field Cartesian coordinate derivatives |
| deg | geometric restraint Cartesian coordinate derivatives |
| dex | extra energy term Cartesian coordinate derivatives |

DIPOLE

| | |
|---------|--|
| bdpl | magnitude of each of the dipoles (Debyes) |
| sdpl | position of each dipole between defining atoms |
| ndipole | total number of dipoles in the system |
| idpl | numbers of atoms that define each dipole |

DISGEO

| | |
|------------|---|
| bnd | distance geometry upper and lower bounds matrix |
| vdwrad | hard sphere radii for distance geometry atoms |
| vhir | signed volume values for chirality constraints |
| compact | index of local distance compaction on embedding |
| pathmax | maximum value of upper bound after smoothing |
| vdwmax | maximum value of hard sphere sum for an atom pair |
| nhir | total number of chirality constraints |
| ichir | numbers of atoms in each chirality constraint |
| use_invert | flag to use enantiomer closest to input structure |
| use_anneal | flag to use simulated annealing refinement |

DOMEGA

| | |
|------|--|
| teb | bond stretch derivatives over torsions |
| tea | angle bend derivatives over torsions |
| teba | stretch-bend derivatives over torsions |
| teub | Urey-Bradley derivatives over torsions |

Cartesian coordinate derivative components

atom & bond dipoles for current structure

distance geometry bounds and parameters

derivative components over dihedrals

| | |
|-------|---|
| teaa | angle-angle derivatives over torsions |
| teopb | out-of-plane bend derivatives over torsions |
| teopd | out-of-plane distance derivatives over torsions |
| teid | improper dihedral derivatives over torsions |
| teit | improper torsion derivatives over torsions |
| tet | torsional derivatives over torsions |
| tebt | stretch-torsion derivatives over torsions |
| tett | torsion-torsion derivatives over torsions |
| tev | van der Waals derivatives over torsions |
| tec | charge-charge derivatives over torsions |
| tecd | charge-dipole derivatives over torsions |
| ted | dipole-dipole derivatives over torsions |
| tem | atomic multipole derivatives over torsions |
| tep | polarization derivatives over torsions |
| ter | reaction field derivatives over torsions |
| tes | solvation derivatives over torsions |
| telf | metal ligand field derivatives over torsions |
| teg | geometric restraint derivatives over torsions |
| tex | extra energy term derivatives over torsions |

ENERGI

individual potential energy components

| | |
|------|--|
| eb | bond stretch potential energy of the system |
| ea | angle bend potential energy of the system |
| eba | stretch-bend potential energy of the system |
| eub | Urey-Bradley potential energy of the system |
| ea | angle-angle potential energy of the system |
| eopb | out-of-plane bend potential energy of the system |
| eopd | out-of-plane distance potential energy of the system |
| eid | improper dihedral potential energy of the system |
| eit | improper torsion potential energy of the system |
| et | torsional potential energy of the system |
| ebt | stretch-torsion potential energy of the system |
| ett | torsion-torsion potential energy of the system |
| ev | van der Waals potential energy of the system |
| ec | charge-charge potential energy of the system |
| ecd | charge-dipole potential energy of the system |
| ed | dipole-dipole potential energy of the system |
| em | atomic multipole potential energy of the system |
| ep | polarization potential energy of the system |
| er | reaction field potential energy of the system |
| es | solvation potential energy of the system |
| elf | metal ligand field potential energy of the system |
| eg | geometric restraint potential energy of the system |
| ex | extra term potential energy of the system |

EWALD

parameters for regular or PM Ewald summation

| | |
|---------|---|
| aewald | Ewald convergence coefficient value (Ang-1) |
| frecip | fractional cutoff value for reciprocal sphere |
| tinfoil | flag governing use of tinfoil boundary conditions |

EWREG

maxvec
ejc
ejs
ekc
eks
elc
els

exponential factors for regular Ewald sum

maximum number of k-vectors per reciprocal axis
exponential factors for cosine along the j-axis
exponential factors for sine along the j-axis
exponential factors for cosine along the k-axis
exponential factors for sine along the k-axis
exponential factors for cosine along the l-axis
exponential factors for sine along the l-axis

FACES

maxnbr
maxtt
maxt
maxp
maxv
maxen
maxfn
maxc
maxep
maxfs
maxcy
mxcyep
maxfp
mxfpCY

variables for Connolly area and volume

maximum number of neighboring atom pairs
maximum number of temporary tori
maximum number of total tori
maximum number of probe positions
maximum number of vertices
maximum number of concave edges
maximum number of concave faces
maximum number of circles
maximum number of convex edges
maximum number of saddle faces
maximum number of cycles
maximum number of cycle convex edges
maximum number of convex faces
maximum number of convex face cycles

FIELDS

biotyp
forcefield

molecular mechanics force field description

force field atom type of each biopolymer type
string used to describe the current forcefield

FILES

nprior
ldir
leng
filename
outfile

name and number of current structure files

number of previously existing cycle files
length in characters of the directory name
length in characters of the base filename
base filename used by default for all files
output filename used for intermediate results

FRACS

xfrac
yfrac
zfrac

atom distances to molecular center of mass

fractional coordinate along a-axis of center of mass
fractional coordinate along b-axis of center of mass
fractional coordinate along c-axis of center of mass

GROUP

grpmass
wgrp
grpnum
ngrp

partitioning of system into atom groups

total mass of all the atoms in each group
weight for each set of group-group interactions
original group number for each nonempty group
total number of atom groups in the system

| | |
|----------------|---|
| kgrp | contiguous list of the atoms in each group |
| igrp | first and last atom of each group in the list |
| grplist | number of the group to which each atom belongs |
| use_group | flag to use partitioning of system into groups |
| use_intra | flag to include only intragroup interactions |
| use_inter | flag to include only intergroup interactions |
| HES CUT | cutoff value for Hessian matrix elements |
| hesscut | magnitude of smallest allowed Hessian element |
| HESS N | Cartesian Hessian elements for a single atom |
| hessx | Hessian elements for x-component of current atom |
| hessy | Hessian elements for y-component of current atom |
| hessz | Hessian elements for z-component of current atom |
| IMPROP | improper dihedrals in the current structure |
| kprop | force constant values for improper dihedral angles |
| vprop | ideal improper dihedral angle value in degrees |
| niprop | total number of improper dihedral angles in the system |
| iiprop | numbers of the atoms in each improper dihedral angle |
| IMPTOR | improper torsions in the current structure |
| itors1 | 1-fold amplitude and phase for each improper torsion |
| itors2 | 2-fold amplitude and phase for each improper torsion |
| itors3 | 3-fold amplitude and phase for each improper torsion |
| nitors | total number of improper torsional angles in the system |
| iitors | numbers of the atoms in each improper torsional angle |
| INFORM | control values for I/O and program flow |
| digits | decimal places output for energy and coordinates |
| iprint | steps between status printing (0=no printing) |
| iwrite | steps between coordinate dumps (0=no dumps) |
| verbose | logical flag to turn on extra information |
| debug | logical flag to turn on full debug printing |
| holdup | logical flag to wait for carriage return on exit |
| abort | logical flag to stop execution at next chance |
| INTER | sum of intermolecular energy components |
| einter | total intermolecular potential energy |
| IOUNIT | Fortran input/output (I/O) unit numbers |
| iout | Fortran I/O unit for major output (default=6) |
| input | Fortran I/O unit for major input (default=5) |
| KANANG | forcefield parameters for angle-angle terms |

anan

angle-angle cross term parameters for each atom class

KANGS

forcefield parameters for bond angle bending

maxna

maximum number of harmonic angle bend parameter entries

maxna5

maximum number of 5-membered ring angle bend entries

maxna4

maximum number of 4-membered ring angle bend entries

maxna3

maximum number of 3-membered ring angle bend entries

maxnaf

maximum number of Fourier angle bend parameter entries

acon

force constant parameters for harmonic angle bends

acon5

force constant parameters for 5-ring angle bends

acon4

force constant parameters for 4-ring angle bends

acon3

force constant parameters for 3-ring angle bends

aconf

force constant parameters for Fourier angle bends

ang

bond angle parameters for harmonic angle bends

ang5

bond angle parameters for 5-ring angle bends

ang4

bond angle parameters for 4-ring angle bends

ang3

bond angle parameters for 3-ring angle bends

angf

phase shift angle and periodicity for Fourier bends

ka

string of atom classes for harmonic angle bends

ka5

string of atom classes for 5-ring angle bends

ka4

string of atom classes for 4-ring angle bends

ka3

string of atom classes for 3-ring angle bends

kaf

string of atom classes for Fourier angle bends

KATOMS

forcefield parameters for the atom types

weight

average atomic mass of each atom type

atmc1s

atom class number for each of the atom types

atmnum

atomic number for each of the atom types

ligand

number of atoms to be attached to each atom type

symbol

modified atomic symbol for each atom type

describe

string identifying each of the atom types

KBONDS

forcefield parameters for bond stretching

maxnb

maximum number of bond stretch parameter entries

maxnb5

maximum number of 5-membered ring bond stretch entries

maxnb4

maximum number of 4-membered ring bond stretch entries

maxnb3

maximum number of 3-membered ring bond stretch entries

maxnel

maximum number of electronegativity bond corrections

bcon

force constant parameters for harmonic bond stretch

bcon5

force constant parameters for 5-ring bond stretch

bcon4

force constant parameters for 4-ring bond stretch

bcon3

force constant parameters for 3-ring bond stretch

blen

bond length parameters for harmonic bond stretch

blen5

bond length parameters for 5-ring bond stretch

blen4

bond length parameters for 4-ring bond stretch

blen3

bond length parameters for 3-ring bond stretch

dlen

electronegativity bond length correction parameters

kb

string of atom classes for harmonic bond stretch

| | |
|-----|--|
| kb5 | string of atom classes for 5-ring bond stretch |
| kb4 | string of atom classes for 4-ring bond stretch |
| kb3 | string of atom classes for 3-ring bond stretch |
| kel | string of atom classes for electronegativity corrections |

KCHARGE

forcefield parameters for partial charges

| | |
|-----|--|
| chg | partial charge parameters for each atom type |
|-----|--|

KDIPOL

forcefield parameters for bond dipoles

| | |
|--------|--|
| maxnd | maximum number of bond dipole parameter entries |
| maxnd5 | maximum number of 5-membered ring dipole entries |
| maxnd4 | maximum number of 4-membered ring dipole entries |
| maxnd3 | maximum number of 3-membered ring dipole entries |
| dpl | dipole moment parameters for bond dipoles |
| dpl5 | dipole moment parameters for 5-ring dipoles |
| dpl4 | dipole moment parameters for 4-ring dipoles |
| dpl3 | dipole moment parameters for 3-ring dipoles |
| pos | dipole position parameters for bond dipoles |
| pos5 | dipole position parameters for 5-ring dipoles |
| pos4 | dipole position parameters for 4-ring dipoles |
| pos3 | dipole position parameters for 3-ring dipoles |
| kd | string of atom classes for bond dipoles |
| kd5 | string of atom classes for 5-ring dipoles |
| kd4 | string of atom classes for 4-ring dipoles |
| kd3 | string of atom classes for 3-ring dipoles |

KEYS

contents of current keyword parameter file

| | |
|---------|---|
| nkey | number of nonblank lines in the keyword file |
| keyline | contents of each individual keyword file line |

KGEOMS

parameters for the geometrical restraints

| | |
|-------|--|
| depth | depth of shallow Gaussian basin restraint |
| width | exponential width coefficient of Gaussian basin |
| rwall | radius of spherical droplet boundary restraint |
| xpfix | x-coordinate target for each restrained position |
| ypfix | y-coordinate target for each restrained position |
| zpfix | z-coordinate target for each restrained position |
| pfix | flat-well range and force constant for each position |
| dfix | target range and force constant for each distance |
| afix | target range and force constant for each angle |
| tfix | target range and force constant for each torsion |
| npfix | number of position restraints to be applied |
| ipfix | atom number involved in each position restraint |
| ndfix | number of distance restraints to be applied |
| idfix | atom numbers defining each distance restraint |
| nafix | number of angle restraints to be applied |
| iafix | atom numbers defining each angle restraint |
| ntfix | number of torsional restraints to be applied |

| | |
|---------------|--|
| itfix | atom numbers defining each torsional restraint |
| use_basin | logical flag governing use of Gaussian basin |
| use_wall | logical flag governing use of droplet boundary |
| KHBOND | forcefield parameters for H-bonding terms |
| maxnhb | maximum number of hydrogen bonding pair entries |
| radhb | radius parameter for hydrogen bonding pairs |
| epshb | well depth parameter for hydrogen bonding pairs |
| khh | string of atom types for hydrogen bonding pairs |
| KIPROP | forcefield parameters for improper dihedral |
| maxndi | maximum number of improper dihedral parameter entries |
| dcon | force constant parameters for improper dihedrals |
| tdi | ideal dihedral angle values for improper dihedrals |
| kdi | string of atom classes for improper dihedral angles |
| KITORS | forcefield parameters for improper torsions |
| maxnti | maximum number of improper torsion parameter entries |
| ti1 | torsional parameters for improper 1-fold rotation |
| ti2 | torsional parameters for improper 2-fold rotation |
| ti3 | torsional parameters for improper 3-fold rotation |
| kti | string of atom classes for improper torsional parameters |
| KMULTI | forcefield parameters for atomic multipoles |
| maxnmp | maximum number of atomic multipole parameter entries |
| multip | atomic monopole, dipole and quadrupole values |
| mpaxis | type of local axis definition for atomic multipoles |
| kmp | string of atom types for atomic multipoles |
| KOPBND | forcefield parameters for out-of-plane bend |
| maxnopb | maximum number of out-of-plane bending entries |
| copb | force constant parameters for out-of-plane bending |
| kaopb | string of atom classes for out-of-plane bending |
| KOPDST | forcefield parameters for out-plane distance |
| maxnopb | maximum number of out-of-plane distance entries |
| copb | force constant parameters for out-of-plane distance |
| kaopb | string of atom classes for out-of-plane distance |
| KORBS | forcefield parameters for pisystem orbitals |
| maxnpi | maximum number of pisystem bond parameter entries |
| electron | number of pi-electrons for each atom class |
| ionize | ionization potential for each atom class |
| repulse | repulsion integral value for each atom class |
| sslope | slope for bond stretch vs. pi-bond order |

| | |
|---------------|---|
| tslope | slope for 2-fold torsion vs. pi-bond order |
| kpi | string of atom classes for pisystem bonds |
| KPOLR | forcefield parameters for polarizability |
| polr | dipole polarizability parameters for each atom type |
| pgrp | connected types in polarization group of each atom type |
| KSTBND | forcefield parameters for stretch-bending |
| stbn | stretch-bending parameters for each atom class |
| KSTTOR | forcefield parameters for stretch-torsions |
| maxnbt | maximum number of stretch-torsion parameter entries |
| btcon | force constant parameters for stretch-torsion |
| kbt | string of atom classes for bonds in stretch-torsion |
| KTORSN | forcefield parameters for torsional angles |
| maxnt | maximum number of torsional angle parameter entries |
| maxnt5 | maximum number of 5-membered ring torsion entries |
| maxnt4 | maximum number of 4-membered ring torsion entries |
| t1 | torsional parameters for standard 1-fold rotation |
| t2 | torsional parameters for standard 2-fold rotation |
| t3 | torsional parameters for standard 3-fold rotation |
| t4 | torsional parameters for standard 4-fold rotation |
| t5 | torsional parameters for standard 5-fold rotation |
| t6 | torsional parameters for standard 6-fold rotation |
| t15 | torsional parameters for 1-fold rotation in 5-ring |
| t25 | torsional parameters for 2-fold rotation in 5-ring |
| t35 | torsional parameters for 3-fold rotation in 5-ring |
| t45 | torsional parameters for 4-fold rotation in 5-ring |
| t55 | torsional parameters for 5-fold rotation in 5-ring |
| t65 | torsional parameters for 6-fold rotation in 5-ring |
| t14 | torsional parameters for 1-fold rotation in 4-ring |
| t24 | torsional parameters for 2-fold rotation in 4-ring |
| t34 | torsional parameters for 3-fold rotation in 4-ring |
| t44 | torsional parameters for 4-fold rotation in 4-ring |
| t54 | torsional parameters for 5-fold rotation in 4-ring |
| t64 | torsional parameters for 6-fold rotation in 4-ring |
| kt | string of atom classes for torsional angles |
| kt5 | string of atom classes for 5-ring torsions |
| kt4 | string of atom classes for 4-ring torsions |
| KURYBR | forcefield parameters for Urey-Bradley terms |
| maxnu | maximum number of Urey-Bradley parameter entries |
| ucon | force constant parameters for Urey-Bradley terms |
| dst13 | ideal 1-3 distance parameters for Urey-Bradley terms |
| ku | string of atom classes for Urey-Bradley terms |

KVDWPR

maxnvp
radpr
epspr
kvpr

forcefield parameters for special vdw terms

maximum number of special van der Waals pair entries
radius parameter for special van der Waals pairs
well depth parameter for special van der Waals pairs
string of atom classes for special van der Waals pairs

KVDWS

rad
eps
rad4
eps4
reduct

forcefield parameters for van der Waals terms

van der Waals radius parameter for each atom class
van der Waals well depth parameter for each atom class
van der Waals radius parameter in 1-4 interactions
van der Waals well depth parameter in 1-4 interactions
van der Waals reduction factor for each atom class

LIGHT

nlight
kbx
kby
kbz
kex
key
kez
locx
locy
locz
rgx
rgy
rgz

indices for method of lights pair neighbors

total number of sites for method of lights calculation
low index of neighbors of each site in the x-sorted list
low index of neighbors of each site in the y-sorted list
low index of neighbors of each site in the z-sorted list
high index of neighbors of each site in the x-sorted list
high index of neighbors of each site in the y-sorted list
high index of neighbors of each site in the z-sorted list
pointer from x-sorted list into original interaction list
pointer from y-sorted list into original interaction list
pointer from z-sorted list into original interaction list
pointer from original interaction list into x-sorted list
pointer from original interaction list into y-sorted list
pointer from original interaction list into z-sorted list

LINMIN

stpmin
stpmax
cappa
angmax
intmax

parameters for line search minimization

minimum step length in current line search direction
maximum step length in current line search direction
stringency of line search (0=tight < cappa < 1=loose)
maximum angle between search direction and -gradient
maximum number of inner iterations during line search

MATH

radian
pi
sqrtpi
logten
twosix

mathematical and geometrical constants

conversion factor from radians to degrees
numerical value of the geometric constant
numerical value of the square root of Pi
numerical value of the natural log of ten
numerical value of the sixth root of two

MDSTUF

nfree
velsave
uindsave

control of molecular dynamics trajectory

total number of degrees of freedom for a system
flag to save atomic velocity components to a file
flag to save induced atomic dipoles to a file

| | |
|---------------|---|
| integrate | type of molecular dynamics integration algorithm |
| MINIMA | general parameters for minimizations |
| fctmin | value below which function is deemed optimized |
| hguess | initial value for the H-matrix diagonal elements |
| maxiter | maximum number of iterations during optimization |
| nextiter | iteration number to use for the first iteration |
| MOLCUL | individual molecules within current system |
| molmass | molecular weight for each molecule in the system |
| totmass | total weight of all the molecules in the system |
| nmol | total number of separate molecules in the system |
| kmol | contiguous list of the atoms in each molecule |
| imol | first and last atom of each molecule in the list |
| molecule | number of the molecule to which each atom belongs |
| MOLDYN | velocity and acceleration on MD trajectory |
| v | current velocity of each atom along the x,y,z-axes |
| a | current acceleration of each atom along x,y,z-axes |
| aold | previous acceleration of each atom along x,y,z-axes |
| MOMENT | components of electric multipole moments |
| netchg | net electric charge for the total system |
| netdpl | dipole moment magnitude for the total system |
| netqdp | diagonal quadrupole (Qxx, Qyy, Qzz) for system |
| xdpl | dipole vector x-component in the global frame |
| ydpl | dipole vector y-component in the global frame |
| zdpl | dipole vector z-component in the global frame |
| xxqdp | quadrupole tensor xx-component in global frame |
| xyqdp | quadrupole tensor xy-component in global frame |
| xzqdp | quadrupole tensor xz-component in global frame |
| yxqdp | quadrupole tensor yx-component in global frame |
| yyqdp | quadrupole tensor yy-component in global frame |
| yzqdp | quadrupole tensor yz-component in global frame |
| zxqdp | quadrupole tensor zx-component in global frame |
| zyqdp | quadrupole tensor zy-component in global frame |
| zzqdp | quadrupole tensor zz-component in global frame |
| MPLPOT | specifics of atomic multipole functions |
| m2scale | factor by which 1-2 multipole interactions are scaled |
| m3scale | factor by which 1-3 multipole interactions are scaled |
| m4scale | factor by which 1-4 multipole interactions are scaled |
| m5scale | factor by which 1-5 multipole interactions are scaled |
| MPOLE | multipole components for current structure |
| maxpole | max components (monopole=1,dipole=4,quadrupole=13) |

| | |
|--------|---|
| pole | multipole values for each site in the local frame |
| rpole | multipoles rotated to the global coordinate system |
| npole | total number of multipole sites in the system |
| ipole | number of the atom for each multipole site |
| polsiz | number of multipole components at each multipole site |
| zaxis | number of the z-axis defining atom for each site |
| xaxis | number of the x-axis defining atom for each site |
| polaxe | local axis type for each multipole site |

MUTANT

hybrid atoms for free energy perturbation

| | |
|---------|---|
| lambda | weighting of initial state in hybrid Hamiltonian |
| nhybrid | number of atoms mutated from initial to final state |
| ihybrid | atomic sites differing in initial and final state |
| type0 | atom type of each atom in the initial state system |
| class0 | atom class of each atom in the initial state system |
| type1 | atom type of each atom in the final state system |
| class1 | atom class of each atom in the final state system |
| alter | true if an atom is to be mutated, false otherwise |

NUCLEO

parameters for nucleic acid structure

| | |
|---------|--|
| bkbone | phosphate backbone angles for each nucleotide |
| glyco | glycosidic torsional angle for each nucleotide |
| pucker | sugar pucker, either 2=2'-endo or 3=3'-endo |
| dblhlx | flag to mark system as nucleic acid double helix |
| deoxy | flag to mark deoxyribose or ribose sugar units |
| hlxform | helix form (A, B or Z) of polynucleotide strands |

OMEGA

dihedrals for torsional space computations

| | |
|--------|---|
| dihed | current value in radians of each dihedral angle |
| nomega | number of dihedral angles allowed to rotate |
| iomega | numbers of two atoms defining rotation axis |
| zline | line number in Z-matrix of each dihedral angle |

OPBEND

out-of-plane bends in the current structure

| | |
|---------|--|
| kopb | force constant values for out-of-plane bending |
| nopbend | total number of out-of-plane bends in the system |
| iopb | bond angle numbers used in out-of-plane bending |

OPDIST

out-of-plane distances in current structure

| | |
|---------|--|
| kopd | force constant values for out-of-plane distance |
| nopdist | total number of out-of-plane distances in the system |
| iopb | numbers of the atoms in each out-of-plane distance |

ORBITS

orbital energies for conjugated pisystem

| | |
|---|---|
| q | number of pi-electrons contributed by each atom |
| w | ionization potential of each pisystem atom |

| | |
|---------------|--|
| em | repulsion integral for each pisystem atom |
| nfill | number of filled pisystem molecular orbitals |
| OUTPUT | control of coordinate output file format |
| archive | logical flag to save structures in an archive |
| noverion | logical flag governing use of filename versions |
| overwrite | logical flag to overwrite intermediate files inplace |
| cyclesave | logical flag to mark use of numbered cycle files |
| coordtype | selects Cartesian, internal, rigid body or none |
| PATHS | parameters for Elber reaction path method |
| p0 | reactant Cartesian coordinates as variables |
| p1 | product Cartesian coordinates as variables |
| pmid | midpoint between the reactant and product |
| pvect | vector connecting the reactant and product |
| pstep | step per cycle along reactant-product vector |
| pzet | current projection on reactant-product vector |
| pnorm | length of the reactant-product vector |
| acoeff | transformation matrix 'A' from Elber paper |
| gc | gradients of the path constraints |
| PDB | definition of a Protein Data Bank structure |
| xpdb | x-coordinate of each atom stored in PDB format |
| ypdb | y-coordinate of each atom stored in PDB format |
| zpdb | z-coordinate of each atom stored in PDB format |
| npdb | number of atoms stored in Protein Data Bank format |
| resnum | number of the residue to which each atom belongs |
| npdb12 | number of atoms directly bonded to each CONECT atom |
| ipdb12 | atom numbers of atoms connected to each CONECT atom |
| pdblist | list of the Protein Data Bank atom number of each atom |
| pdbtyp | Protein Data Bank record type assigned to each atom |
| atmnam | Protein Data Bank atom name assigned to each atom |
| resnam | Protein Data Bank residue name assigned to each atom |
| PHIPSI | phi-psi-omega-chi angles for a protein |
| phi | value of the phi angle for each amino acid residue |
| psi | value of the psi angle for each amino acid residue |
| omega | value of the omega angle for each amino acid residue |
| chi | values of the chi angles for each amino acid residue |
| chiral | chirality of each amino acid residue (1=L, -1=D) |
| disulf | residue joined to each residue via a disulfide link |
| PIORBS | conjugated system in the current structure |
| norbit | total number of pisystem orbitals in the system |
| iorbit | numbers of the atoms containing pisystem orbitals |
| reorbit | number of evaluations between orbital updates |
| piperp | atoms defining a normal plane to each orbital |

| | |
|---------|---|
| npibond | total number of bonds affected by the pisystem |
| pibond | bond and pibond numbers for each pisystem bond |
| npitors | total number of torsions affected by the pisystem |
| pitors | torsion and pibond numbers for each pisystem torsion |
| listpi | atom list indicating whether each atom has an orbital |

PISTUF

bonds and torsions in the current pisystem

| | |
|--------|--|
| bkpi | bond stretch force constants for pi-bond order of 1.0 |
| blpi | ideal bond length values for a pi-bond order of 1.0 |
| kslope | rate of force constant decrease with bond order decrease |
| lslope | rate of bond length increase with a bond order decrease |
| torspi | 2-fold torsional energy barrier for pi-bond order of 1.0 |

PME

parameters for particle mesh Ewald summation

| | |
|----------|---|
| maxfft | maximum number of points along each FFT direction |
| maxorder | maximum order of the B-spline approximation |
| maxtable | maximum size of the FFT table array |
| maxgrid | maximum dimension of the PME charge grid array |
| bsmod1 | B-spline moduli along the a-axis direction |
| bsmod2 | B-spline moduli along the b-axis direction |
| bsmod3 | B-spline moduli along the c-axis direction |
| table | intermediate array used by the FFT calculation |
| nfft1 | number of grid points along the a-axis direction |
| nfft2 | number of grid points along the b-axis direction |
| nfft3 | number of grid points along the c-axis direction |
| bsorder | order of the PME B-spline approximation |

POLAR

polarizabilities and induced dipole moments

| | |
|----------|--|
| polarity | dipole polarizability for each multipole site (Ang**3) |
| pdamp | value of polarizability damping factor for each site |
| uind | induced dipole components at each multipole site |
| uinp | induced dipoles in field used for energy interactions |
| npolar | total number of polarizable sites in the system |

POLGRP

polarizable site group connectivity lists

| | |
|--------|--|
| maxp11 | maximum number of atoms in a polarization group |
| maxp12 | maximum number of atoms in groups 1-2 to an atom |
| maxp13 | maximum number of atoms in groups 1-3 to an atom |
| maxp14 | maximum number of atoms in groups 1-4 to an atom |
| np11 | number of atoms in polarization group of each atom |
| ip11 | atom numbers of atoms in same group as each atom |
| np12 | number of atoms in groups 1-2 to each atom |
| ip12 | atom numbers of atoms in groups 1-2 to each atom |
| np13 | number of atoms in groups 1-3 to each atom |
| ip13 | atom numbers of atoms in groups 1-3 to each atom |
| np14 | number of atoms in groups 1-4 to each atom |
| ip14 | atom numbers of atoms in groups 1-4 to each atom |

POLPOT

poleps
polsor
pgamma
p1scale
p2scale
p3scale
p4scale
d1scale
d2scale
d3scale
d4scale
u1scale
u2scale
u3scale
u4scale
polold
poltyp

specifics of polarization functional form

induced dipole convergence criterion (rms Debyes/atom)
induced dipole SOR convergence acceleration factor
prefactor in exponential polarization damping term
field intra-group scale factor for energy evaluations
field 1-2 group scale factor for energy evaluations
field 1-3 group scale factor for energy evaluations
field 1-4 group scale factor for energy evaluations
field intra-group scale factor for direct induced
field 1-2 group scale factor for direct induced
field 1-3 group scale factor for direct induced
field 1-4 group scale factor for direct induced
field intra-group scale factor for mutual induced
field 1-2 group scale factor for mutual induced
field 1-3 group scale factor for mutual induced
field 1-4 group scale factor for mutual induced
logical flag for scale factor instead of Thole damping
type of polarization potential (direct or mutual)

POTENT

use_bond
use_angle
use_strbnd
use_urey
use_angang
use_opbnd
use_opdist
use_improp
use_imptor
use_tors
use_strtor
use_tortor
use_vdw
use_charge
use_chgdpl
use_dipole
use_mpole
use_polar
use_rxnflld
use_solv
use_gbsa
use_metal
use_geom
use_extra
use_orbit

usage of each potential energy component

logical flag governing use of bond stretch potential
logical flag governing use of angle bend potential
logical flag governing use of stretch-bend potential
logical flag governing use of Urey-Bradley potential
logical flag governing use of angle-angle cross term
logical flag governing use of out-of-plane bend term
logical flag governing use of out-of-plane distance
logical flag governing use of improper dihedral term
logical flag governing use of improper torsion term
logical flag governing use of torsional potential
logical flag governing use of stretch-torsion term
logical flag governing use of torsion-torsion term
logical flag governing use of vdw der Waals potential
logical flag governing use of charge-charge potential
logical flag governing use of charge-dipole potential
logical flag governing use of dipole-dipole potential
logical flag governing use of multipole potential
logical flag governing use of polarization term
logical flag governing use of reaction field term
logical flag governing use of surface area solvation
logical flag governing use of GB/SA solvation term
logical flag governing use of ligand field term
logical flag governing use of geometric restraints
logical flag governing use of extra potential term
logical flag governing use of pssystem computation

PRECIS

tiny
small

values of machine precision tolerances

the smallest positive floating point value
the smallest relative floating point spacing

| | |
|---------------|---|
| huge | the largest relative floating point spacing |
| REFER | storage of reference atomic coordinate set |
| xref | reference x-coordinate for each atom in the system |
| yref | reference y-coordinate for each atom in the system |
| zref | reference z-coordinate for each atom in the system |
| nref | total number of atoms in the reference system |
| reftyp | atom type for each atom in the reference system |
| n12ref | number of atoms bonded to each reference atom |
| i12ref | atom numbers of atoms 1-2 connected to each atom |
| refleng | length in characters of the reference filename |
| refltile | length in characters of the reference title string |
| refnam | atom name for each atom in the reference system |
| reffile | base filename for the reference structure |
| reftitle | title used to describe the reference structure |
| RESDUE | standard biopolymer residue abbreviations |
| amino | three-letter abbreviations for amino acids types |
| nuclz | three-letter abbreviations for nucleic acids types |
| amino1 | one-letter abbreviations for amino acids types |
| nuclz1 | one-letter abbreviations for nucleic acids types |
| RGDDYN | velocities and momenta for rigid-body MD |
| vcm | current translational velocity of each rigid-body |
| wcm | current angular velocity of each rigid-body |
| lm | current angular momenta of each rigid-body |
| linear | logical flag to mark group as linear or nonlinear |
| RIGID | rigid body coordinates for atom groups |
| xrb | rigid body reference x-coordinate for each atom |
| yrb | rigid body reference y-coordinate for each atom |
| zrb | rigid body reference z-coordinate for each atom |
| rbc | current rigid body coordinates for each atom group |
| RING | number and location of small ring structures |
| nring3 | total number of 3-membered rings in the system |
| iring3 | numbers of the atoms involved in each 3-ring |
| nring4 | total number of 4-membered rings in the system |
| iring4 | numbers of the atoms involved in each 4-ring |
| nring5 | total number of 5-membered rings in the system |
| iring5 | numbers of the atoms involved in each 5-ring |
| nring6 | total number of 6-membered rings in the system |
| iring6 | numbers of the atoms involved in each 6-ring |
| ROTATE | molecule partitions for rotation of a bond |
| nrot | total number of atoms moving when bond rotates |

| | |
|---------------|--|
| rot | atom numbers of atoms moving when bond rotates |
| use_short | logical flag governing use of shortest atom list |
| RXNFLD | reaction field matrix elements and indices |
| b1 | |
| b2 | |
| ijk | |
| RXNPOT | specifics of reaction field functional form |
| rfsiz | radius of reaction field sphere centered at origin |
| rfbulkd | bulk dielectric constant of reaction field continuum |
| rfterms | number of terms to use in reaction field summation |
| SCALES | parameter scale factors for optimization |
| scale | multiplicative factor for each optimization parameter |
| set_scale | logical flag to show if scale factors have been set |
| SEQUEN | sequence information for a biopolymer |
| nseq | total number of residues in biopolymer sequences |
| nchain | number of separate biopolymer sequence chains |
| ichain | first and last residue in each biopolymer chain |
| seqtyp | residue type for each residue in the sequence |
| seq | three-letter code for each residue in the sequence |
| chnnam | one-letter identifier for each sequence chain |
| SHAKE | definition of Shake/Rattle constraints |
| krat | ideal distance value for rattle constraint |
| nrat | number of rattle constraints to be applied |
| irat | atom numbers of atoms in a rattle constraint |
| ratimage | flag to use minimum image for rattle constraint |
| use_rattle | logical flag to set use of rattle constraints |
| SHUNT | polynomial switching function coefficients |
| off | distance at which the potential energy goes to zero |
| off2 | square of distance at which the potential goes to zero |
| cut | distance at which switching of the potential begins |
| cut2 | square of distance at which the switching begins |
| c0 | zeroth order coefficient of multiplicative switch |
| c1 | first order coefficient of multiplicative switch |
| c2 | second order coefficient of multiplicative switch |
| c3 | third order coefficient of multiplicative switch |
| c4 | fourth order coefficient of multiplicative switch |
| c5 | fifth order coefficient of multiplicative switch |
| f0 | zeroth order coefficient of additive switch function |
| f1 | first order coefficient of additive switch function |
| f2 | second order coefficient of additive switch function |

| | |
|----|---|
| f3 | third order coefficient of additive switch function |
| f4 | fourth order coefficient of additive switch function |
| f5 | fifth order coefficient of additive switch function |
| f6 | sixth order coefficient of additive switch function |
| f7 | seventh order coefficient of additive switch function |

SIZES

parameter values to set array dimensions

"sizes.i" sets values for critical array dimensions used throughout the software; these parameters will fix the size of the largest systems that can be handled; values too large for the computer's memory and/or swap space to accomodate will result in poor performance or outright failure

| parameter: | maximum allowed number of: |
|------------|---|
| maxatm | atoms in the molecular system |
| maxval | atoms directly bonded to an atom |
| maxgrp | user-defined groups of atoms |
| maxtyp | force field atom type definitions |
| maxclass | force field atom class definitions |
| maxkey | lines in the keyword file |
| maxrot | bonds for torsional rotation |
| maxvar | optimization variables (vector storage) |
| maxopt | optimization variables (matrix storage) |
| maxhess | off-diagonal Hessian elements |
| maxlight | sites for method of lights neighbors |
| maxvib | vibrational frequencies |
| maxgeo | distance geometry points |
| maxcell | unit cells in replicated crystal |
| maxring | 3-, 4-, or 5-membered rings |
| maxfix | geometric restraints |
| maxbio | biopolymer atom definitions |
| maxres | residues in the macromolecule |
| maxamino | amino acid residue types |
| maxnuc | nucleic acid residue types |
| maxbnd | covalent bonds in molecular system |
| maxang | bond angles in molecular system |
| maxtors | torsional angles in molecular system |
| maxpi | atoms in conjugated pisystem |
| maxpib | covalent bonds involving pisystem |
| maxpit | torsional angles involving pisystem |

SOLUTE

parameters for continuum solvation models

| | |
|---------|---|
| rsolv | atomic radius of each atom for continuum solvation |
| vsolv | atomic volume of each atom for continuum solvation |
| asolv | atomic solvation parameters (kcal/mole/Ang**2) |
| rborn | Born radius of each atom for GB/SA solvation |
| drb | solvation derivatives with respect to Born radii |
| doffset | dielectric offset to continuum solvation atomic radii |
| p1 | single-atom scale factor for analytical Still GB/SA |
| p2 | 1-2 interaction scale factor for analytical Still GB/SA |

| | |
|---------|--|
| p3 | 1-3 interaction scale factor for analytical Still GB/SA |
| p4 | nonbonded scale factor for analytical Still GB/SA |
| p5 | soft cutoff parameter for analytical Still GB/SA |
| gpol | polarization self-energy values for each atom |
| shct | overlap scaling factors for Hawkins-Cramer-Truhlar GB/SA |
| wace | "omega" values for atom class pairs for use with ACE |
| s2ace | "sigma^2" values for atom class pairs for use with ACE |
| uace | "mu" values for atom class pairs for use with ACE |
| solvtyp | solvation model (ASP, SASA, ONION, STILL, HCT, ACE) |

STODYN

frictional coefficients for SD trajectory

| | |
|------------|--|
| friction | global frictional coefficient for exposed particle |
| gamma | atomic frictional coefficients for each atom |
| use_sdarea | logical flag to use surface area friction scaling |

STRBND

stretch-bends in the current structure

| | |
|---------|---|
| ksb | force constant for stretch-bend terms |
| nstrbnd | total number of stretch-bend interactions |
| isb | angle and bond numbers used in stretch-bend |

STRTOR

stretch-torsions in the current structure

| | |
|---------|---|
| kst | 1-, 2- and 3-fold stretch-torsion force constants |
| nstrtor | total number of stretch-torsion interactions |
| ist | torsion and bond numbers used in stretch-torsion |

SYNTRN

definition of synchronous transit path

| | |
|-------|---|
| t | value of the path coordinate (0=reactant, 1=product) |
| pm | path coordinate for extra point in quadratic transit |
| xmin1 | reactant coordinates as array of optimization variables |
| xmin2 | product coordinates as array of optimization variables |
| xm | extra coordinate set for quadratic synchronous transit |

TITLES

title for the current molecular system

| | |
|--------|---|
| ltitle | length in characters of the nonblank title string |
| title | title used to describe the current structure |

TORPOT

specifics of torsional functional forms

| | |
|----------|---|
| idihunit | convert improper dihedral force to kcal/mole/deg**2 |
| itorunit | convert improper torsion amplitudes to kcal/mole |
| torsunit | convert torsional parameter amplitudes to kcal/mole |
| storunit | convert stretch-torsion force to kcal/mole/Ang |

TORS

torsional angles within the current structure

| | |
|-------|---|
| tors1 | 1-fold amplitude and phase for each torsional angle |
| tors2 | 2-fold amplitude and phase for each torsional angle |

| | |
|---------------|--|
| tors3 | 3-fold amplitude and phase for each torsional angle |
| tors4 | 4-fold amplitude and phase for each torsional angle |
| tors5 | 5-fold amplitude and phase for each torsional angle |
| tors6 | 6-fold amplitude and phase for each torsional angle |
| ntors | total number of torsional angles in the system |
| itors | numbers of the atoms in each torsional angle |
| TREE | potential smoothing & search tree levels |
| maxpss | maximum number of potential smoothing levels |
| etree | energy reference value at the top of the tree |
| ilevel | smoothing deformation value at each tree level |
| nlevel | number of levels of potential smoothing used |
| UNITS | physical constants and unit conversions |
| avogadro | Avogadro's number (N) in particles/mole |
| boltzmann | Boltzmann constant (kB) in $\text{g}^*\text{Ang}^{**2}/\text{ps}^{**2}/\text{K}/\text{mole}$ |
| gasconst | ideal gas constant (R) in kcal/mole/K |
| lightspd | speed of light in vacuum (c) in cm/ps |
| bohr | conversion from Bohrs to Angstroms |
| joule | conversion from calories to joules |
| evolt | conversion from Hartree to electron-volts |
| hartree | conversion from Hartree to kcal/mole |
| electric | conversion from $\text{electron}^{**2}/\text{Ang}$ to kcal/mole |
| debye | conversion from electron-Ang to Debyes |
| prescon | conversion from kcal/mole/Ang ³ to Atm |
| convert | conversion from kcal to $\text{g}^*\text{Ang}^{**2}/\text{ps}^{**2}$ |
| UREY | Urey-Bradley interactions in the structure |
| uk | Urey-Bradley force constants (kcal/mole/Ang ²) |
| ul | ideal 1-3 distance values in Angstroms |
| nurey | total number of Urey-Bradley terms in the system |
| iury | numbers of the atoms in each Urey-Bradley interaction |
| URYPOT | specifics of Urey-Bradley functional form |
| cury | cubic coefficient in Urey-Bradley potential |
| qury | quartic coefficient in Urey-Bradley potential |
| ureyunit | convert Urey-Bradley constant to kcal/mole/Ang ² |
| USAGE | atoms active during energy computation |
| nuse | number of active atoms used in energy calculation |
| use | true if an atom is active, false if inactive |
| VDW | van der Waals parameters for current structure |
| radmin | minimum energy distance for each atom class pair |
| epsilon | well depth parameter for each atom class pair |
| radmin4 | minimum energy distance for 1-4 interaction pairs |

| | |
|----------|---|
| epsilon4 | well depth parameter for 1-4 interaction pairs |
| radhbnd | minimum energy distance for hydrogen bonding pairs |
| epshbnd | well depth parameter for hydrogen bonding pairs |
| kred | value of reduction factor parameter for each atom |
| ired | attached atom from which reduction factor is applied |
| nvdw | total number van der Waals active sites in the system |
| ivdw | number of the atom for each van der Waals active site |

VDWPOT

| | |
|----------|--|
| abuck | value of "A" constant in Buckingham vdw potential |
| bbuck | value of "B" constant in Buckingham vdw potential |
| cbuck | value of "C" constant in Buckingham vdw potential |
| ghal | value of "gamma" in buffered 14-7 vdw potential |
| dhal | value of "delta" in buffered 14-7 vdw potential |
| v2scale | factor by which 1-2 vdw interactions are scaled |
| v3scale | factor by which 1-3 vdw interactions are scaled |
| v4scale | factor by which 1-4 vdw interactions are scaled |
| v5scale | factor by which 1-5 vdw interactions are scaled |
| igauss | coefficients of Gaussian fit to vdw potential |
| ngauss | number of Gaussians used in fit to vdw potential |
| vdwtyp | type of van der Waals potential energy function |
| radtyp | type of parameter (sigma or R-min) for atomic size |
| radsiz | atomic size provided as radius or diameter |
| radrule | combining rule for atomic size parameters |
| epsrule | combining rule for vdw well depth parameters |
| gausstyp | type of Gaussian fit to van der Waals potential |

VIRIAL

| | |
|------|--|
| virx | x-component of the total internal virial |
| viry | y-component of the total internal virial |
| virz | z-component of the total internal virial |

WARP

| | |
|------------|--|
| m2 | second moment of Gaussian representing each atom |
| deform | value of diffusional smoothing deformation parameter |
| diff1 | diffusion coefficient for torsional potential |
| diffv | diffusion coefficient for van der Waals potential |
| diffc | diffusion coefficient for charge-charge potential |
| use_deform | flag to use diffusion smoothed potential terms |
| use_gda | flag to use Straub's GDA instead of Scheraga's DEM |

XTALS

| | |
|------------|---|
| e0_lattice | ideal lattice energy for the current crystal |
| moment_0 | ideal dipole moment for monomer from crystal |
| nxtal | number of crystal structures to be stored |
| nvary | number of potential parameters to optimize |
| ivary | index for the types of potential parameters |
| vary | atom numbers involved in potential parameters |

specifics of van der Waals functional form

components of the internal virial

parameters for potential surface smoothing

crystal structures for parameter fitting

| | |
|---------------|--|
| iresid | crystal structure to which each residual refers |
| rsdtyp | experimental variable for each of the residuals |
| vartyp | type of potential parameter to be optimized |
| ZCLOSE | ring openings and closures for Z-matrix |
| nadd | number of added bonds between Z-matrix atoms |
| iadd | numbers of the atom pairs defining added bonds |
| ndel | number of bonds between Z-matrix atoms to delete |
| idel | numbers of the atom pairs defining deleted bonds |
| ZCOORD | Z-matrix internal coordinate definitions |
| zbond | bond length used to define each Z-matrix atom |
| zang | bond angle used to define each Z-matrix atom |
| ztors | angle or torsion used to define Z-matrix atom |
| iz | defining atom numbers for each Z-matrix atom |

11. Index of Function & Subroutine Calls

This section contains an alphabetical cross index listing of the routines called by each TINKER program, subroutine and function. Routines not present in the left hand column do not make calls to any other portion of the TINKER package.

| <u>Routine</u> | <u>List of Source Code Units called by this Routine</u> | | | | |
|-----------------------|--|--|--|---|---|
| ACTIVE | GETTEXT | UPCASE | | | |
| ADDBASE | ADDBOND OVERLAP | FINDATM PIALTER | JACOBI PIMOVE | NEWATM PITILT | OLDATM |
| ADDSIDE | ADDBASE JACOBI PIMOVE | ADDBOND NEWATM PITILT | FATAL OLDATM PRTSEQ | FINDATM OVERLAP VERSION | FREEUNIT PIALTER |
| ALCHEMY | ENERGY HATOM READYXYZ | FINAL HYBRID UPCASE | FREEUNIT INITIAL VERSION | GETTEXT MECHANIC | GETXYZ NUMERAL |
| ANALYSIS | BOUNDS ECHARGE3 EHAL3 EMM3HB3 ESOLV3 EXTRA3 | EANGANG3 ECHGDPL3 EIMPROP3 EMPOLE3 ESTRBND3 PISCF | EANGLE3 EDIPOLE3 EIMPTOR3 EOPBEND3 ESTRTOR3 REPLICA | EBOND3 EGAUSS3 ELJ3 EOPDIST3 ETORS3 | EBUCK3 EGEOM3 EMETAL3 ERXNFLD3 EUREY3 |
| ANALYZE | ANALYZ4 FINAL NEXTARG TRIMTEXT | ANALYZ6 FREEUNIT PARAMYZE UPCASE | ANALYZ8 GETXYZ PROPYZE VERSION | ATOMYZE INITIAL READYXYZ | ENRGYZE MECHANIC SUFFIX |
| ANGLES | FATAL | | | | |
| ANNEAL | BEEMAN MDINIT SHAKEUP | FINAL MDREST SIGMOID | GETTEXT MECHANIC UPCASE | GETXYZ NEXTARG VERLET | INITIAL SDSTEP |
| ARCHIVE | ACTIVE INITIAL PRTXYZ VERSION | BASEFILE NEXTARG READYXYZ | FINAL NUMERAL SUFFIX | FREEUNIT PRTMSI TRIMTEXT | GETTEXT PRTXMOL UPCASE |
| ATTACH | SORT | | | | |
| BASEFILE | CONTROL | GETKEY | TRIMTEXT | | |
| BEEMAN | GRADIENT RATTLE2 | MDSAVE TEMPER | MDSTAT | PRESSURE | RATTLE |

| | | | | | |
|----------|--------------------|-----------------|----------|----------|---------|
| BETAI | BETACF | GAMMLN | | | |
| BIGBLOCK | CELLATOM | | | | |
| BONDS | FATAL | | | | |
| BORN | SURFATOM | | | | |
| BSET | BMAX | | | | |
| BSSTEP | FATAL | MMID | PZEXTR | | |
| CALENDAR | IDATE | ITIME | | | |
| CERROR | FATAL | TRIMTEXT | | | |
| CFFTB | CFFTB1 | | | | |
| CFFTB1 | PASSB | PASSB2 | PASSB3 | PASSB4 | PASSB5 |
| CFFTF | CFFTF1 | | | | |
| CFFTF1 | PASSF | PASSF2 | PASSF3 | PASSF4 | PASSF5 |
| CFFTI | CFFTI1 | | | | |
| CHKTREE | LOCALXYZ | | | | |
| CIRPLN | ANORM | DOT | VCROSS | VNORM | |
| CLIMBER | ENERGY | GETREF | LOCALMIN | MAKEINT | MAKEXYZ |
| CLIMBRGD | ENERGY | LOCALRGD | RIGIDXYZ | | |
| CLIMBROT | ENERGY | LOCALROT | MAKEXYZ | | |
| CLIMBTOR | CHKTREE MAKEXYZ | ENERGY | GETREF | LOCALXYZ | MAKEINT |
| CLIMBXYZ | CHKTREE | ENERGY | GETREF | LOCALXYZ | |
| CLUSTER | CUTOFFS SORT3 | FATAL UPCASE | GETNUMB | GETTEXT | SORT |
| COMMAND | GETARG | UPCASE | | | |
| COMPRESS | CERROR | GETTOR | | | |
| CONNECT | SORT | | | | |

| | | | | | |
|-----------|--|--|--|--|---|
| CONNOLLY | COMPRESS TORUS | CONTACT VAM | NEIGHBOR | PLACE | SADDLES |
| CONTACT | ANORM | CERROR | PTINCY | | |
| CONTROL | GETTEXT | UPCASE | | | |
| COORDS | GYRATE | RMSERROR | | | |
| CORRELATE | FINAL TRIMTEXT | INITIAL | NEXTARG | PROPERTY | READBLK |
| CRYSTAL | BIGBLOCK GETTEXT MOLECULE UPCASE | BOUNDS GETXYZ NEXTARG VERSION | FIELD INITIAL PRTXYZ | FINAL KATOM SYMMETRY | FREEUNIT LATTICE UNITCELL |
| CUTOFFS | GETTEXT | UPCASE | | | |
| DEPTH | DOT | VCROSS | VNORM | | |
| DIAGQ | GETIME | SETIME | | | |
| DIFFEQ | BSSTEP | DERIVS | GDASTAT | | |
| DIFFUSE | BASEFILE GETWORD READXYZ | FATAL INITIAL SUFFIX | FIELD KATOM UNITCELL | FINAL MOLECULE VERSION | FREEUNIT NEXTARG |
| DISTGEOM | ACTIVE FATAL GETTEXT KCHIRAL PRTXYZ VERSION | ANGLES FINAL GETXYZ KGEOM SETIME | ATTACH FREEUNIT GRAFIC MAKERE TORSIONS | BONDS GEODESIC IMPOSE NEXTARG TRIFIX | EMBED GETIME INITIAL NUMERAL UPCASE |
| DMDUMP | GRAFIC | | | | |
| DOCUMENT | FINAL INITIAL SORT6 UPCASE | FREEUNIT LOWCASE SORT7 VERSION | GETPRM NEXTARG SORT9 | GETTEXT NEXTTEXT SUFFIX | GETWORD PRTPRM TRIMTEXT |
| DSTMAT | GETIME RANDOM | GETNUMB SETIME | GETTEXT SORT2 | INVBETA TRIFIX | LOWCASE UPCASE |
| DYNAMIC | BEEMAN MDREST SHAKEUP | FINAL MECHANIC VERLET | GETXYZ NEXTARG | INITIAL RGDSTEP | MDINIT SDSTEP |
| EANGANG | GROUPS | IMAGE | | | |

| | | | |
|-----------|-----------|-----------|---------------------|
| EANGANG1 | GROUPS | IMAGE | |
| EANGANG2 | EANGANG2A | GROUP | |
| EANGANG2A | IMAGE | | |
| EANGANG3 | GROUPS | IMAGE | |
| EANGLE | GROUPS | IMAGE | |
| EANGLE1 | GROUPS | IMAGE | |
| EANGLE2 | EANGLE2A | EANGLE2B | GROUPS |
| EANGLE2A | GROUPS | IMAGE | |
| EANGLE2B | IMAGE | | |
| EANGLE3 | GROUPS | IMAGE | |
| EBOND | GROUPS | IMAGE | |
| EBOND1 | GROUPS | IMAGE | |
| EBOND2 | GROUPS | IMAGE | |
| EBOND3 | GROUPS | IMAGE | |
| EBUCK | EBUCK0A | EBUCK0B | |
| EBUCK0A | GROUPS | IMAGE | SWITCH |
| EBUCK0B | GROUPS | LIGHTS | SWITCH |
| EBUCK1 | EBUCK1A | EBUCK1B | |
| EBUCK1A | GROUPS | IMAGE | SWITCH |
| EBUCK1B | GROUPS | LIGHTS | SWITCH |
| EBUCK2 | GROUPS | IMAGE | SWITCH |
| EBUCK3 | EBUCK3A | EBUCK3B | |
| EBUCK3A | GROUPS | IMAGE | SWITCH |
| EBUCK3B | GROUPS | LIGHTS | SWITCH |
| ECHARGE | ECHARGE0A | ECHARGE0B | ECHARGE0C ECHARGE0D |

| | | | | | |
|------------------|-----------|-----------|-----------|-----------|--------|
| ECHARGE0A | GROUPS | IMAGE | SWITCH | | |
| ECHARGE0B | GROUPS | LIGHTS | SWITCH | | |
| ECHARGE0C | ERF | GROUPS | | | |
| ECHARGE0D | EPME | ERFC | GROUPS | IMAGE | SWITCH |
| ECHARGE1 | ECHARGE1A | ECHARGE1B | ECHARGE1C | ECHARGE1D | |
| ECHARGE1A | GROUPS | IMAGE | SWITCH | | |
| ECHARGE1B | GROUPS | LIGHTS | SWITCH | | |
| ECHARGE1C | ERF | GROUPS | | | |
| ECHARGE1D | EPME1 | ERFC | GROUPS | IMAGE | SWITCH |
| ECHARGE2 | ECHARGE2A | ECHARGE2B | ECHARGE2C | | |
| ECHARGE2A | GROUPS | IMAGE | SWITCH | | |
| ECHARGE2B | ERF | GROUPS | | | |
| ECHARGE2C | ERFC | GROUPS | IMAGE | | |
| ECHARGE3 | ECHARGE3A | ECHARGE3B | ECHARGE3C | ECHARGE3D | |
| ECHARGE3A | GROUPS | IMAGE | SWITCH | | |
| ECHARGE3B | GROUPS | LIGHTS | SWITCH | | |
| ECHARGE3C | ERF | GROUPS | | | |
| ECHARGE3D | EPME3 | ERFC | GROUPS | IMAGE | SWITCH |
| ECHGDPL | GROUPS | IMAGE | SWITCH | | |
| ECHGDPL1 | GROUPS | IMAGE | SWITCH | | |
| ECHGDPL2 | GROUPS | IMAGE | SWITCH | | |
| ECHGDPL3 | GROUPS | IMAGE | SWITCH | | |
| EDIPOLE | GROUPS | IMAGE | SWITCH | | |
| EDIPOLE1 | GROUPS | IMAGE | SWITCH | | |
| EDIPOLE2 | GROUPS | IMAGE | SWITCH | | |
| EDIPOLE3 | GROUPS | IMAGE | SWITCH | | |

| | | | |
|----------------|--------|--------|--------|
| EGAUSS | GROUPS | SWITCH | |
| EGAUSS1 | GROUPS | SWITCH | |
| EGAUSS2 | GROUPS | SWITCH | |
| EGAUSS3 | GROUPS | SWITCH | |
| EGBSA0A | GROUPS | SWITCH | |
| EGBSA0B | ERF | GROUPS | |
| EGBSA1A | GROUPS | SWITCH | |
| EGBSA1B | ERF | GROUPS | |
| EGBSA2A | SWITCH | | |
| EGBSA2B | ERF | | |
| EGBSA3A | GROUPS | SWITCH | |
| EGBSA3B | ERF | GROUPS | |
| EGEOM | GROUPS | IMAGE | |
| EGEOM1 | GROUPS | IMAGE | |
| EGEOM2 | GROUPS | IMAGE | |
| EGEOM3 | GROUPS | IMAGE | |
| EHAL | EHAL0A | EHAL0B | |
| EHAL0A | GROUPS | IMAGE | SWITCH |
| EHAL0B | GROUPS | LIGHTS | SWITCH |
| EHAL1 | EHAL1A | EHAL1B | |
| EHAL1A | GROUPS | IMAGE | SWITCH |
| EHAL1B | GROUPS | LIGHTS | SWITCH |
| EHAL2 | GROUPS | IMAGE | SWITCH |
| EHAL3 | EHAL3A | EHAL3B | |
| EHAL3A | GROUPS | IMAGE | SWITCH |

| | | | | | |
|-----------------|--------|---------|---------|--------|--------|
| EHAL3B | GROUPS | LIGHTS | SWITCH | | |
| EIGEN | GETIME | POWER | SETIME | | |
| EIGENRGD | DIAGQ | HESSRGD | | | |
| EIGENROT | DIAGQ | HESSROT | | | |
| EIGENROT | DIAGQ | HESSROT | | | |
| EIGENTOR | DIAGQ | HESSROT | | | |
| EIGENXYZ | DIAGQ | HESSIAN | | | |
| EIMPROP | GROUPS | IMAGE | | | |
| EIMPROP1 | GROUPS | IMAGE | | | |
| EIMPROP2 | GROUPS | IMAGE | | | |
| EIMPROP3 | GROUPS | IMAGE | | | |
| EIMPTOR | GROUPS | IMAGE | | | |
| EIMPTOR1 | GROUPS | IMAGE | | | |
| EIMPTOR2 | GROUPS | IMAGE | | | |
| EIMPTOR3 | GROUPS | IMAGE | | | |
| ELJ | ELJ0A | ELJ0B | | | |
| ELJ0A | GROUPS | IMAGE | SWITCH | | |
| ELJ0B | GROUPS | LIGHTS | SWITCH | | |
| ELJ1 | ELJ1A | ELJ1B | | | |
| ELJ1A | GROUPS | IMAGE | SWITCH | | |
| ELJ1B | GROUPS | LIGHTS | SWITCH | | |
| ELJ2 | GROUPS | IMAGE | SWITCH | | |
| ELJ3 | ELJ3A | ELJ3B | | | |
| ELJ3A | GROUPS | IMAGE | SWITCH | | |
| ELJ3B | GROUPS | LIGHTS | SWITCH | | |
| EMBED | BNDERR | CHIRER | CHKSIZE | COORDS | DMDUMP |

| | | | | | |
|-----------------|--------------------------------------|--------------------------------------|---------------------------------------|------------------------------|----------------------------------|
| | DSTMAT GETIME METRIC SETIME | EIGEN GYRATE NUMERAL TORSER | EXPLORE IMPOSE PRTXYZ VDWERR | FRACDIST LOCERR REFINE | FREEUNIT MAJORIZE RMSError |
| EMETAL | FATAL | | | | |
| EMETAL1 | FATAL | | | | |
| EMETAL3 | EMETAL | | | | |
| EMM3HB | EMM3HB0A | EMM3HB0B | | | |
| EMM3HB0A | GROUPS | IMAGE | SWITCH | | |
| EMM3HB0B | GROUPS | LIGHTS | SWITCH | | |
| EMM3HB1 | EMM3HB1A | EMM3HB1B | | | |
| EMM3HB1A | GROUPS | IMAGE | SWITCH | | |
| EMM3HB1B | GROUPS | LIGHTS | SWITCH | | |
| EMM3HB2 | GROUPS | IMAGE | SWITCH | | |
| EMM3HB3 | EMM3HB3A | EMM3HB3B | | | |
| EMM3HB3A | GROUPS | IMAGE | SWITCH | | |
| EMM3HB3B | GROUPS | LIGHTS | SWITCH | | |
| EMPOLE | EMPOLE0A | EMPOLE0B | | | |
| EMPOLE0A | GROUPS SWITCH | IMAGE | INDUCE | ROTMAT | ROTPOLE |
| EMPOLE0B | EREAL | ERECIP | INDUCE | ROTMAT | ROTPOLE |
| EMPOLE1 | EMPOLE1A | EMPOLE1B | | | |
| EMPOLE1A | GROUPS SWITCH | IMAGE TORQUE | INDUCE TORQUE1 | ROTMAT | ROTPOLE |
| EMPOLE1B | EREAL1 TORQUE | ERECIP1 | INDUCE | ROTMAT | ROTPOLE |
| EMPOLE2 | EMPOLE2A | | | | |
| EMPOLE2A | GROUPS TORQUE | IMAGE | ROTMAT | ROTPOLE | SWITCH |

| | | | | | |
|------------------|---|---|---|--|--|
| EMPOLE3 | EMPOLE3A | EMPOLE3B | | | |
| EMPOLE3A | GROUPS SWITCH | IMAGE | INDUCE | ROTMAT | ROTPOLE |
| EMPOLE3B | EREAL3 | ERECIP3 | INDUCE | ROTMAT | ROTPOLE |
| ENERGY | BOUNDS ECHARGE EHAL EMM3HB ESOLV EXTRA | EANGANG ECHGDPL EIMPROP EMPOLE ESTRBND PISCF | EANGLE EDIPOLE EIMPTOR EOPBEND ESTRTOR REPLICA | EBOND EGAUSS ELJ EOPDIST ETORS | EBUCK EGEOM EMETAL ERXNFLD EUREY |
| ENRGYZE | ANALYSIS | | | | |
| EOPBEND | GROUPS | IMAGE | | | |
| EOPBEND1 | GROUPS | IMAGE | | | |
| EOPBEND2 | EOPBEND2A | GROUPS | | | |
| EOPBEND2A | IMAGE | | | | |
| EOPBEND3 | GROUPS | IMAGE | | | |
| EOPDIST | GROUPS | IMAGE | | | |
| EOPDIST1 | GROUPS | IMAGE | | | |
| EOPDIST2 | GROUPS | IMAGE | | | |
| EOPDIST3 | GROUPS | IMAGE | | | |
| EPME | BSPLINE | FFTFRONT | | | |
| EPME1 | BSPLINE1 | FFTBACK | FFTFRONT | | |
| EPME3 | BSPLINE | FFTFRONT | | | |
| EPUCLC | ANORM | | | | |
| EREAL | ERFC | IMAGE | SWITCH | | |
| EREAL1 | ERFC | IMAGE | SWITCH | TORQUE | TORQUE1 |
| EREAL3 | ERFC | IMAGE | SWITCH | | |
| ERECIP1 | TORQUE | | | | |
| ERF | ERFCORE | | | | |

| | | | | | |
|-----------------|----------------|----------------|----------------|----------------|----------------|
| ERFC | ERFCORE | | | | |
| ERFIK | D1D2 | RFINDEX | | | |
| ERFINV | ERF | FATAL | | | |
| ERXNFLD | ERFIK | IJK_PT | ROTMAT | ROTPOLE | SWITCH |
| ERXNFLD3 | ERFIK | IJK_PT | ROTMAT | ROTPOLE | SWITCH |
| ESOLV | BORN | EGBSA0A | EGBSA0B | SURFACE | |
| ESOLV1 | BORN | BORN1 | EGBSA1A | EGBSA1B | SURFACE |
| ESOLV2 | EGBSA2A | EGBSA2B | | | |
| ESOLV3 | BORN | EGBSA3A | EGBSA3B | SURFACE | |
| ESTRBND | GROUPS | IMAGE | | | |
| ESTRBND1 | GROUPS | IMAGE | | | |
| ESTRBND2 | GROUPS | IMAGE | | | |
| ESTRBND3 | GROUPS | IMAGE | | | |
| ESTRTOR | GROUPS | IMAGE | | | |
| ESTRTOR1 | GROUPS | IMAGE | | | |
| ESTRTOR2 | GROUPS | IMAGE | | | |
| ESTRTOR3 | GROUPS | IMAGE | | | |
| ETORS | GROUPS | IMAGE | | | |
| ETORS1 | GROUPS | IMAGE | | | |
| ETORS2 | GROUPS | IMAGE | | | |
| ETORS3 | GROUPS | IMAGE | | | |
| EUREY | GROUPS | IMAGE | | | |
| EUREY1 | GROUPS | IMAGE | | | |
| EUREY2 | GROUPS | IMAGE | | | |
| EUREY3 | GROUPS | IMAGE | | | |

| | | | | | |
|-----------------|----------------------------|----------------------------|------------------------------|------------------------------|-------------------------------|
| EWALDCOF | ERFC | | | | |
| EXPLORE | INITERR | MIDERR | SIGMOID | TOTERR | |
| FFTBACK | CFFTB | | | | |
| FFTFRONT | CFFTF | | | | |
| FFTSETUP | CFFTI | | | | |
| FIELD | GETPRM | PRMKEY | | | |
| FRACDIST | DIST2 | TRIMTEXT | | | |
| FREEUNIT | FATAL | | | | |
| GDA | DIFFEQ GETXYZ PRTXYZ | FINAL INITIAL RANDOM | FREEUNIT MECHANIC TNCG | GDASTAT NEXTARG UPCASE | GETTEXT NUMERAL VERSION |
| GDA1 | GRADIENT | HESSIAN | | | |
| GDA2 | GRADIENT | | | | |
| GDA3 | HESSIAN | | | | |
| GDASTAT | ENERGY | GYRATE | WRITEOUT | | |
| GEODESIC | MINPATH | SORT3 | | | |
| GETBASE | PDBATM | | | | |
| GETIME | CLOCK | | | | |
| GETINT | BASEFILE MAKEXYZ | CHKXYZ NEXTARG | CONNECT READINT | FATAL SUFFIX | FREEUNIT VERSION |
| GETKEY | FATAL UPCASE | FREEUNIT | GETTEXT | SUFFIX | TRIMTEXT |
| GETMOL2 | BASEFILE VERSION | FREEUNIT | NEXTARG | READMOL2 | SUFFIX |
| GETNUCH | PDBATM | | | | |
| GETNUMB | TRIMTEXT | | | | |
| GETPDB | BASEFILE VERSION | FREEUNIT | NEXTARG | READPDB | SUFFIX |
| GETPRB | DIST2 | DOT | GETTOR | VCROSS | |

| | | | | | |
|----------|--|--|--|---|--|
| GETPRM | FREEUNIT SUFFIX | GETTEXT UPCASE | INITPRM VERSION | NEXTARG | READPRM |
| GETPROH | PDBATM | | | | |
| GETSEQ | GETWORD | TRIMTEXT | UPCASE | | |
| GETSEQN | GETTEXT | GETWORD | TRIMTEXT | UPCASE | |
| GETSIDE | PDBATM | | | | |
| GETTOR | DIST2 | | | | |
| GETXYZ | BASEFILE SUFFIX | FATAL VERSION | FREEUNIT | NEXTARG | READXYZ |
| GRADIENT | BOUNDS ECHARGE1 EHAL1 EMM3HB1 ESOLV1 EXTRA1 | EANGANG1 ECHGDPL1 EIMPROP1 EMPOLE1 ESTRBND1 PISCF | EANGLE1 EDIPOLE1 EIMPTOR1 EOPBEND1 ESTRTOR1 REPLICA | EBOND1 EGAUSS1 ELJ1 EOPDIST1 ETORS1 | EBUCK1 EGEOM1 EMETAL1 ERXNFLD1 EUREY1 |
| GRADRGD | GRADIENT | | | | |
| GRADROT | GRADIENT | ROTLIST | | | |
| HANGLE | NUMERAL | | | | |
| HBOND | NUMERAL | | | | |
| HDIPOLE | NUMERAL | | | | |
| HESSIAN | BORN EBUCK2 EGEOM2 EMETAL2 ERXNFLD2 EUREY2 REPLICA | BOUNDS ECHARGE2 EHAL2 EMM3HB2 ESOLV2 EXTRA2 | EANGANG2 ECHGDPL2 EIMPROP2 EMPOLE2 ESTRBND2 FATAL | EANGLE2 EDIPOLE2 EIMPTOR2 EOPBEND2 ESTRTOR2 INDUCE | EBOND2 EGAUSS2 ELJ2 EOPDIST2 ETORS2 PISCF |
| HESSRGD | GRADRGD | RIGIDXYZ | | | |
| HESSROT | GRADROT | MAKEXYZ | | | |
| HIMPTOR | NUMERAL | | | | |
| HSTRTOR | NUMERAL | | | | |
| HTORS | NUMERAL | | | | |

| | | | | | |
|-----------------|------------------------------|----------------------------|--------------------------------|-----------------------------|----------------------------|
| HYBRID | HANGLE HIMPTOR | HATOM HSTRBND | HBOND HSTRTOR | HCHARGE HTORS | HDIPOLE HVDW |
| IMPOSE | CENTER | QUATFIT | RMSFIT | | |
| INDUCE | INDUCE0A | INDUCE0B | | | |
| INDUCE0A | FATAL | GROUPS | IMAGE | PRTERR | SWITCH |
| INDUCE0B | FATAL UMUTUAL2 | PRTERR | UDIRECT1 | UDIRECT2 | UMUTUAL1 |
| INEDGE | CERROR | | | | |
| INERTIA | JACOBI | | | | |
| INITERR | LOCERR | TORSER | | | |
| INITIAL | COMMAND | INITRES | PRECISE | PROMO | |
| INITROT | FATAL | NEXTARG | ROTCHECK | ROTLIST | |
| INTEDIT | FIELD GETWORD TRIMTEXT | FINAL INITIAL UPCASE | FREEUNIT MAKEXYZ VERSION | GEOMETRY NUMBER ZHELP | GETINT PRTINT ZVALUE |
| INTXYZ | FINAL VERSION | FREEUNIT | GETINT | INITIAL | PRTXYZ |
| INVBETA | BETAI | GAMMLN | | | |
| INVERT | FATAL | | | | |
| IPEDGE | CERROR | | | | |
| KANGANG | GETTEXT | UPCASE | | | |
| KANGLE | GETTEXT | NUMERAL | UPCASE | | |
| KATOM | GETNUMB | GETSTRING | GETTEXT | UPCASE | |
| KBOND | GETTEXT | KENEG | NUMERAL | UPCASE | |
| KCHARGE | GETTEXT | UPCASE | | | |
| KDIPOLE | GETTEXT | NUMERAL | UPCASE | | |
| KENEG | GETTEXT | NUMERAL | UPCASE | | |
| KEWALD | EWALDCOF | FATAL | FFTSETUP | GETTEXT | MODULI |

| | | | | | |
|-----------------|-------------------|----------|----------|---------|----------|
| | UPCASE | | | | |
| KGEOM | FATAL | GEOMETRY | GETTEXT | UPCASE | |
| KIMPROP | GETTEXT | NUMERAL | UPCASE | | |
| KIMPTOR | GETTEXT | NUMERAL | TORPHASE | UPCASE | |
| KMPOLE | GETTEXT UPCASE | NUMBER | NUMERAL | RANDOM | SORT3 |
| KOPBEND | GETTEXT | NUMBER | NUMERAL | UPCASE | |
| KOPDIST | GETTEXT | NUMERAL | UPCASE | | |
| KORBIT | GETTEXT | NUMERAL | UPCASE | | |
| KPOLAR | GETTEXT | POLARGRP | UPCASE | | |
| KSOLV | GETTEXT | GETWORD | KANGLE | KBOND | UPCASE |
| KSTRBND | GETTEXT | UPCASE | | | |
| KSTRTOR | GETTEXT | NUMERAL | UPCASE | | |
| KTORS | GETTEXT | NUMERAL | TORPHASE | UPCASE | |
| KUREY | GETTEXT | NUMERAL | UPCASE | | |
| KVDW | GETTEXT | NUMBER | NUMERAL | UPCASE | |
| LBFGS | COMMENT | GETTEXT | SEARCH | UPCASE | WRITEOUT |
| LIGASE | FINDATM | | | | |
| LIGHTS | FATAL | SORT2 | SORT5 | | |
| LMSTEP | PRECISE | QRSOLVE | | | |
| LOCALMIN | GRADIENT | TNCG | | | |
| LOCALRGD | OCVM | | | | |
| LOCALROT | OCVM | | | | |
| LOCALXYZ | TNCG | | | | |
| MAJORIZE | GETIME | GYRATE | RMSERROR | SETIME | |
| MAKEINT | ADJACENT | FATAL | GEOMETRY | GETTEXT | UPCASE |

| | | | | | |
|------------------|---|---|--|---|---|
| MAKEPDB | ATTACH GETSIDE | FREEUNIT NUMERAL | GETBASE PDBATM | GETNUCH READSEQ | GETPROH VERSION |
| MAKEXYZ | XYZATM | | | | |
| MAPCHECK | FREEUNIT | NUMERAL | PRTXYZ | VERSION | |
| MAXWELL | ERFINV | RANDOM | | | |
| MDINIT | FREEUNIT LATTICE READDYN | GETTEXT MAXWELL UPCASE | GETWORD MDREST VERSION | GRADIENT NUMERAL | GRPLINE RANVEC |
| MDREST | INVERT | | | | |
| MDSAVE | FATAL PRTXYZ | FREEUNIT SUFFIX | NUMERAL VERSION | OPENEND | PRTDYN |
| MEASFN | CERROR | TRIPLE | VCROSS | VECANG | VNORM |
| MEASFP | CERROR | DOT | VCROSS | VECANG | VNORM |
| MEASFS | CERROR | DOT | VECANG | VNORM | |
| MEASPM | VCROSS | | | | |
| MECHANIC | ACTIVE CUTOFFS KATOM KGEOM KOPBEND KSTRBND LATTICE RINGS | ANGLES FATAL KBOND KIMPROP KOPDIST KSTRTOR MOLECULE SMOOTH | ATTACH FIELD KCHARGE KIMPTOR KORBIT KTORS MUTATE TORSIONS | BONDS KANGANG KDIPOLE KMETAL KPOLAR KUREY ORBITAL UNITCELL | CLUSTER KANGLE KEWALD KMPOLE KSOLV KVDW POLYMER |
| MERGE | FATAL | GETREF | | | |
| MIDERR | BNDERR | CHIRER | LOCERR | TORSER | |
| MINIMIZE1 | GRADIENT | | | | |
| MINIMIZE | FINAL LBFGS | FREEUNIT MECHANIC | GETXYZ NEXTARG | GRADIENT PRTXYZ | INITIAL VERSION |
| MINIROT | FINAL INITROT VERSION | FREEUNIT LBFGS | GETINT MECHANIC | GRADROT NEXTARG | INITIAL PRTINT |
| MINIROT1 | GRADROT | MAKEXYZ | | | |
| MINRIGID | FINAL | FREEUNIT | GETXYZ | GRADRGD | INITIAL |

| | | | | | |
|------------------|----------------------------|---------------------------------|-------------------------------|--------------------|-------------------|
| | LBFGS VERSION | MECHANIC | NEXTARG | ORIENT | PRTXYZ |
| MINRIGID1 | GRADRGD | RIGIDXYZ | XYZRIGID | | |
| MMID | DERIVS | | | | |
| MODECART | CLIMBXYZ | EIGENXYZ | GETREF | IMPOSE | MAKEREf |
| MODEROT | CLIMBROT | EIGENROT | MAKEXYZ | | |
| MODESRCH | CLIMBER | EIGENROT | MAKEINT | MAKEREf | MAPCHECK |
| MODETORS | CLIMBTOR MAKEREf | EIGENTOR | GETREF | IMPOSE | MAKEINT |
| MODULI | BSPLINE | DFTMOD | | | |
| MOLECULE | SORT | SORT3 | | | |
| MOLUIND | UFIELD | | | | |
| MOMENTS | INDUCE | JACOBI | ROTMAT | ROTPOLE | |
| MUTATE | GETTEXT | UPCASE | | | |
| NEIGHBOR | CERROR | DIST2 | | | |
| NEWATM | ADDBOND | XYZATM | | | |
| NEWTON | FINAL INITIAL UPCASE | FREEUNIT MECHANIC VERSION | GETTEXT NEXTARG | GETXYZ PRTXYZ | GRADIENT TNCG |
| NEWTON1 | GRADIENT | | | | |
| NEWTON2 | HESSIAN | | | | |
| NEWTROT | FINAL INITIAL TNCG | FREEUNIT INITROT UPCASE | GETINT MECHANIC VERSION | GETTEXT NEXTARG | GRADROT PRTINT |
| NEWTROT1 | GRADROT | MAKEXYZ | | | |
| NEWTROT2 | HESSROT | MAKEXYZ | | | |
| NORMAL | RANDOM | | | | |
| NUCBASE | OCVM | ORIENT | POTOFF | ZATOM | |
| NUCCHAIN | NUCBASE | OCVM | ORIENT | ZATOM | |

| | | | | | |
|-----------|--|---|--|--------------------------------------|-------------------------------|
| NUCLEIC | BASEFILE GETKEY MOLECULE PRTXYZ | CONNECT GETSEQN NEXTARG TRIMTEXT | DELETE INITIAL NUCCHAIN VERSION | FIELD MAKEINT PRTINT WATSON | FREEUNIT MAKEXYZ PRTSEQ |
| NUMBER | TRIMTEXT | | | | |
| OCVM | GETTEXT | PRECISE | UPCASE | WRITEOUT | |
| OLDATM | ADDBOND | FATAL | | | |
| OPTIMIZE1 | GRADIENT | | | | |
| OPTIMIZE | FATAL INITIAL VERSION | FINAL MECHANIC | FREEUNIT NEXTARG | GETXYZ OCVM | GRADIENT PRTXYZ |
| OPTIROT | FATAL INITIAL PRTINT | FINAL INITROT VERSION | FREEUNIT MECHANIC | GETINT NEXTARG | GRADROT OCVM |
| OPTIROT1 | GRADROT | MAKEXYZ | | | |
| OPTRIGID | FATAL INITIAL PRTXYZ | FINAL MECHANIC VERSION | FREEUNIT NEXTARG | GETXYZ OCVM | GRADRGD ORIENT |
| OPTRIGID1 | GRADRGD | RIGIDXYZ | XYZRIGID | | |
| ORBITAL | FATAL | GETTEXT | PIPLANE | UPCASE | |
| ORIENT | XYZRIGID | | | | |
| OVERLAP | SLATER | | | | |
| PATH | FINAL LBFGS WRITEOUT | GETXYZ MECHANIC | IMPOSE NEXTARG | INITIAL ORTHOG | INVERT POTNRG |
| PATH1 | POTNRG | | | | |
| PATHPNT | OCVM | | | | |
| PATHSCAN | PATHPNT | SADDLE1 | TANGENT | | |
| PATHVAL | IMPOSE | | | | |
| PDBXYZ | CHKXYZ GETNUMB RIBOSOME | DELETE GETPDB SORT | FIELD INITIAL UPCASE | FINAL LIGASE VERSION | FREEUNIT PRTXYZ |

| | | | | | |
|-----------------|--|---|--|---|----------------------------|
| PIPLANE | FATAL | | | | |
| PISCF | NEWATM | | | | |
| PITILT | OLDATM | | | | |
| PLACE | CERROR | DIST2 | GETPRB | GETTOR | INEDGE |
| POLARGRP | SORT | SORT8 | | | |
| POLARIZE | FATAL MOLUIND | GETXYZ | INITIAL | JACOBI | MECHANIC |
| POLYMER | FATAL | GETTEXT | IMAGE | UPCASE | |
| POTNRG | GRADIENT | | | | |
| POWER | RANDOM | | | | |
| PRECOND | CHOLSKY | COLUMN | | | |
| PRESSURE | LATTICE | | | | |
| PRMKEY | GETTEXT | GETWORD | POTOFF | UPCASE | |
| PROCHAIN | GETTEXT | PROSIDE | UPCASE | ZATOM | |
| PROJCT | DOT | | | | |
| PROPYZE | GYRATE | INERTIA | MOMENTS | | |
| PROSIDE | FREEUNIT | PRTINT | PRTXYZ | VERSION | ZATOM |
| PROTEIN | BASEFILE FINAL MAKEINT PRTSEQ | CHKXYZ FREEUNIT MAKEXYZ PRTXYZ | CONNECT GETKEY NEXTARG TRIMTEXT | DELETE GETSEQ PROCHAIN VERSION | FIELD INITIAL PRTINT |
| PRTDYN | ZATOM | | | | |
| PRTERR | ZATOM | | | | |
| PRTINT | VERSION | | | | |
| PRTMOL2 | NUMBER | VERSION | | | |
| PRTMSI | VERSION | | | | |
| PRTPDB | VERSION | | | | |

| | | | | | |
|-----------------|--|---|--|------------------------------|-------------------------------|
| PRTPRM | NUMBER | | | | |
| PRTSEQ | VERSION | | | | |
| PRTXMOL | VERSION | | | | |
| PRTXYZ | VERSION | | | | |
| PSS | ACTIVE INITIAL MECHANIC SIGMOID | FINAL INITROT MODECART UPCASE | GETTEXT LOCALXYZ MODETORS | GETXYZ MAKEINT NEXTARG | IMPOSE MAKEREf PSSWRITE |
| PSS1 | GRADIENT | | | | |
| PSS2 | HESSIAN | | | | |
| PSSRGD1 | GRADRGD | RIGIDXYZ | | | |
| PSSRIGID | FINAL INITIAL OCVM SIGMOID | FREEUNIT MAKEREf ORIENT UPCASE | GETTEXT MECHANIC PRTXYZ VERSION | GETXYZ NEXTARG RGDSRCH | IMPOSE NUMERAL RIGIDXYZ |
| PSSROT | FINAL INITIAL MODEROT UPCASE | FREEUNIT INITROT NEXTARG VERSION | GETINT MAKEREf NUMERAL | GETTEXT MAKEXYZ OCVM | IMPOSE MECHANIC PRTXYZ |
| PSSROT1 | GRADROT | MAKEXYZ | | | |
| PSSWRITE | FREEUNIT | NUMERAL | PRTXYZ | VERSION | |
| PTINCY | DOT | EPUCLC | PROJCT | ROTANG | |
| QUATFIT | JACOBI | | | | |
| RADIAL | FINAL INITIAL TRIMTEXT | FREEUNIT LATTICE UNITCELL | GETWORD MOLECULE VERSION | GETXYZ NUMERAL | IMAGE READXYZ |
| RANDOM | CALENDAR | GETTEXT | UPCASE | | |
| RANVEC | RANDOM | | | | |
| RATTLE | FATAL | IMAGE | PRTErr | | |
| RATTLE2 | FATAL | IMAGE | PRTErr | | |
| READBLK | FATAL | FREEUNIT | GETWORD | NUMERAL | |

| | | | | | |
|-----------------|---|--|--|---|--|
| READDYN | FATAL | VERSION | | | |
| READINT | FATAL VERSION | GETTEXT | GETWORD | NEXTTEXT | TRIMTEXT |
| READMOL2 | FATAL UPCASE | GETTEXT VERSION | GETWORD | SORT | TRIMTEXT |
| READPDB | FATAL VERSION | FIXPDB | GETTEXT | TRIMTEXT | UPCASE |
| READPRM | FATAL NUMERAL | GETNUMB PRMKEY | GETSTRING TORPHASE | GETTEXT TRIMTEXT | GETWORD UPCASE |
| READSEQ | FATAL VERSION | GETNUMB | GETTEXT | GETWORD | TRIMTEXT |
| READXYZ | CHKXYZ SORT | FATAL TRIMTEXT | GETTEXT VERSION | GETWORD | NEXTTEXT |
| REFINE | LBFGS | | | | |
| REPLICA | FATAL | | | | |
| RGDSRCH | CLIMBRGD | EIGENRGD | RIGIDXYZ | | |
| RGDSTEP | GRADIENT PRESSURE | LINBODY REGBODY | MDSAVE ROTRGD | MDSTAT TEMPER | NEWCRD |
| RIBOSOME | ADDBOND NEWATM | ADDSIDE OLDATM | FATAL PRTSEQ | FINDATM VERSION | FREEUNIT |
| RINGS | ANGLES | BONDS | FATAL | TORSIONS | |
| RMSERROR | TRIMTEXT | | | | |
| ROTANG | DOT | VCROSS | | | |
| ROTCHECK | ROTLIST | | | | |
| ROTLIST | FATAL | | | | |
| SADDLE | COMMENT GETXYZ MECHANIC PRTXYZ UPCASE | FATAL IMPOSE NEXTARG READXYZ VERSION | FINAL INITIAL PATHPNT SADDLE1 | FREEUNIT MAKEINT PATHSCAN SEARCH | GETTEXT MAKEXYZ PATHVAL TANGENT |
| SADDLE1 | GRADIENT | | | | |
| SADDLES | CERROR | IPEDGE | TRIPLE | | |

| | | | | | |
|------------------|-------------------------------|------------------------------------|--------------------------------|--------------------------------|--------------------------------|
| SCAN | ACTIVE INITROT MODESRCH | FINAL LOCALMIN NEXTARG | FREEUNIT MAKEINT NUMERAL | GETXYZ MAPCHECK READYXYZ | INITIAL MECHANIC VERSION |
| SCAN1 | GRADIENT | | | | |
| SCAN2 | HESSIAN | | | | |
| SDAREA | SURFATOM | | | | |
| SDSTEP | GRADIENT SDTERM | MDSAVE | MDSTAT | RATTLE | RATTLE2 |
| SDTERM | NORMAL | SDAREA | | | |
| SETIME | CLOCK | | | | |
| SHAKEUP | GETNUMB | GETTEXT | GETWORD | UPCASE | |
| SLATER | ASET | BSET | CJKM | POLYP | |
| SMOOTH | GETTEXT | NEXTARG | UPCASE | | |
| SNIFFER | FINAL INITIAL SNIFFER1 | FREEUNIT MAKERE F VERSION | GETREF MECHANIC WRITEOUT | GETXYZ NEXTARG | GRADIENT PRTXYZ |
| SNIFFER1 | GRADIENT | | | | |
| SOAK | DELETE MERGE VERSION | FREEUNIT MOLECULE | IMAGE READYXYZ | LATTICE SUFFIX | MAKERE F UNITCELL |
| SPACEFILL | ACTIVE GETTEXT NEXTARG | CONNOLLY GETXYZ READYXYZ | FIELD INITIAL SUFFIX | FINAL KATOM UPCASE | FREEUNIT KVDW VERSION |
| SPECTRUM | BASEFILE VERSION | FREEUNIT | INITIAL | NEXTARG | SUFFIX |
| SQUARE | GETTEXT TRUST | LMSTEP UPCASE | PRECISE WRITEOUT | QRFACT | RSDVALUE |
| SUFFIX | TRIMTEXT | | | | |
| SUPERPOSE | FIELD IMPOSE READYXYZ | FINAL INITIAL SUFFIX | FREEUNIT KATOM TRIMTEXT | GETTEXT NEXTARG UPCASE | GETXYZ PRTXYZ VERSION |
| SURFACE | FATAL | SORT2 | | | |

| | | | | | |
|------------------|---------------------------------------|-------------------------------------|---------------------------------------|--------------------------------------|------------------------------------|
| SURFATOM | FATAL | SORT2 | | | |
| SWITCH | REPLICA | | | | |
| SYBYLXYZ | FINAL VERSION | FREEUNIT | GETMOL2 | INITIAL | PRTXYZ |
| SYMMETRY | CELLATOM | | | | |
| TANGENT | PATHPNT | SADDLE1 | | | |
| TEMPER | MAXWELL | RANDOM | RANVEC | | |
| TESTGRAD | ENERGY INITIAL | FINAL MECHANIC | GETTEXT NEXTARG | GETXYZ UPCASE | GRADIENT |
| TESTHESS | FINAL HESSIAN UPCASE | FREEUNIT INITIAL VERSION | GETTEXT MECHANIC | GETXYZ NEXTARG | GRADIENT NUMGRAD |
| TESTLIGHT | EBUCK EGAUSS1 EMM3HB INITIAL | EBUCK1 EHAL EMM3HB1 LIGHTS | ECHARGE EHAL1 FINAL MECHANIC | ECHARGE1 ELJ GETIME NEXTARG | EGAUSS ELJ1 GETXYZ SETIME |
| TESTROT | ENERGY INITROT | FINAL MAKEXYZ | GETINT MECHANIC | GRADROT NEXTARG | INITIAL |
| TIMER | ENERGY GRADIENT SETIME | FINAL HESSIAN UPCASE | GETIME INITIAL | GETTEXT MECHANIC | GETXYZ NEXTARG |
| TIMEROT | ENERGY GRADROT NEXTARG | FINAL HESSROT SETIME | GETIME INITIAL UPCASE | GETINT INITROT | GETTEXT MECHANIC |
| TNCG | GETTEXT UPCASE | HMATRIX WRITEOUT | PISCF | SEARCH | TNSOLVE |
| TNSOLVE | PRECOND | | | | |
| TORSIONS | FATAL | | | | |
| TORUS | CERROR | GETTOR | | | |
| TOTERR | BNDERR | CHIRER | LOCERR | TORSER | VDWERR |
| TRIANGLE | FATAL | | | | |
| TRIPLE | DOT | VCROSS | | | |

| | | | | | |
|-----------------|----------------------------|-----------------------------|--------------------------|---------------------|-------------------|
| TRUST | PRECISE | RSDVALUE | | | |
| UDIRECT2 | ERFC | IMAGE | SWITCH | | |
| UMUTUAL2 | ERFC | IMAGE | SWITCH | | |
| UNITCELL | FATAL | GETTEXT | GETWORD | UPCASE | |
| VAM | CERROR GENDOT TRIPLE | CIRPLN MEASFN VCROSS | DEPTH MEASFP VNORM | DIST2 MEASFS | DOT MEASPM |
| VDWERR | LIGHTS | | | | |
| VECANG | ANORM | DOT | TRIPLE | | |
| VERLET | GRADIENT RATTLE2 | MDSAVE TEMPER | MDSTAT | PRESSURE | RATTLE |
| VERSION | LOWCASE | NEXTARG | TRIMTEXT | | |
| VIBRATE | DIAGQ HESSIAN PRTXYZ | FATAL INITIAL VERSION | FINAL MECHANIC | FREEUNIT NEXTARG | GETXYZ NUMERAL |
| VIBROT | DIAGQ INITROT | FINAL MECHANIC | GETINT | HESSROT | INITIAL |
| VNORM | ANORM | | | | |
| VOLUME | CONNOLLY | | | | |
| VOLUME1 | FATAL | | | | |
| VOLUME2 | FATAL | | | | |
| WATSON | ZATOM | | | | |
| WATSON1 | GRADRGD | RIGIDXYZ | | | |
| WRITEOUT | FREEUNIT | NUMERAL | PRTINT | PRTXYZ | VERSION |
| XTALERR | ENERGY | XTALMOVE | XTALPRM | | |
| XTALFIT | FINAL SQUARE | GETXYZ XTALPRM | INITIAL | MECHANIC | NEXTARG |
| XTALLAT1 | ENERGY | LATTICE | | | |
| XTALMIN | FINAL | FREEUNIT | GETXYZ | GRADIENT | INITIAL |

| | | | | | |
|-----------------|--|---|------------------------------------|--------------------------------------|------------------------------------|
| | LATTICE TNCG | MECHANIC VERSION | NEXTARG XTALLAT1 | OCVM | PRTXYZ |
| XTALMOL1 | GRADIENT | | | | |
| XTALMOL2 | HESSIAN | | | | |
| XTALMOVE | LATTICE | | | | |
| XTALPRM | BOUNDS | LATTICE | MOLECULE | | |
| XYZEDIT | ACTIVE FREEUNIT INSERT PRTXYZ UNITCELL | CUTOFFS GETXYZ KATOM RANDOM VERSION | DELETE IMAGE LATTICE SOAK | FIELD INERTIA MAKEREFS SORT | FINAL INITIAL MERGE SORT4 |
| XYZINT | FINAL MAKEINT VERSION | FREEUNIT NEXTARG | GETTEXT PRTINT | GETXYZ READINT | INITIAL UPCASE |
| XYZPDB | FIELD KATOM | FINAL MAKEPDB | FREEUNIT MOLECULE | GETXYZ P RTPDB | INITIAL VERSION |
| XYZRIGID | JACOBI | ROTEULER | | | |
| XYZSYBYL | BONDS PRTMOL2 | FINAL VERSION | FREEUNIT | GETXYZ | INITIAL |
| ZATOM | FATAL | | | | |
| ZVALUE | MAKEXYZ | TRIMTEXT | | | |

12. Examples using the TINKER Package

This section contains brief descriptions of the sample calculations found in the EXAMPLE subdirectory of the TINKER distribution. These examples exercise several of the current TINKER programs and are intended to provide a flavor of the capabilities of the package.

ANION Example

Computes an estimation of the free energy of hydration of Cl^- anion vs. Br^- anion via a 2 picosecond simulation on a “hybrid” anion in a box of water followed by a free energy perturbation calculation

ARGON Example

Performs an initial energy minimization on a periodic box containing 150 argon atoms followed by 6 picoseconds of a molecular dynamics using a modified Beeman integration algorithm and a Bersedsen thermostat

CLUSTER Example

Performs a set of 10 Gaussian density annealing (GDA) trials on a cluster of 13 argon atoms in an attempt to locate the global minimum energy structure

CRAMBIN Example

Generates a TINKER file from a PDB file, followed by a single point energy computation and determination of the molecular volume and surface area

CYCLOHEX Example

First approximately locates the transition state between chair and boat cyclohexane, followed by subsequent refinement of the transition state and a final vibrational analysis to show that a single negative frequency is associated with the saddle point

ENKEPHALIN Example

Produces coordinates from the met-enkephalin amino acid sequence and phi/psi angles, followed by truncated Newton energy minimization and determination of the lowest frequency normal mode

FORMAMIDE Example

Converts to a unit cell from fractional coordinates, followed by full crystal energy minimization and determination of optimal carbonyl oxygen energy parameters from a fit to lattice energy and structure

HELIX Example

Performs a rigid-body optimization of the packing of two idealized polyalanine helices using only van der Waals interactions

SALT Example

Converts a sodium chloride assymetric unit to the corresponding unit cell, then runs a crystal minimization starting from the initial diffraction structure using Ewald summation to model the long-range electrostatic interactions.

13. Benchmark Results

The tables in this section provide CPU benchmarks for basic TINKER energy and derivative evaluations, vibrational analysis and molecular dynamics. All times are in seconds and were measured with TINKER executables dimensioned to **maxatm** of 10000 and **maxhess** of 1000000 in the source file **sizes.i**. All calculations were run twice in rapid succession on a quiet machine. The times reported for each benchmark are the results from the second run. If you have built TINKER on an alternative machine type and are able to run the benchmarks on the additional machine type, please send the results for inclusion in a future listing.

BENCHMARK #1: Calmodulin Energy Evaluation

The system is an isolated molecule of the 148-residue protein calmodulin with 2264 atoms using the Amber-94 force field. All interactions are computed with no use of cutoffs. Times listed are for calculation setup followed by a single energy, energy/gradient and Hessian evaluation.

| MACHINE TYPE | MHz | SETUP | ENERGY | GRAD | HESS |
|--------------------------------------|------------|--------------|---------------|-------------|-------------|
| Compaq DS10 (Tru64 5.0) | 466 | 0.35 | 1.28 | 1.90 | 8.05 |
| DEC Alpha 4100 (Tru64 5.0) | 400 | 0.55 | 2.22 | 3.92 | 15.56 |
| SGI IndigoII R10K (Irix 6.5) | 195 | 1.16 | 3.50 | 5.93 | 22.51 |
| Target-USA Athlon (RH 6.2, PGI) | 950 | 0.35 | 0.67 | 1.59 | 8.04 |
| Target-USA Athlon (RH 6.2, Absoft) | 950 | 0.33 | 0.94 | 1.68 | 8.58 |
| Target-USA Athlon (RH 6.2, g77) | 950 | 0.38 | 0.91 | 1.95 | 9.85 |
| Compaq Armada M700 (RH 6.2, PGI) | 650 | 0.46 | 1.35 | 2.69 | 14.20 |
| Compaq Armada M700 (RH 6.2, Absoft) | 650 | 0.39 | 1.46 | 2.70 | 15.45 |
| Compaq Armada M700 (RH 6.2, g77) | 650 | 0.43 | 1.66 | 3.65 | 17.21 |
| Compaq Armada M700 (Win2K, CVF 6.5) | 650 | 0.35 | 1.29 | 2.32 | 12.10 |
| Compaq Armada M700 (Win2K, LF95 5.5) | 650 | 0.32 | 1.83 | 2.88 | 15.90 |

BENCHMARK #2: Crambin Crystal Energy Evaluation

The system is a unit cell of the 46-residue protein crambin containing 2 polypeptide chains, 2 ethanol and 178 water molecules for a total of 1360 atoms using the OPLS-UA force field. Periodic boundaries are used with particle mesh Ewald for electrostatics and a 9.0 Å cutoff for vdW interactions. Times listed are for calculation setup followed by a single energy, energy/gradient and Hessian evaluation.

| MACHINE TYPE | MHz | SETUP | ENERGY | GRAD | HESS |
|------------------------------------|------------|--------------|---------------|-------------|-------------|
| Compaq DS10 (Tru64 5.0) | 466 | 0.28 | 0.48 | 0.63 | 1.78 |
| DEC Alpha 4100 (Tru64 5.0) | 400 | 0.46 | 0.85 | 1.08 | 3.46 |
| SGI IndigoII R10K (Irix 6.5) | 195 | 0.92 | 1.00 | 1.43 | 4.02 |
| Target-USA Athlon (RH 6.2, PGI) | 950 | 0.32 | 0.33 | 0.48 | 1.30 |
| Target-USA Athlon (RH 6.2, Absoft) | 950 | 0.26 | 0.39 | 0.53 | 1.67 |
| Target-USA Athlon (RH 6.2, g77) | 950 | 0.34 | 0.46 | 0.64 | 1.77 |
| Compaq Armada M700 (RH 6.2, PGI) | 650 | 0.42 | 0.52 | 0.76 | 2.62 |

| | | | | | |
|--------------------------------------|-----|------|------|------|------|
| Compaq Armada M700 (RH 6.2, Absoft) | 650 | 0.30 | 0.62 | 0.89 | 3.29 |
| Compaq Armada M700 (RH 6.2, g77) | 650 | 0.37 | 0.75 | 1.02 | 3.55 |
| Compaq Armada M700 (Win2K, CVF 6.5) | 650 | 0.29 | 0.51 | 0.72 | 2.43 |
| Compaq Armada M700 (Win2K, LF95 5.5) | 650 | 0.28 | 0.61 | 0.78 | 3.33 |

BENCHMARK #3: Peptide Normal Mode Calculation

The system is a minimum energy conformation of a 20-residue peptide containing one of each of the standard amino acids for a total of 328 atoms using the OPLS-AA force field without cutoffs. The time reported is for computation of the Hessian and calculation of the normal modes of the Hessian matrix and the vibration frequencies requiring two separate matrix diagonalization steps.

| MACHINE TYPE | MHz | NORMAL MODES |
|--------------------------------------|-----|--------------|
| Compaq DS10 (Tru64 5.0) | 466 | 41 |
| DEC Alpha 4100 (Tru64 5.0) | 400 | 76 |
| SGI IndigoII R10K (Irix 6.5) | 195 | 145 |
| Target-USA Athlon (RH 6.2, PGI) | 950 | 51 |
| Target-USA Athlon (RH 6.2, Absoft) | 950 | 54 |
| Target-USA Athlon (RH 6.2, g77) | 950 | 64 |
| Compaq Armada M700 (RH 6.2, PGI) | 650 | 66 |
| Compaq Armada M700 (RH 6.2, Absoft) | 650 | 71 |
| Compaq Armada M700 (RH 6.2, g77) | 650 | 119 |
| Compaq Armada M700 (Win2K, CVF 6.5) | 650 | 63 |
| Compaq Armada M700 (Win2K, LF95 5.5) | 650 | 86 |

BENCHMARK #4: TIP3P Water Box Molecular Dynamics

The system consists of 216 rigid TIP3P water molecules in a 18.643 Å periodic box, 9.0 Å shifted energy switch cutoffs for nonbonded interactions. The time reported is for 1000 dynamics steps of 1.0 fs each using the modified Beeman integrator and Rattle constraints on all bond lengths.

| MACHINE TYPE | MHz | DYNAMICS |
|--------------------------------------|-----|----------|
| Compaq DS10 (Tru64 5.0) | 466 | 104 |
| DEC Alpha 4100 (Tru64 5.0) | 400 | 195 |
| SGI IndigoII R10K (Irix 6.5) | 195 | 278 |
| Target-USA Athlon (RH 6.2, PGI) | 950 | 70 |
| Target-USA Athlon (RH 6.2, Absoft) | 950 | 77 |
| Target-USA Athlon (RH 6.2, g77) | 950 | 125 |
| Compaq Armada M700 (RH 6.2, PGI) | 650 | 131 |
| Compaq Armada M700 (RH 6.2, Absoft) | 650 | 138 |
| Compaq Armada M700 (RH 6.2, g77) | 650 | 226 |
| Compaq Armada M700 (Win2K, CVF 6.5) | 650 | 130 |
| Compaq Armada M700 (Win2K, LF95 5.5) | 650 | 141 |

BENCHMARK #5: TINKER Water Box Molecular Dynamics

The system consists of 216 TINKER flexible polarizable atomic multipole water molecules in a 18.643 Å periodic box using regular Ewald summation for the electrostatics and a 12.0 Å switched cutoff for vdW interactions. The time reported is for 100 dynamics steps of 1.0 fs each using the modified Beeman integrator and 0.01 Debye rms convergence for induced dipole moments.

| MACHINE TYPE | MHz | DYNAMICS |
|--------------------------------------|-----|----------|
| Compaq DS10 (Tru64 5.0) | 466 | 304 |
| DEC Alpha 4100 (Tru64 5.0) | 400 | 527 |
| SGI IndigoII R10K (Irix 6.5) | 195 | 738 |
| Target-USA Athlon (RH 6.2, PGI) | 950 | 243 |
| Target-USA Athlon (RH 6.2, Absoft) | 950 | 281 |
| Target-USA Athlon (RH 6.2, g77) | 950 | 376 |
| Compaq Armada M700 (RH 6.2, PGI) | 650 | 443 |
| Compaq Armada M700 (RH 6.2, Absoft) | 650 | 525 |
| Compaq Armada M700 (RH 6.2, g77) | 650 | 811 |
| Compaq Armada M700 (Win2K, CVF 6.5) | 650 | 411 |
| Compaq Armada M700 (Win2K, LF95 5.5) | 650 | 535 |

14. Collaborators & Acknowledgments

The TINKER package has developed over a period of many years, very slowly during the late-1980's, and more rapidly since the mid-1990's in Jay Ponder's research group at the Washington University School of Medicine in St. Louis. Many people have played significant roles in the development of the package into its current form. The major contributors are listed below:

| | |
|---|--|
| Stew Rubenstein | coordinate interconversions; original optimization methods and torsional angle manipulation |
| Craig Kundrot | molecular surface area & volume and their derivatives |
| Shawn Huston | original AMBER/OPLS implementation; free energy calculations; time correlation functions |
| Mike Dudek | DMA-derived multipole models for peptides and proteins |
| Yong "Mike" Kong | multipole electrostatics; dipole polarization; reaction field treatment; TINKER water model |
| Reece Hart | potential smoothing methodology; Scheraga's DEM, Straub's GDA and extensions |
| Mike Hodsdon | extension of the TINKER distgeom program and its application to NMR NOE structure determination |
| Rohit Pappu | potential smoothing methodology and PSS algorithms; rigid body optimization; GB/SA solvation derivatives |
| Wijnand Mooij | MM3 directional hydrogen bonding term; crystal lattice minimization code |
| Gerald Loeffler | stochastic/Langevin dynamics implementation |
| Marina Vorobieva Nina Sokolova | nucleic acid building module and parameter translation |
| Peter Bagossi | TINKER force field parameters for alkanes and diatomics |
| Pengyu Ren | Ewald summation for polarizable atomic multipoles; TINKER force field for water, organics and peptides |
| Anders Carlsson | ligand field potential energy term for transition metals |
| Andrey Kutepov | integrator for rigid-body dynamics trajectories |

In addition, we would like to thank **Tom Darden** for making his particle mesh Ewald code generally available to the simulation community.

It is critically important that TINKER's distributed force field parameter sets exactly reproduce the intent of the original force field authors. We would like to thank **Julian Tirado-Rives** (OPLS-AA), **Alex MacKerell** (CHARMM27), and **Adrian Roitberg** and **Carlos Simmerling** (AMBER) for their help in testing TINKER's results against those given by the authentic programs and parameter sets. **Lou Allinger** has provided updated parameters for MM2 and MM3 on several occasions. His very successful methods provided the original inspiration for the development of TINKER.

Finally, we wish to thank the many users of the TINKER package for their suggestions and comments, praise and criticism, which have resulted in a variety of improvements.

15. References & Suggested Reading

This section contains a list of the references to general theory, algorithms and implementation details which have been of use during the development of the TINKER package. Methods described in some of the references have been implemented in detail within the TINKER source code. Other references contain useful background information although the algorithms themselves are now obsolete. Still other papers contain ideas or extensions planned for future inclusion in TINKER. References for specific force field parameter sets are provided in an earlier section of this User's Guide. This list is heavily skewed toward biomolecules in general and proteins in particular. This bias reflects our group's major interests; however an attempt has been made to include methods which should be generally applicable.

PARTIAL LIST OF MOLECULAR MECHANICS SOFTWARE PACKAGES

| | |
|-------------|---|
| AMBER | Peter Kollman, University of California, San Francisco |
| AMMP | Rob Harrison, Thomas Jefferson University, Philadelphia |
| ARGOS | Andy McCammon, University of California, San Diego |
| BOSS | William Jorgensen, Yale University |
| BRUGEL | Shoshona Wodak, Free University of Brussels |
| CFF | Shneior Lifson, Weizmann Institute |
| CHARMM | Martin Karplus, Harvard University |
| CHARMM/GEMM | Bernard Brooks, National Institutes of Health, Bethesda |
| DELPHI | Bastian van de Graaf, Delft University of Technology |
| DISCOVER | Molecular Simulations Inc., San Diego |
| DL_POLY | W. Smith & T. Forester, CCP5, Daresbury Laboratory |
| ECEPP | Harold Scheraga, Cornell University |
| ENCAD | Michael Levitt, Stanford University |
| FANTOM | Werner Braun, University of Texas, Galveston |
| FEDER/2 | Nobuhiro Go, Kyoto University |
| GROMACS | Herman Berendsen, University of Groningen |
| GROMOS | Wilfred van Gunsteren, BIOMOS and ETH, Zurich |
| IMPACT | Ronald Levy, Rutgers University |
| MACROMODEL | Schodinger, Inc., Jersey City, New Jersey |
| MM2/MM3/MM4 | N. Lou Allinger, University of Georgia |
| MMC | Cliff Dykstra, Indiana Univ.-Purdue Univ. at Indianapolis |
| MMFF | Tom Halgren, Merck Research Laboratories, Rahway |
| MMTK | Konrad Hinsen, Inst. of Structural Biology, Grenoble |
| MOIL | Ron Elber, Cornell University |
| MOLARIS | Arieh Warshal, University of Southern California |
| MOLDY | Keith Refson, Oxford University |
| MOSCITO | Dietmar Paschek & Alfons Geiger, Universität Dortmund |
| NAMD | Klaus Schulten, University of Illinois, Urbana |
| OOMPAA | Andy McCammon, University of California, San Diego |
| ORAL | Karel Zimmerman, INRA, Jouy-en-Josas, France |
| ORIENT | Anthony Stone, Cambridge University |
| PCMODEL | Kevin Gilbert, Serena Software, Bloomington, Indiana |
| PEFF | Jan Dillen, University of Pretoria, South Africa |
| Q | Johan Åqvist, Uppsala University |
| SIBFA | Nohad Gresh, INSERM, CNRS, Paris |
| SIGMA | Jan Hermans, University of North Carolina |

SPASIBA
SPASMS
TINKER
XPLOR/CNS
YAMMP
YASP
YETI

Gerard Vergoten, Université de Lille
David Spellmeyer and the Kollman Group, UCSF
Jay Ponder, Washington University, St. Louis
Axel Brünger, Stanford University
Stephen Harvey, University of Alabama, Birmingham
Florian Mueller-Plathe, ETH Zentrum, Zurich
Angelo Vedani, Biografik-Labor 3R, Basel

AMBER D. A. Pearlman, D. A. Case, J. W. Caldwell, W. S. Ross, T. E. Cheatham III, S. DeBolt, D. Ferguson, G. Seibel and P. Kollman, AMBER, a Package of Computer Programs for Applying Molecular Mechanics, Normal Mode Analysis, Molecular Dynamics and Free Energy Calculations to Simulate the Structural and Energetic Properties of Molecules, *Comp. Phys. Commun.*, **91**, 1-41 (1995)

ARGOS T. P. Straatsma and J. A. McCammon, ARGOS, a Vectorized General Molecular Dynamics Program, *J. Comput. Chem.*, **11**, 943-951 (1990)

CHARMM B. R. Brooks, R. E. Bruccoleri, B. D. Olafson, D. J. States, S. Swaminathan and M. Karplus, CHARMM: A Program for Macromolecular Energy, Minimization, and Dynamics Calculations, *J. Comput. Chem.*, **4**, 187-217 (1983)

ENCAD M. Levitt, M. Hirshberg, R. Sharon and V. Daggett, Potential Energy Function and Parameters for Simulations for the Molecular Dynamics of Proteins and Nucleic Acids in Solution, *Comp. Phys. Commun.*, **91**, 215-231 (1995)

FANTOM T. Schaumann, W. Braun and K. Wurtrich, The Program FANTOM for Energy Refinement of Polypeptides and Proteins Using a Newton-Raphson Minimizer in Torsion Angle Space, *Biopolymers*, **29**, 679-694 (1990)

FEDER/2 H. Wako, S. Endo, K. Nagayama and N. Go, FEDER/2: Program for Static and Dynamic Conformational Energy Analysis of Macro-molecules in Dihedral Angle Space, *Comp. Phys. Commun.*, **91**, 233-251 (1995)

GROMACS H. J. C. Berendsen, D. van der Spoel and R. van Drunen, GROMACS: A Message-passing Parallel Molecular Dynamics Implementation, *Comp. Phys. Commun.*, **91**, 43-56 (1995)

GROMOS W. R. P. Scott, P. H. Hunenberger, I. G. Tironi, A. E. Mark, S. R. Billeter, J. Fennen, A. E. Torda, T. Huber, P. Kruger, W. F. van Gunsteren, The GROMOS Biomolecular Simulation Program Package, *J. Phys. Chem. A*, **103**, 3596-3607 (1999)

IMPACT D. B. Kitchen, F. Hirata, J. D. Westbrook, R. Levy, D. Kofke and M. Yarmush, Conserving Energy during Molecular Dynamics Simulations of Water, Proteins, and Proteins in Water, *J. Comput. Chem.*, **10**, 1169-1180 (1990)

MACROMODEL F. Mahamadi, N. G. J. Richards, W. C. Guida, R. Liskamp, M. Lipton, C. Caufield, G. Chang, T. Hendrickson and W. C. Still, MacroModel—An Integrated Software System for Modeling Organic and Bioorganic Molecules Using Molecular Mechanics, *J. Comput. Chem.*, **11**, 440-467 (1990)

MM2 N. L. Allinger, Conformational Analysis. 130. MM2. A Hydrocarbon Force Field Utilizing V_1 and V_2 Torsional Terms, *J. Am. Chem. Soc.*, **99**, 8127-8134 (1977)

MM3 N. L. Allinger, Y. H. Yuh and J.-H. Lii, Molecular Mechanics. The MM3 Force Field for Hydrocarbons, *J. Am. Chem. Soc.*, *111*, 8551-8566 (1989)

MM4 N. L. Allinger, K. Chen and J.-H. Lii, An Improved Force Field (MM4) for Saturated Hydrocarbons, *J. Comput. Chem.*, *17*, 642-668 (1996)

MMC C. E. Dykstra, Molecular Mechanics for Weakly Interacting Assemblies of Rare Gas Atoms and Small Molecules, *J. Am. Chem. Soc.*, *111*, 6168-6174 (1989)

MMFF T. A. Halgren, Merck Molecular Force Field. I. Basis, Form, Scope, Parameterization, and Performance of MMFF94, *J. Comput. Chem.*, *17*, 490-516 (1996)

MOIL R. Elber, A. Roitberg, C. Simmerling, R. Goldstein, H. Li, G. Verkhiver, C. Keasar, J. Zhang and A. Ulitsky, MOIL: A Program for Simulations of Macromolecules, *Comp. Phys. Commun.*, *91*, 159-189 (1995)

MOSCITO See the web site at <http://ganter.chemie.uni-dortmund.de/~pas/moscito.html>

NAMD L. Kalé, R. Skeel, M. Bhandarkar, R. Brunner, A. Gursoy, N. Krawetz, J. Phillips, A. Shinozaki, K. Varadarajan and K. Schulten, NAMD2: Greater Scalability for Parallel Molecular Dynamics, *J. Comput. Phys.*, *151*, 283-312 (1999)

OOMPAA G. A. Huber and J. A. McCammon, OOMPAA—Object-oriented Model for Probing Assemblages of Atoms, *J. Comput. Phys.*, *151*, 264-282 (1999)

ORAL K. Zimmermann, ORAL: All Purpose Molecular Mechanics Simulator and Energy Minimizer, *J. Comput. Chem.*, *12*, 310-319 (1991)

PCMODEL See the web site at <http://www.serenasoft.com>

PEFF J. L. M. Dillen, PEFF: A Program for the Development of Empirical Force Fields, *J. Comput. Chem.*, *13*, 257-267 (1992)

Q See the web site at <http://aqvist.bmc.uu.se/Q>

SIBFA N. Gresh, Inter- and Intramolecular Interactions. Inception and Refinements of the SIBFA, Molecular Mechanics (SMM) Procedure, a Separable, Polarizable Methodology Grounded on *ab Initio* SCF/MP2 Computations. Examples of Applications to Molecular Recognition Problems, *J. Chim. Phys. PCB*, *94*, 1365-1416 (1997)

SIGMA See the web site at <http://femto.med.unc.edu/SIGMA>

SPASIBA P. Derreumaux and G. Vergoten, A New Spectroscopic Molecular Mechanics Force-Field - Parameters For Proteins, *J. Chem. Phys.*, *102*, 8586-8605 (1995)

TINKER See the web site at <http://dasher.wustl.edu/tinker>

YAMMP R. K.-Z. Tan and S. C. Harvey, Yammp: Development of a Molecular Mechanics Program Using the Modular Programming Method, *J. Comput. Chem.*, *14*, 455-470 (1993)

YETI A. Vedani, YETI: An Interactive Molecular Mechanics Program for Small-Molecule Protein Complexes, *J. Comput. Chem.*, 9, 269-280 (1988)

MOLECULAR MECHANICS

U. Burkert and N. L. Allinger, **Molecular Mechanics**, American Chemical Society, Washington, D.C., 1982

K. Rasmussen, **Potential Energy Functions in Conformational Analysis** (Lecture Notes in Chemistry, Vol. 27), Springer-Verlag, Berlin, 1985

A. K. Rappé and C. J. Casewit, **Molecular Mechanics across Chemistry**, University Science Books, Sausalito, CA, 1997

K. Machida, **Principles of Molecular Mechanics**, Kodansha/John Wiley & Sons, Tokyo/New York, 1999

P. Comba and T. W. Hambley, **Molecular Modeling of Inorganic Compounds**, VCH, New York, 1995

COMPUTER SIMULATION METHODS

M. J. Field, **A Practical Introduction to the Simulation of Molecular Systems**, Cambridge Univ. Press, Cambridge, 1999

A. R. Leach, **Molecular Modelling: Principles and Applications**, Addison Wesley Longman, Essex, England, 1996

D. Frankel and B. Smit, **Understanding Molecular Simulation: From Algorithms to Applications**, Academic Press, San Diego, CA, 1996

D. C. Rapaport, **The Art of Molecular Dynamics Simulation**, Cambridge University Press, Cambridge, 1995

J. M. Haile, **Molecular Dynamics Simulation**, John Wiley and Sons, New York, 1992

M. P. Allen and D. J. Tildesley, **Computer Simulation of Liquids**, Oxford University Press, Oxford, 1987

T. Schlick, R. D. Skeel, A. T. Brünger, L. V. Kale, J. A. Board, J. Hermans and K. Schulten, Algorithmic Challenges in Computational Molecular Biophysics, *J. Comput. Phys.*, 151, 9-48 (1999)

MODELING OF BIOLOGICAL MACROMOLECULES

J. A. McCammon and S. Harvey, **Dynamics of Proteins and Nucleic Acids**, Cambridge University Press, Cambridge, 1987

C. L. Brooks III, M. Karplus and B. M. Pettitt, **Proteins: A Theoretical Perspective of Dynamics, Structure, and Thermodynamics**, John Wiley and Sons, New York, 1988

W. F. van Gunsteren, P. K. Weiner and A. J. Wilkinson, **Computer Simulation of Biomolecular Systems, Vol. 1-3**, Kluwer Academic Publishers, Dordrecht, 1989-1997

T. E. Cheatham and B. R. Brooks, Recent Advances in Molecular Dynamics Simulation towards the Realistic Representation of Biomolecules in Solution, *Theor. Chem. Acc.*, **99**, 279-288 (1998)

CONJUGATE GRADIENT AND QUASI-NEWTON OPTIMIZATION

J. Nocedal and S. J. Wright, **Numerical Optimization**, Springer-Verlag, New York, 1999

S. G. Nash and A. Sofer, **Linear and Nonlinear Programming**, McGraw-Hill, New York, 1996

R. Fletcher, **Practical Methods of Optimization**, John Wiley & Sons Ltd., Chichester, 1987

D. G. Luenberger, **Linear and Nonlinear Programming**, 2nd Ed., Addison-Wesley, Reading, MA, 1984

P. E. Gill, W. Murray and M. H. Wright, **Practical Optimization**, Academic Press, New York, 1981

J. Nocedal, Updating Quasi-Newton Matrices with Limited Storage, *Math. Comp.*, **773-782** (1980)

S. J. Watowich, E. S. Meyer, R. Hagstrom and R. Josephs, A Stable, Rapidly Converging Conjugate Gradient Method for Energy Minimization, *J. Comput. Chem.*, **9**, 650-661 (1988)

W. C. Davidon, Optimally Conditioned Optimization Algorithms without Line Searches, *Math. Prog.*, **9**, 1-30 (1975)

TRUNCATED NEWTON OPTIMIZATION

J. W. Ponder and F. M. Richards, An Efficient Newton-like Method for Molecular Mechanics Energy Minimization of Large Molecules, *J. Comput. Chem.*, **8**, 1016-1024 (1987)

R. S. Dembo and T. Steihaug, Truncated-Newton Algorithms for Large-Scale Unconstrained Optimization, *Math. Prog.*, **26**, 190-212 (1983)

S. C. Eisenstat and H. F. Walker, Choosing the Forcing Terms in an Inexact Newton Method, *SIAM J. Sci. Comput.*, **17**, 16-32 (1996)

T. Schlick and M. Overton, A Powerful Truncated Newton Method for Potential Energy Minimization, *J. Comput. Chem.*, **8**, 1025-1039 (1987)

D. S. Kershaw, The Incomplete Cholesky-Conjugate Gradient Method for the Iterative Solution of Systems of Linear Equations, *J. Comput. Phys.*, **26**, 43-65 (1978)

T. A. Manteuffel, An Incomplete Factorization Technique for Positive Definite Linear Systems, *Math. Comp.*, **34**, 473-497 (1980)

P. Derreumaux, G. Zhang and T. Schlick and B. R. Brooks, A Truncated Newton Minimizer Adapted for CHARMM and Biomolecular Applications, *J. Comput. Chem.*, **15**, 532-552 (1994)

I. S. Duff, A. M. Erisman and J. K. Reid, **Direct Methods for Sparse Matrices**, Oxford University Press, Oxford, 1986

POTENTIAL ENERGY SMOOTHING

R. V. Pappu, R. K. Hart and J. W. Ponder, Analysis and Application of Potential Energy Smoothing Methods for Global Optimization, *J. Phys. Chem. B*, **102**, 9725-9742 (1998)

L. Piela, J. Kostrowicki and H. A. Scheraga, The Multiple-Minima Problem in the Conformational Analysis of Molecules. Deformation of the Potential Energy Hypersurface by the Diffusion Equation Method, *J. Phys. Chem.*, **93**, 3339-3346 (1989)

J. Ma and J. E. Straub, Simulated Annealing Using the Classical Density Distribution, *J. Chem. Phys.*, **101**, 533-541 (1994)

C. Tsao and C. L. Brooks, Cluster Structure Determination Using Gaussian Density Distribution Global Minimization Methods, *J. Chem. Phys.*, **101**, 6405-6411 (1994)

S. Nakamura, H. Hirose, M. Ikeguchi and J. Doi, Conformational Energy Minimization Using a Two-Stage Method, *J. Phys. Chem.*, **99**, 8374-8378 (1995)

T. Huber, A. E. Torda and W. F. van Gunsteren, Structure Optimization Combining Soft-Core Interaction Functions, the Diffusion Equation Method, and Molecular Dynamics, *J. Phys. Chem. A*, **101**, 5926-5930 (1997)

S. Schelstraete and H. Verschelde, Finding Minimum-Energy Configurations of Lennard-Jones Clusters Using an Effective Potential, *J. Phys. Chem. A*, **101**, 310-315 (1998)

I. Andricioaei and J. E. Straub, Global Optimization Using Bad Derivatives: Derivative-Free Method for Molecular Energy Minimization, *J. Comput. Chem.*, **19**, 1445-1455 (1998)

L. Piela, Search for the Most Stable Structures on Potential Energy Surfaces, *Coll. Czech. Chem. Commun.*, **63**, 1368-1380 (1998)

“SNIFFER” GLOBAL OPTIMIZATION

A. O. Griewank, Generalized Descent for Global Optimization, *J. Opt. Theor. Appl.*, **34**, 11-39 (1981)

R. A. R. Butler and E. E. Slaminka, An Evaluation of the Sniffer Global Optimization Algorithm Using Standard Test Functions, *J. Comput. Phys.*, **99**, 28-32 (1993)

J. W. Rogers and R. A. Donnelly, Potential Transformation Methods for Large-Scale Global Optimization, *SIAM J. Optim.*, **5**, 871-891 (1995)

INTEGRATION METHODS FOR MOLECULAR DYNAMICS

D. Beeman, Some Multistep Methods for Use in Molecular Dynamics Calculations, *J. Comput. Phys.*, **20**, 130-139 (1976)

M. Levitt and H. Meirovitch, Integrating the Equations of Motion, *J. Mol. Biol.*, **168**, 617-620 (1983)

J. Aqvist, W. F. van Gunsteren, M. Leijonmarck and O. Tapia, A Molecular Dynamics Study of the C-Terminal Fragment of the L7/L12 Ribosomal Protein, *J. Mol. Biol.*, **183**, 461-477 (1985)

W. C. Swope, H. C. Andersen, P. H. Berens and K. R. Wilson, A Computer Simulation Method for the Calculation of Equilibrium Constants for the Formation of Physical Clusters of Molecules: Application to Small Water Clusters, *J. Chem. Phys.*, **76**, 637-649 (1982)

CONSTRAINT DYNAMICS

W. F. van Gunsteren and H. J. C. Berendsen, Algorithms for Macromolecular Dynamics and Constraint Dynamics, *Mol. Phys.*, **34**, 1311-1327 (1977)

G. Ciccotti, M. Ferrario and J.-P. Ryckaert, Molecular Dynamics of Rigid Systems in Cartesian Coordinates: A General Formulation, *Mol. Phys.*, **47**, 1253-1264 (1982)

H. C. Andersen, Rattle: A "Velocity" Version of the Shake Algorithm for Molecular Dynamics Calculations, *J. Comput. Phys.*, **52**, 24-34 (1983)

R. Kutteh, RATTLE Recipe for General Holonomic Constraints: Angle and Torsion Constraints, *CCP5 Newsletter*, **46**, 9-17 (1998) [available from the web site at http://www.dl.ac.uk/CCP/CCP5/newsletter_index.html]

B. J. Palmer, Direct Application of SHAKE to the Velocity Verlet Algorithm, *J. Comput. Phys.*, **104**, 470-472 (1993)

S. Miyamoto and P. A. Kollman, SETTLE: An Analytical Version of the SHAKE and RATTLE Algorithm for Rigid Water Models, *J. Comput. Chem.*, **13**, 952-962 (1992)

B. Hess, H. Bekker, H. J. C. Berendsen and J. G. E. M. Fraaije, LINCS: A Linear Constraint Solver for Molecular Simulations, *J. Comput. Chem.*, **18**, 1463-1472 (1997)

J. T. Slusher and P. T. Cummings, Non-Iterative Constraint Dynamics using Velocity-Explicit Verlet Methods, *Mol. Simul.*, **18**, 213-224 (1996)

LANGEVIN, BROWNIAN AND STOCHASTIC DYNAMICS

M. P. Allen, Brownian Dynamics Simulation of a Chemical Reaction in Solution, *Mol. Phys.*, **40**, 1073-1087 (1980)

W. F. van Gunsteren and H. J. C. Berendsen, Algorithms for Brownian Dynamics, *Mol. Phys.*, **45**, 637-647 (1982)

F. Guarnieri and W. C. Still, A Rapidly Convergent Simulation Method: Mixed Monte Carlo/Stochastic Dynamics, *J. Comput. Chem.*, **15**, 1302-1310 (1994)

M. G. Paterlini and D. M. Ferguson, Constant Temperature Simulations using the Langevin Equation with Velocity Verlet Integration, *Chem. Phys.*, **236**, 243-252 (1998)

CONSTANT TEMPERATURE AND PRESSURE DYNAMICS

H. J. C. Berendsen, J. P. M. Postma, W. F. van Gunsteren, A. DiNola and J. R. Haak, Molecular Dynamics with Coupling to an External Bath, *J. Chem. Phys.*, **81**, 3684-3690 (1984)

W. G. Hoover, Canonical Dynamics: Equilibrium Phase-space Distributions, *Phys. Rev. A*, **31**, 1695-1697 (1985)

J. J. Morales, S. Toxvaerd and L. F. Rull, Computer Simulation of a Phase Transition at Constant Temperature and Pressure, *Phys. Rev. A*, **34**, 1495-1498 (1986)

B. R. Brooks, Algorithms for Molecular Dynamics at Constant Temperature and Pressure, Internal Report of Division of Computer Research and Technology, National Institutes of Health, 1988.

M. Levitt, Molecular Dynamics of Native Protein: Computer Simulation of Trajectories, *J. Mol. Biol.*, **168**, 595-620 (1983)

OUT-OF-PLANE DEFORMATION TERMS

J. R. Maple, U. Dinar and A. T. Hagler, Derivation of Force Fields for Molecular Mechanics and Dynamics from *ab initio* Energy Surfaces, *Proc. Natl. Acad. Sci. USA*, **85**, 5350-5354 (1988)

S.-H. Lee, K. Palmo and S. Krimm, New Out-of-Plane Angle and Bond Angle Internal Coordinates and Related Potential Energy Functions for Molecular Mechanics and Dynamics Simulations, *J. Comput. Chem.*, **20**, 1067-1084 (1999)

ANALYTICAL DERIVATIVES OF POTENTIAL FUNCTIONS

K. J. Miller, R. J. Hinde and J. Anderson, First and Second Derivative Matrix Elements for the Stretching, Bending, and Torsional Energy, *J. Comput. Chem.*, **10**, 63-76 (1989)

D. H. Faber and C. Altona, UTAH5: A Versatile Programme Package for the Calculation of Molecular Properties by Force Field Methods, *Computers & Chemistry*, **1**, 203-213 (1977)

W. C. Swope and D. M. Ferguson, Alternative Expressions for Energies and Forces Due to Angle Bending and Torsional Energy, Report G320-3561, *J. Comput. Chem.*, **13**, 585-594 (1992)

A. Blondel and M. Karplus, New Formulation for Derivatives of Torsion Angles and Improper Torsion Angles in Molecular Mechanics: Elimination of Singularities, *J. Comput. Chem.*, **17**, 1132-1141 (1996)

R. E. Tuzun, D. W. Noid and B. G. Sumpter, Efficient Treatment of Out-of-Plane Bend and Improper Torsion Interactions in MM2, MM3, and MM4 Molecular Mechanics Calculations, *J. Comput. Chem.*, **18**, 1804-1811 (1997)

TORSIONAL SPACE DERIVATIVES AND NORMAL MODES

M. Levitt, C. Sander and P. S. Stern, Protein Normal-mode Dynamics: Trypsin Inhibitor, Crambin, Ribonuclease and Lysozyme, *J. Mol. Biol.*, **181**, 423-447 (1985)

M. Levitt, Protein Folding by Restrained Energy Minimization and Molecular Dynamics, *J. Mol. Biol.*, **170**, 723-764 (1983)

H. Wako and N. Go, Algorithm for Rapid Calculation of Hessian of Conformational Energy Function of Proteins by Supercomputer, *J. Comput. Chem.*, **8**, 625-635 (1987)

H. Abe, W. Braun, T. Noguti and N. Go, Rapid Calculation of First and Second Derivatives of Conformational Energy with Respect to Dihedral Angles for Proteins: General Recurrent Equations, *Computers & Chemistry*, **8**, 239-247 (1984)

T. Noguti and N. Go, A Method of Rapid Calculation of a Second Derivative Matrix of Conformational Energy for Large Molecules, *J. Phys. Soc. Japan*, **52**, 3685-3690 (1983)

ANALYTICAL SURFACE AREA AND VOLUME

M. L. Connolly, Analytical Molecular Surface Calculation, *J. Appl. Cryst.*, **16**, 548-558 (1983)

M. L. Connolly, Computation of Molecular Volume, *J. Am. Chem. Soc.*, **107**, 1118-1124 (1985)

M. L. Connolly, Molecular Surfaces: A Review, available from the web site at <http://www.netsci.org/Science/Compchem/feature14.html>

C. E. Kundrot, J. W. Ponder and F. M. Richards, Algorithms for Calculating Excluded Volume and Its Derivatives as a Function of Molecular Conformation and Their Use in Energy Minimization, *J. Comput. Chem.*, **12**, 402-409 (1991)

T. J. Richmond, Solvent Accessible Surface Area and Excluded Volume in Proteins, *J. Mol. Biol.*, **178**, 63-89 (1984)

L. Wesson and D. Eisenberg, Atomic Solvation Parameters Applied to Molecular Dynamics of Proteins in Solution, *Protein Science*, **1**, 227-235 (1992)

V. Gononea and E. Osawa, Implementation of Solvent Effect in Molecular Mechanics, Part 3. The First- and Second-order Analytical Derivatives of Excluded Volume, *J. Mol. Struct. (Theochem)*, **311** 305-324 (1994)

K. D. Gibson and H. A. Scheraga, Exact Calculation of the Volume and Surface Area of Fused Hard-sphere Molecules with Unequal Atomic Radii, *Mol. Phys.*, **62**, 1247-1265 (1987)

K. D. Gibson and H. A. Scheraga, Surface Area of the Intersection of Three Spheres with Unequal Radii: A Simplified Analytical Formula, *Mol. Phys.*, **64**, 641-644 (1988)

S. Sridharan, A. Nichols and K. A. Sharp, A Rapid Method for Calculating Derivatives of Solvent Accessible Surface Areas of Molecules, *J. Comput. Chem.*, **16**, 1038-1044 (1995)

APPROXIMATE SURFACE AREA AND VOLUME

S. J. Wodak and J. Janin, Analytical Approximation to the Accessible Surface Area of Proteins, *Proc. Natl. Acad. Sci. USA*, **77**, 1736-1740 (1980)

W. Hasel, T. F. Hendrickson and W. C. Still, A Rapid Approximation to the Solvent Accessible Surface Areas of Atoms, *Tetrahedron Comput. Method.*, **1**, 103-116 (1988)

J. Weiser, P. S. Shenkin and W. C. Still, Approximate Solvent-Accessible Surface Areas from Tetrahedrally Directed Neighbor Densities, *Biopolymers*, **50**, 373-380 (1999)

BOUNDARY CONDITIONS AND NEIGHBOR METHODS

W. F. van Gunsteren, H. J. C. Berendsen, F. Colonna, D. Perahia, J. P. Hollenberg and D. Lellouch, On Searching Neighbors in Computer Simulations of Macromolecular Systems, *J. Comput. Chem.*, **5**, 272-279 (1984)

F. Sullivan, R. D. Mountain and J. O'Connell, Molecular Dynamics on Vector Computers, *J. Comput. Phys.*, **61**, 138-153 (1985)

J. Boris, A Vectorized "Near Neighbors" Algorithm of Order N Using a Monotonic Logical Grid, *J. Comput. Phys.*, **66**, 1-20 (1986)

S. G. Lambrakos and J. P. Boris, Geometric Properties of the Monotonic Lagrangian Grid Algorithm for Near Neighbors Calculations, *J. Comput. Phys.*, **73**, 183-202 (1987)

T. A. Andrea, W. C. Swope and H. C. Andersen, The Role of Long Ranged Forces in Determining the Structure and Properties of Liquid Water, *J. Chem. Phys.*, **79**, 4576-4584 (1983)

D. N. Theodorou and U. W. Suter, Geometrical Considerations in Model Systems with Periodic Boundary Conditions, *J. Chem. Phys.*, **82**, 955-966 (1985)

J. Barnes and P. Hut, A Hierarchical O(NlogN) Force-calculation Algorithm, *Nature*, **234**, 446-449 (1986)

CUTOFF AND TRUNCATION METHODS

P. J. Steinbach and B. R. Brooks, New Spherical-Cutoff Methods for Long-Range Forces in Macromolecular Simulation, *J. Comput. Chem.*, **15**, 667-683 (1993)

R. J. Loncharich and B. R. Brooks, The Effects of Truncating Long-Range Forces on Protein Dynamics, *Proteins*, **6**, 32-45 (1989)

C. L. Brooks III, B. M. Pettitt and M. Karplus, Structural and Energetic Effects of Truncating Long Ranged Interactions in Ionic and Polar Fluids, *J. Chem. Phys.*, **83**, 5897-5908 (1985)

EWALD SUMMATION TECHNIQUES

A. Y. Toukmaji and J. A. Board, Jr., Ewald Summation Techniques in Perspective: A Survey, *Comp. Phys. Commun.*, **95**, 73-92 (1996)

T. Darden, L. Perera, L. Li and L. Pedersen, New Tricks for Modelers from the Crystallography Toolkit: The Particle Mesh Ewald Algorithm and its Use in Nucleic Acid Simulations, *Structure*, **7**, R550-R60 (1999)

T. Darden, D. York and L. G. Pedersen, Particle Mesh Ewald: An $N \cdot \log(N)$ Method for Ewald Sums in Large Systems, *J. Chem. Phys.*, **98**, 10089-10092 (1993)

U. Essmann, L. Perera, M. L. Berkowitz, T. Darden, H. Lee and L. G. Pedersen, A Smooth Particle Mesh Ewald Method, *J. Chem. Phys.*, **103**, 8577-8593 (1995)

W. Smith, Point Multipoles in the Ewald Summation (Revisited), *CCP5 Newsletter*, **46**, 18-30 (1998) [available from http://www.dl.ac.uk/CCP/CCP5/newsletter_index.html]

S. E. Feller, R. W. Pastor, A. Rojnuckarin, S. Bogusz and B. R. Brooks, Effect of Electrostatic Force Truncation on Interfacial and Transport Properties of Water, *J. Phys. Chem.*, **100**, 17011-17020 (1996)

W. Weber, P. H. Hünenberger and J. A. McCammon, Molecular Dynamics Simulations of a Polyalanine Octapeptide under Ewald Boundary Conditions: Influence of Artificial Periodicity on Peptide Conformation, *J. Phys. Chem. B*, **104**, 3668-3675 (2000)

CONJUGATED AND AROMATIC SYSTEMS

N. L. Allinger, F. Li, L. Yan and J. C. Tai, Molecular Mechanics (MM3) Calculations on Conjugated Hydrocarbons, *J. Comput. Chem.*, **11**, 868-895 (1990)

J. T. Sprague, J. C. Tai, Y. Yuh and N. L. Allinger, The MMP2 Computational Method, *J. Comput. Chem.*, **8**, 581-603 (1987)

J. Kao, A Molecular Orbital Based Molecular Mechanics Approach to Study Conjugated Hydrocarbons, *J. Am. Chem. Soc.*, **109**, 3818-3829 (1987)

J. Kao and N. L. Allinger, Conformational Analysis: Heats of Formation of Conjugated Hydrocarbons by the Force Field Method, *J. Am. Chem. Soc.*, **99**, 975-986 (1977)

D. H. Lo and M. A. Whitehead, Accurate Heats of Atomization and Accurate Bond Lengths: Benzenoid Hydrocarbons, *Can. J. Chem.*, **46**, 2027-2040 (1968)

G. D. Zeiss and M. A. Whitehead, Hetero-atomic Molecules: Semi-empirical Molecular Orbital Calculations and Prediction of Physical Properties, *J. Chem. Soc. A*, 1727-1738 (1971)

FREE ENERGY SIMULATION METHODS

P. Kollman, Free Energy Calculations: Applications to Chemical and Biochemical Phenomena, *Chem. Rev.*, **93**, 2395-2417 (1993)

B. L. Tembe and J. A. McCammon, Ligand-Receptor Interactions, *Computers & Chemistry*, **8**, 281-283 (1984)

W. L. Jorgensen and C. Ravimohan, Monte Carlo Simulation of Differences in Free Energy of Hydration, *J. Chem. Phys.*, **83**, 3050-3054 (1985)

W. L. Jorgensen, J. K. Buckner, S. Boudon and J. Tirado-Rives, Efficient Computation of Absolute Free Energies of Binding by Computer Simulations: Application to the Methane Dimer in Water, *J. Chem. Phys.*, **89**, 3742-3746 (1988)

S. H. Fleischman and C. L. Brooks III, Thermodynamics of Aqueous Solvation: Solution Properties of Alcohols and Alkanes, *J. Chem. Phys.*, **87**, 3029-3037 (1987)

U. C. Singh, F. K. Brown, P. A. Bash and P. A. Kollman, An Approach to the Application of Free Energy Perturbation Methods Using Molecular Dynamics, *J. Am. Chem. Soc.*, **109**, 1607-1614 (1987)

D. A. Pearlman and P. A. Kollman, A New Method for Carrying out Free Energy Perturbation Calculations: Dynamically Modified Windows, *J. Chem. Phys.*, **90**, 2460-2470 (1989)

T. P. Straatsma, H. J. C. Berendsen and J. P. M. Postma, Free Energy of Hydrophobic Hydration: A Molecular Dynamics Study of Noble Gases in Water, *J. Chem. Phys.*, **85**, 6720-6727 (1986)

T. P. Straatsma and H. J. C. Berendsen, Free Energy of Ionic Hydration: Analysis of a Thermodynamic Integration Technique to Evaluate Free Energy Differences by Molecular Dynamics Simulations, *J. Chem. Phys.*, **89**, 5876-5886 (1988)

M. Mezei, The Finite Difference Thermodynamic Integration, Tested on Calculating the Hydration Free Energy Difference between Acetone and Dimethylamine in Water, *J. Chem. Phys.*, **86**, 7084-7088 (1987)

A. E. Mark and W. F. van Gunsteren, Decomposition of the Free Energy of a System in Terms of Specific Interactions, *J. Mol. Biol.*, **240**, 167-176 (1994)

S. Boresch and M. Karplus, The Meaning of Copmponent Analysis: Decomposition of the Free Energy in Terms of Specific Interactions, *J. Mol. Biol.*, **254**, 801-807 (1995)

METHODS FOR PARAMETER DETERMINATION

N. L. Allinger, X. Zhou and J. Bergsma, Molecular Mechanics Parameters, *J. Mol. Struct. (THEOCHEM)*, **312**, 69-83 (1994)

A. J. Pertsin and A. I. Kitaigorodsky, **The Atom-Atom Potential Method: Application to Organic Molecular Solids**, Springer-Verlag, Berlin, 1987

D. E. Williams, Transferable Empirical Nonbonded Potential Functions, in **Crystal Cohesion and Conformational Energies**, Ed. by R. M. Metzger, Springer-Verlag, Berlin, 1981

A. T. Hagler and S. Lifson, A Procedure for Obtaining Energy Parameters from Crystal Packing, *Acta Cryst.*, **B30**, 1336-1341 (1974)

A. T. Hagler, S. Lifson and P. Dauber, Consistent Force Field Studies of Intermolecular Forces in Hydrogen-Bonded Crystals: A Benchmark for the Objective Comparison of Alternative Force Fields, **J. Am. Chem. Soc.**, *101*, 5122-5130 (1979)

W. L. Jorgensen, J. D. Madura and C. J. Swenson, Optimized Intermolecular Potential Functions for Liquid Hydrocarbons, **J. Am. Chem. Soc.**, *106*, 6638-6646 (1984)

W. L. Jorgensen and C. J. Swenson, Optimized Intermolecular Potential Functions for Amides and Peptides: Structure and Properties of Liquid Amides, **J. Am. Chem. Soc.**, *107*, 569-578 (1985)

J. R. Maple, U. Dinur and A. T. Hagler, Derivation of Force Fields for Molecular Mechanics and Dynamics from *ab Initio* Surfaces, **Proc. Nat. Acad. Sci. USA**, *85*, 5350-5354 (1988)

U. Dinur and A. T. Hagler, Direct Evaluation of Nonbonding Interactions from *ab Initio* Calculations, **J. Am. Chem. Soc.**, *111*, 5149-5151 (1989)

ELECTROSTATIC INTERACTIONS

S. L. Price, Towards More Accurate Model Intermolecular Potentials for Organic Molecules, **Rev. Comput. Chem.**, *14*, 225-289 (2000)

C. H. Faerman and S. L. Price, A Transferable Distributed Multipole Model for the Electrostatic Interactions of Peptides and Amides, **J. Am. Chem. Soc.**, *112*, 4915-4926 (1990)

C. E. Dykstra, Electrostatic Interaction Potentials in Molecular Force Fields, **Chem. Rev.**, *93*, 2339-2353 (1993)

M. J. Dudek and J. W. Ponder, Accurate Modeling of the Intramolecular Electrostatic Energy of Proteins, **J. Comput. Chem.**, *16*, 791-816 (1995)

U. Koch and E. Egert, An Improved Description of the Molecular Charge Density in Force Fields with Atomic Multipole Moments, **J. Comput. Chem.**, *16*, 937-944 (1995)

D. E. Williams, Representation of the Molecular Electrostatic Potential by Atomic Multipole and Bond Dipole Models, **J. Comput. Chem.**, *9*, 745-763 (1988)

F. Colonna, E. Evleth and J. G. Angyan, Critical Analysis of Electric Field Modeling: Formamide, **J. Comput. Chem.**, *13*, 1234-1245 (1992)

POLARIZATION EFFECTS

S. Kuwajima and A. Warshel, Incorporating Electric Polarizabilities in Water-Water Interaction Potentials, **J. Phys. Chem.**, *94*, 460-466 (1990)

J. W. Caldwell and P. A. Kollman, Structure and Properties of Neat Liquids Using Nonadditive Molecular Dynamics: Water, Methanol, and N-Methylacetamide, **J. Phys. Chem.**, *99*, 6208-6219 (1995)

D. N. Bernardo, Y. Ding, K. Kroegh-Jespersen and R. M. Levy, An Anisotropic Polarizable Water Model: Incorporation of All-Atom Polarizabilities into Molecular Mechanics Force Fields, *J. Phys. Chem.*, **98**, 4180-4187 (1994)

P. T. van Duijnen and M. Swart, Molecular and Atomic Polarizabilities: Thole's Model Revisited, *J. Phys. Chem. A*, **102**, 2399-2407 (1998)

K. J. Miller, Calculation of the Molecular Polarizability Tensor, *J. Am. Chem. Soc.*, **112**, 8543-8551 (1990)

J. Applequist, J. R. Carl and K.-K. Fung, An Atom Dipole Interaction Model for Molecular Polarizability. Application to Polyatomic Molecules and Determination of Atom Polarizabilities, *J. Am. Chem. Soc.*, **94**, 2952-2960 (1972)

J. Applequist, Atom Charge Transfer in Molecular Polarizabilities. Application of the Olson-Sundberg Model to Aliphatic and Aromatic Hydrocarbons, *J. Phys. Chem.*, **97**, 6016-6023 (1993)

A. J. Stone, Distributed Polarizabilities, *Mol. Phys.*, **56**, 1065-1082 (1985)

J. M. Stout and C. E. Dykstra, A Distributed Model of the Electrical Response of Organic Molecules, *J. Phys. Chem. A*, **102**, 1576-1582 (1998)

MACROSCOPIC TREATMENT OF SOLVENT

C. J. Cramer and D. G. Truhlar, Continuum Solvation Models: Classical and Quantum Mechanical Implementations, *Rev. Comput. Chem.*, **6**, 1-72 (1995)

B. Roux and T. Simonson, Implicit Solvation Models, *Biophys. Chem.*, **78**, 1-20 (1999)

M. K. Gilson, Introduction to Continuum Electrostatics with Molecular Applications, available from <http://gilsonlab.umbi.umd.edu>

SURFACE AREA-BASED SOLVATION MODELS

D. Eisenberg and A. D. McLachlan, Solvation Energy in Protein Folding and Binding, *Nature*, **319**, 199-203 (1986)

L. Wesson and D. Eisenberg, Atomic Solvation Parameters Applied to Molecular Dynamics of Proteins in Solution, *Prot. Sci.*, **1**, 227-235 (1992)

T. Ooi, M. Oobatake, G. Nemethy and H. A. Scheraga, Accessible Surface Areas as a Measure of the Thermodynamic Parameters of Hydration of Peptides, *Proc. Natl. Acad. Sci. USA*, **84**, 3086-3090 (1987)

J. D. Augspurger and H. A. Scheraga, An Efficient, Differentiable Hydration Potential for Peptides and Proteins, *J. Comput. Chem.*, **17**, 1549-1558 (1996)

GENERALIZED BORN SOLVATION MODELS

W. C. Still, A. Tempczyk, R. C. Hawley and T. Hendrickson, A Semiempirical Treatment of Solvation for Molecular Mechanics and Dynamics, *J. Am. Chem. Soc.*, **112**, 6127-6129 (1990)

D. Qiu, P. S. Shenkin, F. P. Hollinger and W. C. Still, The GB/SA Continuum Model for Solvation. A Fast Analytical Method for the Calculation of Approximate Born Radii, *J. Phys. Chem. A*, **101**, 3005-3014 (1997)

G. D. Hawkins, C. J. Cramer and D. G. Truhlar, Pairwise Solute Descreening of Solute Charges from a Dielectric Medium, *Chem. Phys. Lett.*, **246**, 122-129 (1995)

G. D. Hawkins, C. J. Cramer and D. G. Truhlar, Parametrized Models of Aqueous Free Energies of Solvation Based on Pairwise Descreening of Solute Atomic Charges from a Dielectric Medium, *J. Phys. Chem.*, **100**, 19824-19839 (1996)

A. Onufriev, D. Bashford and D. A. Case, Modification of the Generalized Born Model Suitable for Macromolecules, *J. Phys. Chem. B*, **104**, 3712-3720 (2000)

M. Schaefer and M. Karplus, A Comprehensive Analytical Treatment of Continuum Electrostatics, *J. Phys. Chem.*, **100**, 1578-1599 (1996)

M. Schaefer, C. Bartels and M. Karplus, Solution Conformations and Thermodynamics of Structured Peptides: Molecular Dynamics Simulation with an Implicit Solvation Model, *J. Mol. Biol.*, **284**, 835-848 (1998)

SUPERPOSITION OF COORDINATE SETS

S. J. Kearsley, An Algorithm for the Simultaneous Superposition of a Structural Series, *J. Comput. Chem.*, **11**, 1187-1192 (1990)

R. Diamond, A Note on the Rotational Superposition Problem, *Acta Cryst.*, **A44**, 211-216 (1988)

A. D. McLachlan, Rapid Comparison of Protein Structures, *Acta Cryst.*, **A38**, 871-873 (1982)

S. C. Nyburg, Some Uses of a Best Molecular Fit Routine, *Acta Cryst.*, **B30**, 251-253 (1974)

LOCATION OF TRANSITION STATES

R. Czerminski and R. Elber, Reaction Path Study of Conformational Transitions and Helix Formation in a Tetrapeptide, *Proc. Nat. Acad. Sci. USA*, **86**, 6963 (1989)

R. S. Berry, H. L. Davis and T. L. Beck, Finding Saddles on Multidimensional Potential Surfaces, *Chem. Phys. Lett.*, **147**, 13 (1988)

K. Muller, Reaction Paths on Multidimensional Energy Hypersurfaces, *Ang. Chem. Int. Ed. Engl.*, **19**, 1-13 (1980)

S. Bell and J. S. Crighton, Locating Transition States, *J. Chem. Phys.*, **80**, 2464-2475 (1984)

S. Fischer and M. Karplus, Conjugate Peak Refinement: An Algorithm for Finding Reaction Paths and Accurate Transition States in Systems with Many Degrees of Freedom, **Chem. Phys. Lett.**, 194, 252-261 (1992)

J. E. Sinclair and R. Fletcher, A New Method of Saddle-Point Location for the Calculation of Defect Migration Energies, **J. Phys. C**, 7, 864-870 (1974)

R. Elber and M. Karplus, A Method for Determining Reaction Paths in Large Molecules: Application to Myoglobin, **Chem. Phys. Lett.**, 139, 375-380 (1987)

D. T. Nguyen and D. A. Case, On Finding Stationary States on Large-Molecule Potential Energy Surfaces, **J. Phys. Chem.**, 89, 4020-4026 (1985)

T. A. Halgren and W. N. Lipscomb, The Synchronous-Transit Method for Determining Reaction Pathways and Locating Molecular Transition States, **Chem. Phys. Lett.**, 49, 225-232 (1977)

G. T. Barkema and N. Mousseau, Event-Based Relaxation of Continuous Disordered Systems, **Phys. Rev. Lett.**, 77, 4358-4361 (1996)